

Manuscript Number:	BOPH-D-17-00582R1
Full Title:	Chemoreduction combined with partial exenteration for orbital retinoblastoma
Article Type:	Research article
Abstract:	<p>Aim: To evaluate the role of chemotherapy combined with partial exenteration on orbital retinoblastoma patients.</p> <p>Methods: Six orbital retinoblastoma patients were enrolled in the study after ethics committee approval and written informed consent. They were treated by chemotherapy using carboplatin, vincristine and etoposide, then partial exenteration surgery (enucleating the eye and tumor mass in the orbit) was conducted according to the response of the lesion. All patients were asked to visit the doctor on time. The survival rate and growing development were evaluated.</p> <p>Results: Good tumor response was observed in all of patients. Among the 6 patients, 4 patients with no signs of progression received operations after 6 courses of chemotherapy and were still alive after 5 years follow-up. Unfortunately, one patient died of brain metastases after five courses of chemotherapy and the other patient died of abandoning treatment. Postoperative histological examination of the enucleation specimens showed largely fibrosis tissue and a small amount of calcification tissue in the orbit, while retinoblastoma in eyeball. Optic nerve resection margins were healthy in three cases. Only one patient received postoperative chemotherapy because of the involvement of microscopically optic nerve section.</p> <p>Conclusion: Chemoreduction combined with partial exenteration can be useful in orbital retinoblastoma.</p> <p>Trial registration: Registration number: chiCTR-ORC-17013231 ; registration time: 3 /11/2017. Retrospectively registered.</p>
Response to Reviewers:	<p>1. We note that your study assigned a treatment strategy to a set of participants to evaluate the outcome. Therefore, your study falls within the International Committee of Medical Journal Editors (ICMJE)' definition of a clinical trial: any research study that prospectively assigns human subjects to one or more health related interventions to evaluate the effects on health or biological outcomes. As such, Biomed Central requires that a Trial Registration Number (TRN) is provided in order for the manuscript to be published.</p> <p>Although a TRN is usually required prior to the start of the peer-review process, the BMC-series journals does accept retrospectively registered trials. If you have not registered the trial, we therefore request that you do so as soon as possible so that your study can be accepted for publication. All trials must be registered with an ICMJE approved registry, as listed in the ICMJE guide: http://www.icmje.org/about-icmje/faqs/</p> <p>Once you confirm that you will register this study as a clinical trial, we will be able to take note of this and not delay the review process any further.</p> <p>Once you know your trial registration number, please submit a revised version of your manuscript with the number and date of registration included in the abstract. The last section of the abstract should be Trial Registration: listing the trial registry and the unique identifying number, e.g. Trial registration: Current Controlled Trials ISRCTN73824458, as well as the date of registration. Please note that there should be no space between the letters and numbers of the trial registration number. If registration took place after the first participant was enrolled, please state also "Retrospectively registered" at the end of this section</p> <p>We have registrated the study in Chinese Clinical Trial Registry and the Trial Registration Number (TRN) is chiCTR-ORC-17013231.</p> <p>2. Please could you clarify whether your study was randomised or non-randomised.</p> <p>If your study was randomised, could you please ensure your manuscript reporting</p>

adheres to CONSORT guidelines (<http://www.consort-statement.org/>) for reporting clinical trials. Please include a statement within your manuscript to indicate that your study adheres to CONSORT guidelines and include a completed CONSORT checklist as an additional file when submitting your revised manuscript.

If your study was non-randomised, could you please ensure that your manuscript adheres to TREND guidelines (<https://www.cdc.gov/trendstatement/>) for reporting non-randomised trials. Please include a statement within your manuscript to indicate that your study adheres to TREND guidelines and include a completed TREND checklist as an additional file when submitting your revised manuscript.

This is so your methodology can be fully evaluated and utilised.

This study was non-randomised and adhered to Transparent Reporting of Evaluations with Nonrandomized Designs (TREND) guidelines .

3. The Availability of data and materials section refers to the raw data used in your study and presenting tables and figures is not sufficient to state that all data is contained within the manuscript and additional files. Please only use this statement if you have indeed provided all raw data on which your study is based. We strongly encourage all authors to share their raw data, either by providing it in a supplementary file or depositing it in a public repository and providing the details on how to access it in this section. If you do not wish to share your data, please clearly state this in this section along with a justification. Data availability statements can take one of the following forms (or a combination of more than one if required for multiple datasets):

- The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.
- All data generated or analysed during this study are included in this published article [and its supplementary information files].
- The datasets generated and/or analysed during the current study are not publicly available due [REASON WHY DATA ARE NOT PUBLIC] but are available from the corresponding author on reasonable request.
- Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.
- Not applicable. If your manuscript does not contain any data, please state 'Not applicable' in this section.

Please note that if you do wish to share your raw data and do not have consent from all patients to publish this data it will need to be de-identified.

Please also note that if you include your raw data as a supplementary file you will need to provide, after the References, a section titled “Additional files” where you list the following information about each of your supplementary files: * File name (e.g. Additional file 1), * Title of data, * Description of data. All additional files will also need to have been cited in the main manuscript.

The datasets used during the current study are available from the corresponding author on reasonable request.

4. Please represent authors' names using their full initials, not their full name, in the Authors' Contributions section. If there are any duplicated initials, please differentiate them to make it clear that the initials refer to separate authors
We have represent authors' names using their full initials in the Authors' Contributions section.

5. Please include the institutional addresses for all authors on the title page. The corresponding author should still be indicated.
We have included the institutional addresses for all authors on the title page.

6. We note that some abbreviations used in the manuscript text have not been included in the 'List of Abbreviations' section. Please ensure that all abbreviations

present in the text are listed in this section.
All abbreviations used in the manuscript text are listed in 'List of Abbreviations' section.

Please submit your revised manuscript by accessing the following site:

<http://boph.edmgr.com/>

If you have forgotten your username or password please use the "Send Login Details" link to get your login information. For security reasons, your password will be reset.

Please note, if your manuscript is accepted you will not be able to make any changes to the authors, or order of authors, of your manuscript once the editor has accepted your manuscript for publication. If you wish to make any changes to authorship before you resubmit your revisions, please reply to this email and ask for a 'Request for change in authorship' form which should be completed by all authors (including those to be removed) and returned to this email address. Please ensure that any changes in authorship fulfil the criteria for authorship as outlined in BioMed Central's editorial policies (<http://www.biomedcentral.com/about/editorialpolicies#authorship>).

Once you have completed and returned the form, your request will be considered and you will be advised whether the requested changes will be allowed.

By resubmitting your manuscript you confirm that all author details on the revised version are correct, that all authors have agreed to authorship and order of authorship for this manuscript and that all authors have the appropriate permissions and rights to the reported data.

Please be aware that we may investigate, or ask your institute to investigate, any unauthorised attempts to change authorship or discrepancies in authorship between the submitted and revised versions of your manuscript.

We look forward to receiving your revised manuscript before 06 Nov 2017.

Best wishes,

Amy Branch-Hollis
BMC Ophthalmology
<https://bmcophthalmol.biomedcentral.com/>

[Click here to view linked References](#)

Title page

Title:Chemoreduction combined with partial exenteration for orbital retinoblastoma

corresponding author:Jia Tan , Ophthalmology of Xiangya Hospital Central South University, 87 Xiangya Road ,Changsha, China, CN 410008 Tel:073189753021 ,
e-mail: jasmintj@126@.com

first author: Dan Liu, Ophthalmology of Xiangya Hospital Central South University, 87 Xiangya Road ,Changsha, China, CN 410008 ; e-mail: 184710173@qq.com

co-author: Xue-liang Xu, Ophthalmology of Xiangya Hospital Central South University, 87 Xiangya Road ,Changsha, China, CN 410008 ; e-mail:xuxueliang8341@163.com

co-author: Sha Wang, Ophthalmology of Xiangya Hospital Central South University, 87 Xiangya Road ,Changsha, China, CN 410008 e-mail:wangsha_1982@126.com

Abstract

Aim: To evaluate the role of chemotherapy combined with partial exenteration on orbital retinoblastoma patients.

Methods: Six orbital retinoblastoma patients were enrolled in the study after ethics committee approval and written informed consent. They were treated by chemotherapy using carboplatin, vincristine and etoposide, then partial exenteration surgery (enucleating the eye and tumor mass in the orbit) was conducted according to the response of the lesion. All patients were asked to visit the doctor on time. The survival rate and growing development were evaluated.

Results: Good tumor response was observed in all of patients. Among the 6 patients, 4 patients with no signs of progression received operations after 6 courses of chemotherapy and were still alive after 5 years follow-up. Unfortunately, one patient died of brain metastases after five courses of chemotherapy and the other patient died of abandoning treatment. Postoperative histological examination of the enucleation specimens showed largely fibrosis tissue and a small amount of calcification tissue in the orbit, while retinoblastoma in eyeball. Optic nerve resection margins were healthy in three cases. Only one patient received postoperative chemotherapy because of the involvement of microscopically optic nerve section.

Conclusion: Chemoreduction combined with partial exenteration can be useful in orbital retinoblastoma.

Trial registration: Registration number: chiCTR-ORC-17013231 ; registration time: 3 /11/2017. Retrospectively registered.

Retinoblastoma (RB) is the most frequent intraocular malignant tumor in children with a reported incidence ranging from 1 in 15,000 to 1 in 18,000 live births in the United States[1], while is higher in the developing countries, particularly in Asia and some Latin American countries[2]. In neglected or untreated RB patients, the tumor can break through the eye wall and becomes a huge mass occupying the orbit, that is called orbital RB and classified as stage \square RB according to the International Retinoblastoma Staging System (IRSS)[3]. The incidence of orbital RB is becoming rare in industrialized countries, ranging from 6.3% to 7.6% [4], but it is not an unusual feature in developing and under developed world [5][6]. Orbital RB is one of the major contributors to mortality and carries a poor prognosis for life [7][8]. The presence of

1 orbital invasion is associated with 10-27 times higher risk of metastasis when compared
2 to cases without orbital extension [9]. However, there is no proven definitive therapy
3 or management protocol for orbital RB. It continues to remain a challenging disease to
4 treat because of its complex nature. The traditional treatment is orbital exenteration
5 alone which is unlikely achieve surgical clearance and also leads to unrecoverable
6 damage of appearance. In recent years, systemic chemotherapy has been reported to
7 improve overall survival (OS) from less than 20% to 60% to 70% in patients with overt
8 orbital retinoblastoma [10][11][12] , but the RB cells may be still exist in the eyes and
9 the orbits. So chemotherapy combined with surgery are needed to achieve reasonable
10 results. In this study, we collected orbital RB patients in our hospital, and prospectively
11 investigated the outcome and compliance of the chemotherapy combined with partial
12 exenteration in orbital RB patients.
13
14
15
16
17
18
19
20
21
22
23

24 **Patients and methods**

25 Between July 2008 and August 2010, among 92 patients referred for RB to our
26 department, six (6.52%) presented with extensive RB with one case of bilateral orbital
27 mass and five cases with unilateral orbital mass, one of whom had ipsilateral parotid
28 gland mass. This population comprised five boys and one girl, aged 2 to 4 years. Three
29 received eyeball enucleation before, and three didn't receive any treatment. An initial
30 staging assessment was systematically performed at diagnosis (bone marrow
31 aspirate/bone marrow biopsy, lumbar puncture with cerebrospinal fluid cytology, and
32 axial magnetic resonance imaging), and the assessments were negative in all six
33 patients.
34
35
36
37
38
39
40
41

42 Initial treatment consisted of chemotherapy combining of carboplatin, vincristine and
43 etoposide(VEC), as described in the treatment of extraocular forms of RB [13]. All
44 patients received six courses of carboplatin-vincristine-etoposide (Table 1). A first
45 evaluation was performed after completion of chemotherapy. For all patients the
46 efficacy of chemotherapy was assessed by computed tomography (CT) and/or magnetic
47 resonance imaging (MRI).
48
49
50
51
52

53 **Table 1** Dose and usage of VEC chemotherapy
54
55
56
57
58
59
60
61
62
63
64
65

Drug	Daily dose		Period
	≤ 36months	> 36 months	
carboplatin	18.6mg/kg	560mg/m ²	D ₁
vincristine	0.05mg/kg	1.5 mg/m ²	D ₁
etoposide	5mg/kg	150 mg/m ²	D ₁ ,D ₂

Patients with no signs of progression were performed the partial exenteration six months after completion of the chemotherapy. However, the partial orbital exenteration in our study was different from traditional surgery. After the chemotherapy, the orbital mass was significantly smaller so that we can only enucleated the tumor mass in the orbit and tried to keep the patient's conjunctiva and other healthy orbital tissue for the secondary implantation. The intraoperative assessment determined the involvement of conjunctiva and the integrity of the zone of posterior section.

Postoperative treatment was determined by the results of surgery and histological examination of the surgical specimen. The indication for postoperative chemotherapy was based on the finding of the optic nerve sections and the histological results. Post-treatment follow up was based on clinical and radiological examinations.

To restore the patient's appearance, about 2 years after partial exenteration, the patients will receive secondary hydroxyapatite orbital implantation and wear artificial eye.

All patients were enrolled after written informed consent from their guardian, and the institute's ethics committee approved the study protocol (Ethics Committee of Xiangya Hospital).

This study was non-randomised and adhered to Transparent Reporting of Evaluations with Nonrandomized Designs (TREND) guidelines .

Results

The clinical courses of the patients were summarized in Table 2. The preoperative assessments showed clinical regression of all cases (Figure 1, 2). Four patients with no signs of progression received the operation. The intraoperative assessment showed that parts of the conjunctiva was involved, and calcified tissue connected to the eyeball. The orbital soft tissue around the eyeball and calcified tissue was fibrotic and adhesive to

the lesion. The optic nerve section was sufficiency posterior. Postoperative histological examination of the surgical specimens showed largely fibrosis tissue with a small amount of calcified component in the orbit, while RB tumor still existed in the eyeball. Optic nerve resection margins were healthy in three cases. The tumor invasion was visible in the optic nerve resection in one patient. This patient received additional three courses of chemotherapy after operation. Four patients received secondary hydroxyapatite orbital implantation and wore the prosthetic eyes about two years after the partial exenteration. The results were satisfactory in appearance (Figure 1, 2). These four patients were followed up for five years, and they all developed as well as others of the same age.

Unfortunately, one patient died of intracranial metastasis after five courses of chemotherapy, and one patient died of abandoning treatment. Of course, there are certain side effects of chemotherapy, such as hair loss, vomiting, diarrhea, granulocyte reduction which can be self-recovery after drug withdrawal.

Table 2 Clinical summary of the patients

Patients (age)	Clinicoradiological features	Neoadjuvant treatment	Evaluation	Exenteration histology	Postoperative treatment	Post-treatment response
1 (2.8 years)	Bilateral orbital mass	6 carboplatin-vincristine-etoposide	Clinical: regression; CT: tumor reduction	Retinoblastoma, No residual optic nerve invasion	None	22 months Survive
2 (2 years)	Unilateral obital mass	6 carboplatin-vincristine-etoposide	Clinical: regression; CT: tumor reduction	residual calcified tissues, no tumor	None	14 months Survive
3 (4.3 years)	Orbital portion of optic nerve involved and ipsilateral parotid gland mass	6 carboplatin-vincristine-etoposide	Clinical: regression; CT: tumor reduction CSF(-)	Retinoblastoma, Macroscopic invasion of optic nerve resection margins	3 VEC	16 months Survive
4 (3.8 years)	Unilateral obital mass	5 carboplatin-vincristine-etoposide	Clinical: regression; CT: tumor replase; brain metastasis	/	/	Died
5 (2.5 years)	Unilateral obital mass	6 carboplatin-vincristine-etoposide	Clinical: regression; CT: tumor reduction	Retinoblastoma, No residual optic nerve invasion	None	20 months Survive
6 (3.5 years)	Unilateral obital mass	1 carboplatin-vincristine-etoposide; then abandoned	Clinical: progression; CT: not done	/	/	Died

Discussion

1 For the orbital RB, traditional treatment method is the orbital exenteration
2 combined with external beam radiotherapy, which can lead to significant appearance
3 damaging and high risk of life-threatening. The secondary plastic surgery is extremely
4 difficult due to significant soft tissue atrophy of the facial region. Both physical and
5 mental developments of children are seriously affected. How to treat the orbital RB
6 effectively, meanwhile, to improve the survival rate and quality of life of patients, is
7 becoming an important issue of the ophthalmologists in the developing countries.
8

9
10
11
12
13 Chemotherapy used in RB patients began in the 1990s, and now it is becoming a
14 first-line treatment for intraocular RB. Chemotherapies for intraocular phase of RB
15 currently include of vincristine, etoposide and carboplatin, those dose are different
16 among the research institutions. The chemotherapy we used in the study is referring to
17 Oncology Department, Children's Hospital of Philadelphia in U.S [13]. Our results
18 showed the orbital tumors were significantly reduced to near extinction after VEC
19 chemotherapy in all 6 patients (7 eyes, 100%), four of the six children received partial
20 exenteration instead of exenteration and remained alive after five-year follow-up.
21 However, the drugs can't pass through the blood-brain barrier very well, so the effect
22 on central neuron system metastases is poor. One patient in our hospital died of brain
23 metastases after five courses of chemotherapy, and was presumed to have brain
24 metastases before the last chemotherapy. Although we didn't detect any cancer cell in
25 this patients' cerebrospinal fluid before chemotherapy, but it may be false negative
26 result. Therefore, we recommended multiple examinations of cerebrospinal fluid for
27 orbital RB patients. And unfortunately, one patient in our study died because his
28 parents gave up the treatment.
29

30
31
32
33
34 Although the orbital tumor of some children could be significantly reduced after
35 chemotherapy, there still existed the residual tumor in the eye[12]. It was also
36 certified by pathological examination in the study. So further surgery was needed to
37 eradicate the tumor. Radhakrishnan et al. [12]recommended that only enucleation was
38 needed after neoadjuvant chemotherapy. In our opinion, we recommended partial
39 orbital exenteration was a better way since the tumor may not only exist in the eyeball
40 but also in the orbit. On the consideration of the secondary plastic surgery, the partial
41 orbital enucleation performed in the study was different from traditional surgery. We
42 kept the patient's conjunctiva and other healthy orbital tissue maximally, which could
43 create favorable conditions for the secondary plastic surgery.
44

45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60 However, the treatment of RB does not end with enucleation or exenteration.
61
62
63
64
65

Prosthetic rehabilitation of the patients is of the equal importance. Improving appearance will help them to reintegrate in today's esthetic conscious society, thereby improving their quality of life. Previous studies about orbital RB were only concerned about the survival rate, little was reported whether patients accepted cosmetic surgeries and restored better appearance. In our study, about 2 years after partial exenteration, four patients received secondary hydroxyapatite orbital implantation and wore artificial eyes. The facial appearances are improved by the above operations. Therefore, Chemoreduction combined with partial exenteration may be a good method for orbital Rb patients. It can not only save the RB patients life, but also improve these patients appearance and quality of life.

The main side effect of chemotherapy is ototoxicity or renal toxicity after the application of carboplatin. Smits and his colleagues tested 25 RB children's hearing receiving carboplatin chemotherapy, and found no abnormalities during follow-up of 25 months[14]. The similar results are also observed in China. Patients received VEC chemotherapy have normal hearing and renal function, while they might have mild bone marrow suppression, but no serious infections. There are also reports the application of etoposide may result in the secondary tumors[15], especially in children who are hereditary RB. In the 5 years follow-up, we have not yet found the occurrence of secondary tumors in children.

Since orbital RB is a rare disease, and will occur more rarely because of the widely use of chemotherapy on intraocular RB, we could not obtain the enough sample to reach to the scientific conclusion about the survival rate of orbital RB patients and effectiveness of the treatment reported in this study. It is the limitation of the study. But we think it is still of importance to similar case, and further observation is needed to evaluate the survival condition and life quality of the patients.

List of abbreviations

RB: retinoblastoma; VEC: carboplatin, vincristine and etoposide

Declarations:

Acknowledgments

We are grateful to the patients and their families for participating in this study. We also thank Doctor Zhu for the help of writing this manuscript.

Funding

No source of funding was received for this research.

Availability of data and materials

The datasets used during the current study available from the corresponding author on reasonable request.

Authors' contributions:

Professor Tan , Professor Xu and Doctor Liu contributed significantly to the patients' treatment and follow-up; Professor Tan, Doctor Liu , Doctor Wang contributed significantly to the acquisition of data , data analysis and manuscript writing. All the authors have read and approved of the final version of the manuscript.

Ethics approval and consent to participate

The study was conducted in compliance with informed consent regulations and the Declaration of Helsinki. The study protocol was approved by Xiangya Hospital Ethics Committee. Informed consent was obtained from the parents of the patients for the study.

Consent for publication

Consent for publish was obtained from the patients'parents in this study.

Competing interests

The authors declare that they have no competing interests.

References

- [1] Tamboli A, Podgor MJ, Horm JW. The incidence of retinoblastoma in the United States: 1974 through 1985. Arch Ophthalmol. 1990;108(1):128-132.
- [2] Leal-Leal C, Flores-Rojas M, Medina-Sanson A, Cerecedo-Diaz F, Sanchez-Felix S, Gonzalez-Ramella O, Perez-Perez F, Gomez-Martinez R, Quero-Hernandez A, Altamirano-Alvarez E, Alejo-Gonzalez F, Figueroa-Carbajal J, Ellis-Irigoven A, Tejocote-Romero I, Cervantes-Paz R, Pantoja-Guillen F, Vega-Vega L, Carrete-Ramirez F. A multicentre report from the Mexican Retinoblastoma Group. Br J Ophthalmol. 2004;88(8):1074-1077.
- [3] Chantada G, Doz F, Antoneli CB, Grundy R, Clare Stannard FF, Dunkel IJ, Grabowski E, Leal-Leal C, Rodríguez-Galindo C, Schwartzman E, Popovic MB, Kremens B, Meadows AT, Zucker JM. A proposal for an international retinoblastoma staging system. Pediatr Blood Cancer. 2006 Nov;47(6):801-5.

- [4]. Grabowski EF, Abramson DH. Intraocular and extra ocular retinoblastoma. *Hematol Oncol Clin North Am.*1987;1(4):721–35. [PubMed]]
- [5]Ali MJ, Reddy VA, Honavar SG, Naik M. Orbital retinoblastoma: where do we go from here? *J Cancer Res Ther*, 2011, 7(1):11-14.
- [6] Chantada GL, Qaddoumi I, Canturk S, Khetan V, Ma Z, Kimani K, Yeniad B, Sultan I, Sitorus RS, Tacyildiz N, Abramson DH. Strategies to manage retinoblastoma in developing countries. *Pediatr Blood Cancer*, 2011, 56(3):341-348.
- [7]. Abramson DH, Beaverson K, Sangani P. Screening for retinoblastoma: presenting signs as prognosticators of patient and ocular survival. *Paediatrics*. 2003;112(6):1248–55.
- [8] Swathi K, Carol L. Retinoblastoma: achieving new standards with methods of chemotherapy. *Indian J Ophthalmol*. 2015;63(2):103–9.
- [9]Singh AD, Shields CL, Shields JA. Prognostic factors in retinoblastoma. *J Pediatr Ophthalmol Strabismus* 2000;37:134-41. †
- [10]Antoneli CB1, Ribeiro KB, Rodriguez-Galindo C, Soares FA, Arias VA, Novaes PE, Chojniak MM, Malogolowkin M. The addition of ifosfamide/etoposide to cisplatin/teniposide improves the survival of children with retinoblastoma and orbital involvement.[*J Pediatr Hematol Oncol*. 2007 Oct;29(10):700-4.
- [11]Chantada G, Fandiño A, Casak S, Manzitti J, Raslawski E, Schvartzman E. Treatment of overt extraocular retinoblastoma.*Med Pediatr Oncol*. 2003 Mar;40(3):158-61.
- [12]Radhakrishnan V, Kashyap S, Pushker N, Sharma S, Pathy S, Mohanti BK, Vishnubhatla S, Ghose S, Bakhshi S.Outcome, pathologic findings, and compliance in orbital retinoblastoma (InternationalRetinoblastoma Staging System stage III) treated with neoadjuvant chemotherapy: a prospective study.*Ophthalmology*. **2012** Jul;119(7):1470-7.
- [13] Friedman DL, Himelstein B, Shields CL, Shields JA, Needle M, Miller D, Bunin GR, Meadow AT. Chemoreduction and local ophthalmic therapy for intraocular retinoblastoma. *J Clin Oncol*. 2000,18(1):12-17.

[14] Smits C, Swen SJ, Theo Goverts S, Moll AC, Imhof SM, Schouten-van Meeteren AY. Assessment of hearing in very young children receiving carboplatin for retinoblastoma. Eur J Cancer. 2006;42(4):492-500.

[15] Ezoe S. Secondary leukemia associated with the anti-cancer agent, etoposide, a topoisomerase II inhibitor. Int J Environ Res Public-Health. 2012;9(7):2444-2453.

Legends:

Fig 1: The appearance and CT performance of a patient before, after chemotherapy and after artificial eyes wearing

A1-A5: before chemotherapy ;B1-B5:after chemotherapy;C1-C5:after artificial eyes wearing. A2, B2 and C2: CT scanning with axial position; A3, B3 and C3: CT scanning with coronal position; A4; B4 and C4:CT scanning with sagittal position of right eye; A5, B5 and C5: CT scanning with sagittal position of left eye.

Fig 2: The appearance of a patient before and after complete treatment.

A: before chemotherapy; B: after chemotherapy; C: after partial exenteration ; D: after secondary hydroxyapatite orbital implantation and artificial eye wearing.

Figure 1




Figure 2

[Click here to download Figure fig.2.tif](#)

