

Review Article

Orbital retinoblastoma: Where do we go from here?

ABSTRACT

Diagnosis of orbital retinoblastoma traditionally carries a dismal prognosis. Although its incidence is less in the developed countries, it continues to contribute to an epidemic of extraocular disease at diagnosis in the developing world. Orbital retinoblastoma encompasses a wide range of distinct clinical entities with varying tumor load. There are no standard treatment protocols as of now but the current preferred management is multimodal with a combination of initial high-dose chemotherapy, surgery, external beam radiotherapy and prolonged chemotherapy for 12 cycles. Though orbital retinoblastoma is a catastrophic event, rapid advances on many fronts, especially the genetic, makes the future appear brighter than what it is now. This review looks at all the new frontiers that are in store in the near as well as the distant future. Looking at the ever expanding horizons makes one believe of a definite hope that one day we will conquer this disease as we have conquered many others in the past.

KEY WORDS: Extraocular, future, orbit, RB1 gene, retinoblastoma

INTRODUCTION

Retinoblastoma is the most common intraocular malignancy in children, with a reported incidence ranging from 1 in 15,000 to 1 in 18,000 live births.^[1] It is bilateral in about 25–35% of cases.^[2] The average age at diagnosis is 18 months, unilateral cases being diagnosed at around 24 months and bilateral cases before 12 months.^[2] Retinoblastoma was associated with near certain death just over a century ago. Early tumor recognition aided by indirect ophthalmoscopy and refined enucleation technique contributed to an improved survival from 5% in 1896 to 81% in 1967. Advances in external beam radiotherapy in the 1970s, followed by the era of chemoreduction in the late 1990s, resulted in further substantial eye salvage.^[2-5]

WHERE DO WE STAND NOW?

In neglected or untreated cases, retinoblastoma can demonstrate extraocular spread primarily through optic nerve^[6] and also through the sclera.^[7] Though it is a rare clinical presentation in developed countries, ranging from 6.3 to 7.6%,^[8,9] it is not an unusual feature in developing and underdeveloped world. Leal-Leal *et al.*^[10] reported an incidence of 18% in a large multicenter study from Mexico. Kao *et al.*^[11] from Taiwan reported the incidence of orbital retinoblastoma to be 36% in a large study. The incidence is even higher (around 40%) from Nepal where Badhu *et al.*^[12] reported proptosis

to be the most common presenting feature of retinoblastoma.

Orbital retinoblastoma is one of the major contributors to mortality and carries a poor prognosis for life.^[13-17] The presence of orbital invasion is associated with 10–27 times higher risk of metastasis when compared to cases without orbital extension.^[18-20] The 5-year survival rates of orbital retinoblastoma has been reported to be 88% from the United Kingdom,^[21] 91% from Japan^[22] and 93% from the United States.^[23,24] However, the mortality in developing countries is still high owing to late presentations compounded by socioeconomic factors, with the mortality reported as high as 50–90%.^[12,17,25-27]

There is no proven definitive therapy or management protocol for orbital retinoblastoma. It continues to remain a challenging disease to treat because of its complex nature and usually various combination therapies are needed to achieve reasonable results. Multimodal therapies are probably the next step forward. There are numerous reasons for this, which are as follows:

- Systemic chemotherapy alone is unlikely to eradicate residual orbital disease.^[28,29]
- Orbital exenteration alone is unlikely to achieve surgical clearance.^[30,31]
- External beam radiotherapy is unlikely to prevent systemic metastasis.^[32,33]
- Histopathologic evidence of viable tumor cells present even in phthisical eyes following neoadjuvant chemotherapy.^[34]

Mohammad Javed Ali, Vijay Anand P. Reddy¹, Santosh G. Honavar, Milind Naik

Ocular Oncology Service, L.V. Prasad Eye Institute, Road No. 2, Banjara Hills, ¹Apollo Cancer Hospital, Jubilee Hills, Hyderabad – 500 034, India

For correspondence:

Dr. Santosh G. Honavar, Ocular Oncology Service, L.V. Prasad Eye Institute, Road No. 2, Banjara Hills, Hyderabad – 500 034, India.
E-mail: honavar@lvpai.org

Access this article online

Website: www.cancerjournal.net

DOI: 10.4103/0973-1482.80429

PMID: 21546735

Quick Response Code:



Based on the current evidence, as mentioned above, Honavar *et al.*^[35] developed a treatment protocol comprising initial triple drug high dose chemotherapy (3–6 cycles) followed by appropriate surgery, orbital radiotherapy and an additional 12-cycle standard dose chemotherapy. In this series, six cases of orbital retinoblastoma without intracranial extension and systemic metastasis underwent the protocol as described above and the authors reported dramatic resolution of orbital involvement and a mean event-free survival of 36 months. Most of the eyes following chemotherapy become phthisical. The authors agree that their encouraging protocol needs validation and further studies are needed to know whether fewer cycles are as equally effective since there are concerns of the long-term effects of high dose chemotherapy.

NEW FRONTIERS OF THE FUTURE

When reviewing the literature, one fact that stands out clearly is the increasing survival of patients with orbital retinoblastoma. Progress has been made and is continuing on all fronts including medical, surgical, diagnostic, genetic, social and rehabilitative.^[36,37]

Medical frontier

Multimodal therapies for advanced retinoblastoma are picking up pace and support from all spheres as amply elucidated in this review. Histopathologic evaluations of eyes following neoadjuvant chemotherapy for orbital retinoblastoma have further vouched for multimodal approaches.^[34] The introduction of stem cell rescue along with high dose chemotherapy has added another dimension to the treatment of orbital retinoblastoma.^[38,39] The Children's Oncology Group trials (COG trials), currently taking place, has made a great effort to standardize the treatment protocols worldwide. Their well-designed COG ARET 0321 trial of intensive multimodal therapy for extraocular retinoblastoma will probably lay to rest most of the confusion revolving around management protocols. Other areas being investigated include pharmacologic enhancement of radiotherapy, use of tumor cell targeting techniques, differentiating agents and immunotherapy.^[40]

Radiotherapy frontier

Advances in external beam radiotherapy for retinoblastoma with more precise control of the beam through better collimation and tighter isodose curves strongly argue in favor of its continuing supportive role in the management. The modern approaches that are being investigated include stereotactic conformal radiotherapy using a micromultileaf collimator, proton therapy using a fixed horizontal beam and tantalum localization or a rotating gantry with spot scanning.^[41]

Surgical frontier

Better survival has led to increasing research and advances on the surgical front. Newer implants with focus on orbital development and, hence, better cosmesis are coming up.^[42]

Increasing developments in ophthalmology with focus on newer materials and techniques are leading to better implant retention and less exposures or extrusions.^[43] Upcoming concepts in orbital reconstruction, particularly in children, and newer modalities like free tissue transfer for anophthalmic orbit syndrome may pave way for good cosmesis in patients of orbital retinoblastoma who undergo exenteration and radiotherapy.^[44,45]

Diagnostic frontier

On the diagnostic front, exploring the fetal eye is a new frontier. Fetal magnetic resonance imaging (MRI) and fetal three-dimensional ultrasound are being increasingly explored for prenatal diagnosis. The only two cases reported of *in utero* diagnosis using fetal ultrasound had massive extraocular extension.^[46,47] MRI is also being increasingly used to study the biocolonization of the orbital implant.^[48] Whole body bone scans using technetium-99, as shown by Kiratli *et al.*,^[49] and fluorine-18 fluorodeoxyglucose positron emission tomography (PET CT), as demonstrated by Moll *et al.*,^[50] reflect a glimpse of what possibly lies in store for early detection of metastasis.

Genetic frontier

On the genetic front, there has been a distressingly little progress in spite of the fact that *RB1* was the first human cancer gene to be cloned. Development of an automated, inexpensive screening examination for *RB1* mutations has been a long-term need. However, recently, Parsam *et al.*,^[51] from the authors group, have developed and published a combinatorial and less expensive approach for the detection of *RB1* mutations, which is likely to have applications as a screening tool. Ali *et al.*^[52] have for the first time explored the possible correlations between different types of mutations on the *RB1* gene and clinical presentations. It is interesting to note that large deletions were found to correlate with extraocular extension and metastasis at presentation. Further efforts are needed to make it a routine part of patient care. There is also an increasing realization of using a different approach for patients at high risk of metastatic disease. Newer researchers have described a class of genes called the metastasis genes and the metastasis suppressor genes.^[53-57] These genes affect the ability of cancer cells to establish growth foci in locations distant from the primary cancer but do not affect the primary tumor itself. The exploitation of these genes and the pathways to block the growth at distant sites holds promise for precisely targeted therapy to prevent metastasis.^[58-60]

Social frontier

On the social front, awareness through education and outreach to the community has helped to prevent delayed presentations and promote an early referral. Impact of educational programs in certain developing nations of Central America where it was linked with the vaccination programs is continuing to yield encouraging results with the rate of orbital retinoblastoma diagnosis reducing by almost half in the post-campaign period.^[61,62] Similar programs could be cloned to developing

countries in Africa and southern Asia, specifically targeting the education of general population and primary care providers.

CONCLUSIONS

Orbital retinoblastoma is no doubt still a huge challenge by itself. Although the survival has increased over the last few years, lack of access to medical facilities, lack of education about the need for early medical attention and cultural resistance to enucleation continue to contribute to an epidemic of extraocular disease at diagnosis in the developing world. Multimodal therapies and advancement on all fronts elucidated above appear as rays of light in a dark forest which is providing us with clues to a glittering light ahead to fight the darkness surrounding us at present. There is a definite hope that one day we will conquer this disease as we have conquered many others in the past.

REFERENCES

- Bishop JO, Madsen EC. Retinoblastoma: Review of current status. *Surv Ophthalmol* 1975;19:342-66.
- Shields JA, Shields CL. Retinoblastoma. In: Shields CL, Shields JA, editors. *Intraocular tumors – A text and Atlas*. 2nd ed. Philadelphia, United States: Lippincott Williams and Wilkins; 2007. p. 294-318.
- Kingston JE, Hungerford JL, Madreperla SA, Plowman PN. Results of combined chemotherapy and radiotherapy for advanced intraocular retinoblastoma. *Arch Ophthalmol* 1996;114:1339-43.
- Gallie BL, Budning A, DeBoer G, Thiessen JJ, Koren G, Verjee Z, *et al.* Chemotherapy with focal therapy can cure retinoblastoma without radiotherapy. *Arch Ophthalmol* 1996;114:1321-8.
- Shields CL, De Potter P, Himelstein BP, Shields JA, Meadows AT, Maris JM. Chemoreduction in the initial management of intraocular retinoblastoma. *Arch Ophthalmol* 1996;114:1330-8.
- Magrann I, Abramson DH, Ellsworth RM. Optic nerve involvement in retinoblastoma. *Ophthalmology* 1989;96:217-22.
- Rootman J, Ellsworth RM, Hofbauer J, Kitchen D. Orbital extension of retinoblastoma: A clinicopathological study. *Can J Ophthalmol* 1978;13:72-80.
- Grabowski EF, Abramson DH. Intraocular and extra ocular retinoblastoma. *Hematol Oncol Clin North Am* 1987;1:721-35.
- Ellsworth RM. Orbital Retinoblastoma. *Trans Am Ophthalmol Soc* 1974;72:79-88.
- Leal-Leal C, Flores-Rojas M, Medina-Sanson A, Cerecedo-Diaz F, Sanchez-Felix S, Gonzalez-Ramella O, *et al.* A multi-centre report from the Mexican retinoblastoma group. *Br J Ophthalmol* 2004;88:1074-7.
- Kao LY, Su WW, Lin YW. Retinoblastoma in Taiwan: survival and clinical characteristics 1978-2000. *Jpn J Ophthalmol* 2002;46:577-80.
- Badhu B, Sah SP, Thakur SK, Dulal S, Kumar S, Sood A, *et al.* Clinical presentation of retinoblastoma in Eastern Nepal. *Clin Exp Ophthalmol* 2005;33:386-9.
- Stannard C, Lipper S, Sealy R, Sevel D. Retinoblastoma: Correlation of invasion of the optic nerve and choroids with prognosis and metastases. *Br J Ophthalmol* 1979;63:560-70.
- Hungerford J. Factors influencing metastasis in retinoblastoma. *Br J Ophthalmol* 1993;77:541.
- Finger PT, Harbour JW, Karciglu ZA. Risk factors for metastasis in retinoblastoma. *Surv Ophthalmol* 2002;47:1-16.
- Abramson DH, Beaverson K, Sangani P, Vora RA, Lee TC, Hochberg HM, *et al.* Screening for retinoblastoma: Presenting signs as prognosticators of patient and ocular survival. *Paediatrics* 2003;112:1248-55.
- Schvartzman E, Chantada G, Fandino A, de Davila MT, Raslawski E, Manzitti J. Results of a stage-based approach for the treatment of retinoblastoma. *J Clin Oncol* 1996;14:1532-6.
- Singh AD, Shields CL, Shields JA. Prognostic factors in retinoblastoma. *J Pediatr Ophthalmol Strabismus* 2000;37:134-41.
- Kopelman JE, McLean IW, Rosenberg SH. Multivariate analysis of risk factors for metastasis in retinoblastoma treated by enucleation. *Ophthalmology* 1987;94:371-7.
- Khelfaoui F, Validire P, Auperin A, Quintana E, Michon J, Pacquement H, *et al.* Histopathologic risk factors in retinoblastoma: a retrospective study of 172 patients treated in a single institution. *Cancer* 1996;77:1206-13.
- Sanders BM, Draper GJ, Kingston JE. Retinoblastoma in Great Britain 1969-80: Incidence, treatment and survival. *Br J Ophthalmol* 1988;72:576-83.
- Survival rate and risk factors for patients with retinoblastoma in Japan. The committee for the National Registry of Retinoblastoma. *Jpn J Ophthalmol* 1992;36:121-31.
- Abramson DH, Niksarli K, Ellsworth RM, Servodidio CA. Changing trends in the management of retinoblastoma: 1951-1965 vs 1966-1980. *J Pediatr Ophthalmol Strabismus* 1994;31:32-7.
- Young J, Smith MA, Roffers SD. Retinoblastoma. In: Ries LA, Smith MA, Gurney JG, editors. *Cancer incidence and survival among children and adolescents*. SEER Program. United States: NIH Pub; 1999. p. 99-4649.
- Ajaiyeoba IA, Akang EE, Campbell OB, Olurin IO, Aghadiuno PU. Retinoblastomas in Ibadan: Treatment and prognosis. *West Africa J Med* 1993;12:223-7.
- Chantada G, Fandino A, Manzitti J, Urrutia L, Schvartzman E. Late diagnosis of retinoblastomas in a developing country. *Arch Dis Child* 1999;80:171-4.
- Kodilinye HC. Retinoblastoma in Nigeria: Problems of treatment. *Am J Ophthalmol* 1967;63:469-81.
- Antoneli CB, Steinhof F, de Cassia Braga RK, Novaes PE, Chojniak MM, Arias V, *et al.* Extra ocular retinoblastoma: A 13-year experience. *Cancer* 2003;98:1292-8.
- Doz F, Khelfaoui F, Mosseri V, Validire P, Quintana E, Michon J, *et al.* The role of chemotherapy in orbital involvement of retinoblastoma. The experience of single institution with 33 patients. *Cancer* 1994;74:722-32.
- Hungerford JL, Kingston JE, Plowman PN. Orbital recurrence of retinoblastoma. *Ophthalmic Pediatr Genet* 1987;8:63-8.
- Reese AB. Retinoblastoma. In: Reese AB, editor. *Tumors of the eye*. 1st ed. New York: Harper and Row; 1963. p. 155-6.
- Amendola BE, Lamm RF, Markoe AM, Karlsson UL, Shields JA, Shields CL, *et al.* Radiotherapy of retinoblastoma. A review of 63 children treated with different irradiation techniques. *Cancer* 1990;66:21-6.
- Scott IU, Murray TG, Feuer WJ, Van Quill K, Markoe AM, Ling S, *et al.* External beam radiotherapy in retinoblastoma. Tumor control and comparison of two techniques. *Arch Ophthalmol* 1999;117:766-70.
- Ali MJ, Gupta R, Vemuganti GK, Honavar SG. Histopathology of retinoblastoma after primary chemotherapy. *Proceedings of the XIV International Congress of Ocular Oncology*. Cambridge; 2009. p. 296.
- Honavar SG, Singh AD. Management of advanced retinoblastoma. *Ophthalmol Clin North Am* 2005;18:65-73.
- Shields CL, Shields JA. Retinoblastoma management: Advances in enucleation, intravenous chemoreduction and intra-arterial chemotherapy. *Curr Opin Ophthalmol* 2010;21:203-12.
- Bakhshi S, Meel R, Mohanti BK, Hasan Naqvi SG. Treatment and outcomes of non-metastatic extra-ocular retinoblastoma with uniform chemotherapy protocol. *J Pediatr Hematol Oncol* 2010;32:42-5.
- Kremens B, Wieland R, Reinhart H, Neubert D, Beck JD, Klingebiel T, *et al.* High dose chemotherapy with autologous stem cell in children with retinoblastoma. *Bone Marrow Transplant* 2003;31:281-4.
- Lee SH, Yoo KH, Sung KW, Kim JY, Cho EJ, Koo HH, *et al.* Tandem High-dose chemotherapy and autologous stem cell rescue in children

- with bilateral advanced retinoblastoma. *Bone Marrow Transplant* 2008;42:385-91.
40. White L. Chemotherapy in retinoblastoma: Current status and future directions. *Am J Pediatr Hematol Oncol* 1991;13:189-201.
 41. Munier FL, Verwey J, Pica A, Balmer A, Zografos L, Abouzied H, *et al.* New developments in external beam radiotherapy in retinoblastoma: From lens to normal tissue-sparing techniques. *Clin Exp Ophthalmol* 2008;36:78-89.
 42. Lin HY, Liao SL. Orbital development in survivors of retinoblastoma treated with enucleation with hydroxyapatite implant. *Br J Ophthalmol* 2010.
 43. Williams BK Jr, Scheffler AC, Garonzik SN, Gologorsky D, Shi W, Cavalcante LL, *et al.* Frequent prosthesis refitting to prevent implant exposure in patients with retinoblastoma. *J Pediatr Ophthalmol Strabismus* 2010;23:1-9.
 44. Stricker M, Simon E. Orbital reconstruction in children. *Neurochirurgie* 2010;56:287-93.
 45. Baj A, Lagana F, Beltramini GA. Anophthalmic socket syndrome: A new free tissue transfer. *J Oral Maxillofac Surg* 2010;68:2593-7.
 46. Maat-Kevit JA, Oepkes D, Hartwing NG, Vermeij-Keers C, Van Kamp IL, Van de Kamp JJ. A large retinoblastoma detected in a fetus at 21 weeks of gestation. *Prenat Diagn* 1993;13:377-84.
 47. Salim A, Wiknjosastro GH, Danukusumo D, Barnas B, Zalud I. Fetal retinoblastoma. *J Ultrasound Med* 1998;17:717-20.
 48. Douira-Khoms W, Korchane N, Louati H, Bhouiri L, Kchaou I, Ben Hassine L, *et al.* MRI exploration for the evaluation of orbital implant biocolonization in children enucleated for retinoblastoma. *J Fr Ophthalmol* 2009;32:540-3.
 49. Kiratli PO, Kiratli H, Ercan MT. Visualization of orbital retinoblastoma with technetium-99m (V) dimeraptosuccinic acid. *Ann Nucl Med* 1998;12:157-9.
 50. Moll AC, Hoekstra OS, Imhof SM, Comans EF, Schouten-van Meeteren AY, Van der Valk P, *et al.* Fluorine-18 fluorodeoxyglucose positron emission tomography (PET) to detect vital retinoblastoma in the eye: Preliminary experience. *Ophthalmic Genet* 2004;25:31-5.
 51. Parsam VL, Kannabiran C, Honavar S, Vemuganti GK, Ali MJ. A comprehensive, sensitive and economical approach for the detection of mutations in RB1 gene in retinoblastoma. *J Genet* 2009;88:517-27.
 52. Ali MJ, Parsam VL, Honavar SG, Kannabiran C, Reddy VA. RB1 gene mutations in retinoblastoma and its clinical correlation. *Saudi J Ophthalmol* 2010;24:119-23.
 53. Zhang G, He B, Weber GF. Growth factor signaling induces metastasis genes in transformed cells: Molecular connections between Akt Kinases and osteopontin in breast cancer. *Mol Cell Biol* 2003;23:6507-19.
 54. Wang J, Levenson AS, Satcher RL Jr. Identification of unique sets of genes altered during cell-cell contact in an in-vitro model of prostate cancer bone metastasis. *Int J Mol Med* 2006;17:849-56.
 55. Hartl M, Karagiannidis AI, Birster K. Cooperative cell transformation by Myc/Mil(Raf) involves induction of AP-1 and activations of genes implicated in cell motility and metastasis. *Oncogene* 2006;25:4043-55.
 56. Weber GF. The metastasis gene osteopontin: A candidate target for cancer therapy. *Biochim Biophys Acta* 2001;1552:61-85.
 57. Yu Y, Davicioni E, Triche TJ, Merlino G. The homeoprotein six1 transcriptionally activates multiple protumorigenic genes but requires ezrin to promote metastasis. *Cancer Res* 2006;66:1982-9.
 58. Berger JC, Vander Griend DJ, Robinson VL, Hickson JA, Rinker-Schaeffer CW. Metastasis suppressor genes: From gene identification to protein function and regulation. *Cancer Biol Ther* 2005;4:805-12.
 59. Keller ET. Metastasis suppressor genes: A role for raf kinase inhibitor protein (RKIP). *Anticancer Drugs* 2004;15:663-9.
 60. Kauffman EC, Robinson VL, Stadler WM, Sokoloff MH, Rinker-Schaeffer CW. Metastasis suppression: The evolving role of metastasis suppressor genes for regulating cancer cell growth at the secondary site. *J Urol* 2003;169:1122-33.
 61. Leander C, Fu LC, Pena A. Impact of an education programme on the late diagnosis of retinoblastoma in Honduras. *Pediatr Blood Cancer* 2007;49:817-9.
 62. Wilimas JA, Wilson MW, Haik BG, Barnoya M, Fu L, Castellanos M, *et al.* Development of retinoblastoma programs in Central America. *Pediatr Blood Cancer* 2009;53:42-6.

Cite this article as: Ali MJ, Reddy VP, Honavar SG, Naik M. Orbital retinoblastoma: Where do we go from here?. *J Can Res Ther* 2011;7:11-4.

Source of Support: Nil, **Conflict of Interest:** None declared.

Announcement

"Quick Response Code" link for full text articles

The journal issue has a unique new feature for reaching to the journal's website without typing a single letter. Each article on its first page has a "Quick Response Code". Using any mobile or other hand-held device with camera and GPRS/other internet source, one can reach to the full text of that particular article on the journal's website. Start a QR-code reading software (see list of free applications from <http://tinyurl.com/yzlh2tc>) and point the camera to the QR-code printed in the journal. It will automatically take you to the HTML full text of that article. One can also use a desktop or laptop with web camera for similar functionality. See <http://tinyurl.com/2bw7fn3> or <http://tinyurl.com/3ysr3me> for the free applications.