#### Elsevier Editorial System(tm) for Brachytherapy Manuscript Draft

Manuscript Number: 2013-H14

Title: The 2013 American Brachytherapy Society (ABS) Guidelines for Plaque Brachytherapy of Uveal

Melanoma and Retinoblastoma

Article Type: Regular Article

Keywords: Plaque; brachytherapy; radiation; guidelines; methods; ABS; AJCC; consensus

Corresponding Author: Dr. Paul T. Finger, MD

Corresponding Author's Institution: The New York Eye Cancer Center

First Author: Paul T. Finger, MD

Order of Authors: Paul T. Finger, MD

Manuscript Region of Origin: USA

Abstract: Purpose: To present the American Brachytherapy Society (ABS) guidelines for plaque brachytherapy of choroidal melanoma and retinoblastoma.

Methods and Materials: An international, multicenter Ophthalmic Oncology Task Force (OOTF) was assembled to include 47 radiation oncologists, medical physicists and ophthalmic oncologists from 10 countries. The ABS-OOTF produced collaborative guidelines, based on their eye cancer specific clinical experience and knowledge of the literature. This work was reviewed and approved by the ABS Board of Directors.

Results: The ABS-OOTF reached consensus that ophthalmic plaque radiation therapy is best performed in subspecialty brachytherapy centers. Quality assurance, methods of plaque construction and dosimetry should be consistent with the 2012 joint guidelines of the American Association for Physicists in Medicine and ABS. Implantation of plaque sources should be performed by subspecialty trained surgeons. Though there exist select restrictions related to tumor-size and location; the ABS-OOTF agreed that most uveal melanomas of the iris, ciliary body and choroid could be treated with plaque brachytherapy. The ABS-OOTF reached consensus that tumors with gross orbital extension, blind painful eyes and those with no light perception vision are unsuitable for brachytherapy. In contrast, only select retinoblastomas are eligible for plaque brachytherapy. Radionuclide available sources currently include assemblies of gold-shells with low-energy photon seeds (iodine-125, palladium-103, cesium-131), or solid ruthenium-106 or strontium-90 beta-emitting plaques. Prescription doses, dose-rates, treatment durations and clinical methods are described. Conclusions: Plaque brachytherapy is an effective, eye and vision-sparing method to treat patients with intraocular tumors. Practitioners are encouraged to use ABS-OOTF guidelines to enhance their practice.

#### Conclusion

# Conclusions

The American Brachytherapy Society-Ophthalmic Oncology Task Force, comprised of 47 eye cancer specialists from 10 countries, present our current guidelines and methods of plaque brachytherapy for uveal melanoma and retinoblastoma. We point out what is currently accepted as known, unknown as well as a need for standardization, staging and future research.



Sunday, September 8, 13

M. J. Zelefsky, MD
Editor-in-Chief
Brachytherapy
Memorial Sloan-Kettering Cancer Center
Department of Radiation Oncology
1275 York Avenue, Box 22
New York, NY 10065

Dear Dr. Zelefsky:

Please consider our manuscript titled, "The 2013 American Brachytherapy Society (ABS) Guidelines for Plaque Brachytherapy of Uveal Melanoma and Retinoblastoma," for publication in your journal, *Brachytherapy*.

When the board of The American Brachytherapy Society asked me form a committee to develop guidelines for ophthalmic plaque brachytherapy, I took this opportunity to greatly expand and internationalize this effort. I hope you will agree that this process has strengthened the guidelines and made them more broadly applicable and valuable.

However, in that each and every author has contributed to this work and the ABS agreed that they should be recognized for their contributions. Therefore, should you find these guidelines worthy of publication, I respectfully request that every authors name be recognized/linked to the article in PubMed and other such services.

Thank you for your time and consideration.

Best regards,

Paul

Paul T. Finger, MD

"This is original work which has never been published before. The author and all coauthors affirm that this manuscript has not been nor will be submitted elsewhere while under consideration by Brachytherapy."

The New York Eye Cancer Center 115 East 61st St, Suite #5B New York City, New York 10065 Tel: (212) 832-8170 Fax: (212) 888-4030 http://www.eyecancer.com pfinger@eyecancer.com

Director
Ocular Tumor Services:
The New York Eye and Ear Infirmary
NYU School of Medicine
Manhattan Eye, Ear and Throat Hospital

#### List of Abbreviations

ABS - American Brachytherapy Society OOTF - Ophthalmic Oncology Task Force PTF - Paul T Finger FAF – fundus autofluorescence CT – computed tomography MRI - magnetic resonance imaging PET/CT – positron emission tomography / computed tomography T - tumor N – node M -metastasis TTT – transpupillary thermotherapy Gy – gray KeV - kilo-electron volt <sup>60</sup>Co – cobalt-60 <sup>103</sup>Pd – palladium-103 <sup>125</sup>I – iodine-125  $^{106}$ Ru – ruthenium-106 <sup>131</sup>Cs – cesium-131 <sup>90</sup>Sr – strontium-90 VEGF – vascular endothelial growth factor NCI - National Cancer Institute NIH - National Institute of Health DHHS - Department of Health and Human Services AJCC- American Joint Committee on Cancer UICC - Union International for Cancer Control

RPT – ruthenium plaque therapy

1 **ABSTRACT** 2 **Purpose:** To present the American Brachytherapy Society (ABS) guidelines for plaque 3 brachytherapy of choroidal melanoma and retinoblastoma. 4 Methods and Materials: An international, multicenter Ophthalmic Oncology Task Force 5 (OOTF) was assembled to include 47 radiation oncologists, medical physicists and ophthalmic 6 oncologists from 10 countries. The ABS-OOTF produced collaborative guidelines, based on 7 their eye cancer specific clinical experience and knowledge of the literature. This work was 8 reviewed and approved by the ABS Board of Directors. 9 **Results:** The ABS-OOTF reached consensus that ophthalmic plaque radiation therapy is best 10 performed in subspecialty brachytherapy centers. Quality assurance, methods of plaque 11 construction and dosimetry should be consistent with the 2012 joint guidelines of the American 12 Association for Physicists in Medicine and ABS. Implantation of plaque sources should be 13 performed by subspecialty trained surgeons. Though there exist select restrictions related to 14 tumor-size and location; the ABS-OOTF agreed that most uveal melanomas of the iris, ciliary 15 body and choroid could be treated with plaque brachytherapy. The ABS-OOTF reached 16 consensus that tumors with gross orbital extension, blind painful eyes and those with no light 17 perception vision are unsuitable for brachytherapy. In contrast, only select retinoblastomas are 18 eligible for plaque brachytherapy. Radionuclide available sources currently include assemblies of 19 gold-shells with low-energy photon seeds (iodine-125, palladium-103, cesium-131), or solid 20 ruthenium-106 or strontium-90 beta-emitting plaques. Prescription doses, dose-rates, treatment 21 durations and clinical methods are described. 22 **Conclusions:** Plaque brachytherapy is an effective, eye and vision-sparing method to treat 23 patients with intraocular tumors. Practitioners are encouraged to use ABS-OOTF guidelines to

24 enhance their practice. 25 **KEY WORDS:** Plaque, brachytherapy, radiation, guidelines, methods, ABS, AJCC, consensus 26 **INTRODUCTION** Brachytherapy has been used to treat intraocular tumors since 1930. Subsequent reports 27 28 described cobalt-60 (<sup>60</sup>Co), ruthenium-106 (<sup>106</sup>Ru), iodine-125 (<sup>125</sup>I), palladium-103 (<sup>103</sup>Pd), strontium-90 (<sup>90</sup>Sr) and cesium-131 (<sup>131</sup>Cs) plaque sources.<sup>2-12</sup> Despite the international use of 29 30 ophthalmic brachytherapy for both uveal melanoma and retinoblastoma, there exist no 31 prospective randomized or case matched clinical trials comparing the clinical effectiveness or 32 side effects related to these radionuclides. The sole standardized clinical trial for choroidal melanoma, The Collaborative Ocular Melanoma Study (COMS), was restricted to the use of <sup>125</sup>I 33 plagues. 13 34 35 In 1985 the COMS provided the first standardized methods for multicenter tumor diagnosis, plaque construction and <sup>125</sup>I plaque dosimetry. <sup>14</sup> Then the COMS conducted a 12-year study that 36 demonstrated the relative equivalence of <sup>125</sup>I plaque compared to enucleation (removal of the 37 eye) for prevention of metastatic melanoma for a specific cohort of select medium-sized 38 choroidal melanoma. <sup>13</sup> An unintended consequence was that the method of using <sup>125</sup>I seeds in 39 40 COMS-shaped gold carrier plaques was established as the most common plaque method in North America. <sup>15</sup> Similarly, Lommatzsch and others have established a long tradition of using <sup>106</sup>Ru 41 plaque therapy in Europe. 16-21 This difference in practice currently persists. 42 43 The guidelines defined herein will exclude general aspects recently published by the American Association for Physicist in Medicine (AAPM) and the American Brachytherapy 44 Society (ABS). 22,23 The AAPM Task Group-129 (TG-129) has recently provided medical physics 45 guidelines in two publications. The first compared the currently available methods of plaque 46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

treatment planning and contrasted the patterns of intraocular dose deposition of <sup>103</sup>Pd and <sup>125</sup>I plaques for an average sized hypothetical intraocular tumor located at a variety of positions within the eye. <sup>23</sup> Therein, comparative dosimetry revealed that the lower energy photons from <sup>103</sup>Pd irradiation were more rapidly absorbed within the target zone (hypothetical tumor and 2 mm surround) with less irradiation to most normal ocular structures. The second AAPM TG-129 report was published with the ABS and offers preferred methods for dose calculation, plaque handling and quality assurance.<sup>22</sup> That report also includes an appendix describing current clinical controversies and applications. Herein, we supplement the aforementioned work with an ABS sanctioned study of clinical eye plaque brachytherapy. A panel of eye cancer specialists was assembled to broadly reflect current multicenter, international practice patterns. Thus, the ABS Ophthalmic Oncology Task Force (ABS-OOTF) includes a total of 47 ophthalmic oncologists, medical physicists and radiation oncologists from Canada, Finland, France, Germany, India, Japan, United Kingdom, United States of America, Russia and Sweden. Charged with developing modern guidelines for the use of plaque brachytherapy for uveal melanoma and retinoblastoma, consensus methods and indications for treatment are presented.

#### **METHODS AND MATERIALS**

#### Formation of the Committee

This study involved a review of the literature. This included but was not limited to searching PubMed for the terms: brachytherapy, choroid, iris, ciliary body, orbit, melanoma, retinoblastoma, <sup>125</sup>I, <sup>103</sup>Pd, <sup>106</sup>Ru, <sup>90</sup>Sr, <sup>60</sup>Co, <sup>131</sup>Cs, radionuclide, plaque, slotted, notched, proton beam, helium-ion, cyberknife, stereotactic radiosurgery, intensity modulated radiation therapy, extrascleral extension, COMS, dose, dose rate and side effects. This review was supplemented

by the participating authors' general working knowledge of the literature.

In addition, internet-based surveys (SurveyMonkey, Palo Alto, California, USA) of the subjects explored herein were sent to the participating eye cancer specialists. The results of the literature review and survey was adapted to the journal, Brachytherapy's instructions for authors by the corresponding author (PTF). Then, every ABS-OOTF member was allowed at least one opportunity to review and comment. Based on this feedback, the report was edited and returned to at least one representative from each center for a second review. As possible, all comments and suggestions were included in this report. In addition, the report was submitted to the ABS for additional review prior to submission for publication.

Many important recommendations of the ABS-OOTF were graded using levels of consensus modified from the 2003 ABS levels of Nag et al (Table 1).<sup>24</sup>

#### ABS-OOTF Recommended Methods

The ABS-OOTF recommends that plaque procedures be performed in specialized medical centers with expertise in ophthalmic brachytherapy (Level 1 Consensus). Further, it was agreed that these centers read and become familiar with the 2011 and 2012 published eye plaque dosimetry, construction and quality assurance guidelines published by the TG-129 and ABS. <sup>22,23</sup> In addition, each program should have written quality assurance guidelines functionally in place at their institutions. The results of the ABS-OOTF review of the literature, our clinical experience and collective judgment are as follows.

#### Case Selection

The diagnosis of uveal melanoma and retinoblastoma is complex. However, modern methods have greatly improved the accuracy of clinical diagnosis. Though patient history and physical examination (slit-lamp, ophthalmoscopy) are indispensible, state of the art ophthalmic oncology

services also utilize high and low-frequency ultrasound imaging, photography, intraocular angiography, fundus autofluorescence imaging (FAF), optical coherence tomography (OCT), computed tomography (CT), magnetic resonance imaging (MRI), positron emission tomography/computed tomography (PET/CT) and biopsy. In addition, wide-field fundus photography (RetCam, Clarity Medical Systems, Pleasanton, California, USA) has become indispensible for the diagnosis, staging and for monitoring the effects of retinoblastoma treatment. Though beyond the scope of this work, multimodality ophthalmic imaging plays an increasingly integral role in tumor diagnosis and follow up. While the initial diagnosis, follow-up for tumor control and intraocular side effects are best revealed by the ophthalmic oncologist; these results should be periodically examined and reported by each brachytherapy center.

#### Uveal Melanoma

Indications for the use of plaque therapy have expanded since the 2003 ABS guidance (Table 2). Page 12. Page 20. Page 20

#### Special Circumstances: Uveal Melanoma

1) There exists a controversy (Level 3 Consensus) about treatment of certain uveal melanomas.

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

For example, in diagnosis of "small" American Joint Committee on Cancer (AJCC) T1 uveal melanomas, the ABS-OOTF recommends (Level 2 Consensus) that in the absence of thickness >2 mm, subretinal exudative fluid and superficial orange pigment lipofuscin tumors; patients could be offered the alternative of "observation" for evidence of change (within 6 months) typically for documented growth prior to intervention. <sup>39-42</sup> This is particularly applicable for tumors near the fovea and optic nerve, or monocular patients where treatment is likely to cause radiation related vision morbidity. <sup>25,43-45</sup> Patients should also be counseled concerning the as yet unquantified albeit small risk of metastasis related to "observation as treatment." 2) Ocular melanosis, the Nevus of Ota and even natural pigmentation can darken the uvea and can prevent successful intra-operative tumor transillumination. This (in turn) makes definition of the targeted zone and plaque placement particularly difficult. These cases typically require experience and skills in scleral depression, focal transscleral transillumination (fiber optic or HeNe) and intraoperative ultrasound imaging to confirm proper plaque placement. 3) Select centers routinely biopsy uveal melanomas for pathology, genetic and molecular biologic analysis. 46,47 However, patients must be counseled that studies of the ocular and metastatic risks of biopsy have been small, limited in follow up, single center and thus did not reach level 2 consensus.<sup>48</sup> 4) Brachytherapy for tumors near, touching or surrounding the optic disc is also controversial.<sup>25</sup> As seen within the eye, the optic disc diameter is typically 1.8 mm. However as the optic nerve exits the eye into the orbit, it is surrounded by additional components such as the optic nerve sheath and widens to 5 to 6 mm. Thus, if a round plaque is perfectly placed against the retrobulbar optic nerve sheath, its posterior extent will be at least 1.5 mm from the edge of the optic disc. Therefore, the orbital optic nerve size prevents standard plaque positioning as to cover

139 the tumor and safety margin. In the past, 4-mm notches were placed in plaques to compensate. 140 However, 4-mm notches cannot overcome the 5 to 6 mm optic nerve sheath obstruction to allow 141 proper plaque positioning. In 2005, slotted plaques were devised with 8-mm openings. 25 Slots allow the optic nerve 142 143 sheath to enter the plaque, thus more posteriorly locate the seed sources and move the targeted 144 zone into a normalized position (surrounding the choroidal melanoma). It is important to note 145 that plaque slots make dosimetry more complex. In these cases, medical physicists must locate 146 seed sources to both "fill-in" the gap created by the slot and complete the targeted zone. However, the ABS-OOTF also recognizes that the penumbra at the edge of beta (106Ru and 147 <sup>90</sup>Sr) plagues is relatively sharp compared to the low energy gamma of <sup>125</sup>I and <sup>103</sup>Pd 148 plagues. 22,33,49,50 Thus, tumor tissue within the slot is likely to receive less radiation with slotted 149 <sup>106</sup>Ru and <sup>90</sup>Sr plaques compared to <sup>125</sup>I and <sup>103</sup>Pd slotted plaques in treatment of juxtapapillary 150 151 and circumpapillary tumors. 152 Uveal Melanoma Metastasis 153 The ABS-OOTF recommends (Level 2 Consensus) that all patients with uveal melanoma should be evaluated for metastatic disease prior to treatment.<sup>51</sup> However, staging methods vary 154 155 throughout the world. They range from relatively non-specific hematologic surveys, chest x-rays, 156 ultrasonographic or radiographic imaging of the abdomen (MRI or CT) to total body positron emission tomography/computed tomography (PET/CT). <sup>51,52</sup> The ABS-OOTF notes a trend 157 158 toward greater use of abdominal ultrasound screening in Europe and Russia. However, all 159 regimens focus on the liver as primary or sentinel organ at risk. We agree that early detection of 160 metastatic melanoma often allows for adjunctive systemic therapy. A statistically significant 161 comparison of the efficacy of each form of metastatic survey has not been performed.

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

The ABS-OOTF recommends (Level 2 Consensus) that the presence of metastatic disease from uveal melanoma is not an absolute contraindication for brachytherapy. For example, there exist ocular situations where brachytherapy may limit or prevent vision loss from tumorassociated retinal detachment or when tumor growth will soon cause secondary angle closure glaucoma. In addition, brachytherapy of the primary tumor may allow the patient to enter systemic treatment trial where a small proportion will survive. The ABS-OOTF does not recommend brachytherapy for patients whose death is imminent or those who cannot tolerate surgery. Retinoblastoma Brachytherapy is less commonly used as a primary treatment for retinoblastoma. <sup>20,53</sup> More frequently, radioactive plaques are used after local treatment failure (after cryotherapy, chemotherapy (systemic or ophthalmic artery perfusion), transpupillary thermotherapy (TTT), external beam radiation therapy or a combination thereof).<sup>54</sup> For example, a specific indication for plaque treatment may be found in cases were there is residual macular Rb following optimal chemoreduction. In these cases, TTT laser would surely affect the patients potential for vision. The ABS-OOTF recommends (Level 2 Consensus) that ideal tumors for primary brachytherapy are located anterior to the eyes equator and found in unilaterally affected children. For secondary treatment, residual or recurrent tumors are treated irrespective of location. However, in consideration of unusual tumor locations, anterior segment involvement is typically an indication for enucleation and there exist no reports of slotted plaque therapy for juxtapapillary retinoblastoma. Reflecting a generalized effort to avoid, EBRT, non-plaque brachytherapy implants have been used for orbital recurrence of retinoblastoma. 55,56

Systemic evaluations for retinoblastoma vary widely but typically consist of orbital and

intracranial MRI imaging. Due to its ionizing radiation component, CT imaging is employed only when MRI is not available. In high-risk patients, imaging is coupled with lumbar puncture and bone marrow aspiration biopsy.

Determinations of metastatic risk are typically based on clinical and histopathologic staging of the enucleated eye. <sup>57,58</sup> However, fewer eyes are being enucleated due to chemoreduction with subsequent alternative treatments and the recent use of selective ophthalmic arterial perfusion for advanced intraocular disease. Both these techniques likely result in down-staging, where histopathologic markers for metastasis may disappear leaving only clinical staging. <sup>58-60</sup> Therefore, prior to plaque therapy being considered, the ABS-OOTF recommends (Level 2 Consensus) that children whose retinoblastomas may have had access to aqueous outflow passages, massive (> 3mm) involvement of the choroid, post-laminar invasion of the optic nerve, extraocular extension and bilateral disease undergo systemic staging.

# Plaque Treatment Planning

Communication between the radiation oncologist, ophthalmic oncologist and medical physicist is critical for any successful brachytherapy program (Level 2 Consensus). In order to facilitate this communication, a treatment form and fundus diagram should be available to all participating specialists. It should be made part of the radiation oncology medical record and be available to the surgeon in the operating room.

1. The treatment form contains demographic identifying information about the patient, laterality of the involved eye, the largest basal dimension of the tumor, when treatment is scheduled and contact information for the treating eye cancer specialists. Each tumor should be staged according to the latest American Joint Committee on Cancer or equivalent Union for International Cancer Control staging system (currently the 7<sup>th</sup>

208	edition). 61,62			
209	2. The fundus diagram should be created as to demonstrate the tumors clock hour			
210	orientation within the eye, its longitudinal and transverse diameters and its largest basal			
211	diameter. It should include measurements from the tumor to the fovea, optic nerve, lens			
212	and opposite eye wall. This information is typically derived from judgments correlating			
213	the ophthalmic examination, ultrasound findings and photographic images. The ABS-			
214	OOTF agreed (Level 2 Consensus) that neither computed tomography nor magnetic			
215	resonance imaging currently offer superior tumor measurements.			
216	The medical physicist transfers this information to a computerized treatment planning system.			
217	Though described by the joint AAPM/ABS TG-129 report, this process also requires a			
218	determination of the radionuclide, prescription dose and dose rate. For those centers using			
219	radioactive seeds, there must also be seed selection and orientation. The ABS-OOTF			
220	recommends that all centers perform pre-implant treatment planning with documentation of			
221	doses to critical structures. <sup>23</sup> The ABS-OOTF also recommends that each plaque dosimetry plan			
222	undergo independent verification by a qualified medical physicist. The methods of preplanning,			
223	dose calculation, plaque design, plaque handling and quality assurance are recently described in			
224	the TG-129 reports. <sup>22,23</sup>			
225	Radionuclide Selection			
226	The ABS-OOTF found that <sup>125</sup> I and <sup>103</sup> Pd plaques are used by 3 or more centers in North			
227	America, <sup>125</sup> I or <sup>106</sup> Ru in Europe, solely <sup>106</sup> Ru in Japan and both <sup>106</sup> Ru or <sup>90</sup> Sr sources in Russia.			
228	Russian <sup>90</sup> Sr plaques are currently used for uveal melanoma up to 2.5 mm in height and			
229	retinoblastoma up to 3 mm. <sup>10</sup>			
230	In that normal ocular tissue side effects are dose related (Level 1 Consensus), the ABS-OOTF			

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

suggests that each center should engage in an intraocular dose distribution comparison (tumor apex, tumor base, lens, fovea, optic nerve, opposite eye wall) of available radionuclide sources prior to radiation source selection. We also agree (Level 1 Consensus) that each radionuclide offers different energies, intraocular dose distributions and requirements for handling (Table 3). The ABS-OOTF recommends (Level 2 Consensus) the goal of treatment to be delivery of a curative dose to the tumor while offering the least possible radiation to normal ocular structures. Dose Prescription In survey of customs and practice of the ABS-OOTF centers, there exists significant variation in radionuclide characteristics, selection and prescription dose. We recognize the significant differences in dose distribution patterns and a lack of internationally accepted dosimetry standards for each radionuclide. Further, the ABS-OOTF could find no prospective randomized or case matched studies comparing the efficacy or side effects of available plaque radionuclide techniques. Therefore, specific ABS-OOTF recommendations concerning the relative risks and benefits of each technique were considered beyond the scope of this report. The ABS-OOTF guidelines offer an overview of the committee's current practices and published results. 6,17,19-21,37,38,63,42 Dose prescriptions for uveal melanoma typically range from 70 to 100 Gy to the tumors apex. Two ABS-OOTF centers report employing a minimum <sup>106</sup>Ru dose to the sclera and one center continues to use the COMS-mandated minimum 85 Gy of <sup>125</sup>I to 5 axial intraocular millimeters. Depending on the ABS-OOTF center, even higher tumor apex and minimum scleral "base" doses have been employed for both <sup>106</sup>Ru and <sup>90</sup>Sr plaques. The ABS-OOTF recommends (Level 1 Consensus) that the tumor apex or point of maximal thickness remains the prescription point. However, the prescription isodose line should

encompass the entire tumor. In that it may affect local control, dose rates should not be less than

0.60 Gy/h for iodine-125 or palladium-103 plaques.<sup>64</sup> Dose modifications may be appropriate to account for different tumor sizes, implant durations, threshold doses to critical normal ocular structures and the use of alternate radionuclide sources.

ABS-OOTF centers using <sup>106</sup>Ru plaques (Bebig, Eckert and Zeigler Corp., Berlin, Germany)

#### Plaque Selection

range from 5 to 7 days.

typically restrict tumor apical height less than a mean 6 mm and rarely use commercially available <sup>106</sup>Ru plaques larger than 20 mm in diameter. In contrast, centers using <sup>125</sup>I or <sup>103</sup>Pd plaques do not as closely restrict their treatments based on tumor thickness. These patients with tumors greater than 12 mm in apical height or 20 mm in base are advised of their guarded prognosis for retaining useful vision and are counseled regarding alternative therapies. The largest commercially available gold COMS-type plaque (Trachsel Dental Laboratory, Rochester, Minnesota, USA) is 22 mm in diameter.

The ABS-OOTF recommends (Level 1 Consensus) that tumor diameters should not exceed the diameter of the planning target volume to prevent geographic miss. Thus, plaque apertures should exceed the largest tumor diameter as to create a tumor-free margin of safety to prevent geographic miss. That said, centers that utilize <sup>106</sup>Ru plaques must adjust for the 1-mm rim of silver designed to surround the periphery of the source aperture or "window." For small tumors,

#### Retinoblastoma Brachytherapy Practice Patterns

particularly those treated with <sup>106</sup>Ru plaques, durations may be as short as 3 days. However, in

survey of ABS-OOTF centers, brachytherapy for uveal melanoma treatment durations typically

Eligible retinoblastomas are typically less than 15 mm in base and no more that 10 mm in thickness. <sup>20,53,54,65-67</sup> Some describe Group B (International Classification) as being the most

commonly applicable stage. The ABS-OOTF recommends (Level 2 Consensus) that vitreous seeding should be absent or within 2 mm of the tumor surface. Either low energy <sup>103</sup>Pd, <sup>125</sup>I (for thicker tumors) or <sup>106</sup>Ru plaques (for thinner tumors) have been employed. Using low energy plaques, a solitary retinoblastoma is typically treated with a dose of 40 - 50 Gy to the tumor apex over 3 to 5 days. Depending on the ABS-OOTF center, typically higher tumor apex doses have been used for both <sup>106</sup>Ru and <sup>90</sup>Sr plaques.

Murphree noted that a history of or synchronous treatment with chemotherapy potentiates radiation-related intraocular vasculopathy (retinopathy and optic neuropathy). <sup>65</sup> In these cases,

he advocated reduced apical <sup>125</sup>I prescription doses 20-25 Gy or allowing several months

287 PLAQUE SURGERY

between chemotherapy and brachytherapy. <sup>65</sup>

Survey of ABS-OOTF centers suggests that brachytherapy using both low energy photon-emitting sources (103Pd, 125I) and beta-emitting 103Ru have been performed as outpatient procedures. However, centers must comply with local government regulations. The surgeries should be performed under either general or regional anesthesia by a surgeon experienced in plaque insertion. Ocular muscles should be relocated if they interfere with plaque position. This includes both rectus and oblique muscles.

Typically localized by transpupillary or trans-ocular illumination of the globe, the tumor base shadows its subjacent sclera. The edges of the shadow are marked on the sclera with tissue dye. An additional 2-3 mm "free-margin" is typically measured and marked around the tumor base. The ABS-OOTF defines "normal plaque position" (Level 1 Consensus) that the targeted zone includes the tumors base and safety margin. The ABS-OOTF survey found that in comparison to  $^{103}$ Pd and  $^{125}$ I plaques, larger physical safety margins are typically used with  $^{106}$ Ru.

Extra care must be taken in transilluminating thicker (e.g. >5 mm thick) uveal melanomas. Here the tumor can cast eccentric shadows, thus yielding false tumor base diameters. Small posterior and amelanotic tumors can also be a challenge to mark. Here, two techniques are helpful including: posterior point source illumination (e.g. fiber optic or HeNe light sources or scleral depression combined with indirect ophthalmoscopy) with intraoperative ophthalmic ultrasound verification is employed. When this is not possible (e.g. iris and iridociliary melanoma) high frequency ultrasound imaging and direct transcorneal visualization play a more important role during intraoperative tumor localization.

In all cases, the plaque is sutured as to cover the marked, episcleral target zone. Then the extraocular muscles and conjunctiva are re-attached as not to disturb brachytherapy. When utilizing plaque with low-energy seeds, the eye is typically covered with a lead patch shield. Typically, after 5 to 7 days, the patient is returned to the operating room, where the plaque is removed under regional or general anesthesia. The ABS-OOTF agreed (Level 2 Consensus) that displaced muscles should be re-attached into their insertions after plaque removal. However one ABS-OOTF center did not find it necessary to reattach the inferior oblique muscle. If an amniotic membrane is used to buffer the cornea during brachytherapy, it should be removed prior to conjunctival closure.<sup>71</sup>

#### FOLLOW UP AFTER BRACHYTHERAPY

After brachytherapy patients are followed for local control, complications and systemic disease. Most ABS-OOTF centers examine treated eyes every 3 to 6 months. This time interval can be modulated based on the likelihood of secondary complications. For example, intervals are shorter for patients with posteriorly located tumors, at higher risk of radiation maculopathy and radiation optic neuropathy that typically occurs within the first 3 years follow up (see radiation

complications below). <sup>8,38,44,45,72</sup> Similarly, most local tumor recurrence occurs during the first 5 years. Therefore, larger and juxtapapillary tumors (at higher risk for regrowth) may require closer follow up. In addition, patients should be periodically re-examined for evidence of metastatic disease and second non-ocular primary cancers. <sup>51,52,73</sup> The ABS-OOTF agrees (Level 1 Consensus) that periodic radiographic abdominal imaging of the liver can be used to detect hepatic melanoma metastasis. We also concur that early detection yields patients with smaller tumor burdens who would more likely benefit from systemic treatment.

#### ALTERNATIVE SURGICAL TECHNIQUES

Uveal melanomas are alternatively be treated by enucleation or exenteration. The former is used when the tumor is confined to the eye and the latter considered in the presence of gross orbital tumor extension. Pre-enucleation external beam radiation therapy (PERT) has not been widely performed since the COMS large-melanoma study found no statistically significant survival advantage. <sup>52,74</sup> In contrast, most centers continue to apply post-exenteration radiation therapy in cases where there is residual orbital melanoma and retinoblastoma.

Local resection (internal evacuation or external lamellar sclerouvectomy) is employed to remove select (typically select medium-sized or large) uveal melanoma but not retinoblastoma. Some centers irradiate the uveal melanoma prior to endoresection or place a radioactive plaque over the tumors base after transscleral resection. Such adjunctive radiotherapy targets presumed residual melanoma that may seed the orbit or locally recur. Other centers consider vitreous melanoma seeds to be an indication for enucleation. The ABS- OOTF recognizes (Level 3 Consensus) that adjuvant radiotherapy may reduce the risk of local tumor recurrence. However, we also recognize there exist no prospective comparative or case-matched studies examining the relative risks and benefits of resection techniques compared to primary brachytherapy or

346 enucleation.

Retinoblastoma of stage AJCC T4 or International Classification D and E are not candidates for brachytherapy and are typically treated by enucleation.<sup>67</sup> The ABS-OOTF achieved Level 1 Consensus that primary enucleation prior to extraocular extension, optic nerve invasion and or massive choroidal infiltration offers greater than 95% primary tumor free survival.<sup>57,58,67</sup> While retinoblastomas with extrascleral tumor extension are treated with combinations of systemic chemotherapy, surgical excision (enucleation or exenteration) and external beam irradiation as well as systemic surveillance. There exists Level 1 Consensus that if possible, external beam radiation therapy should be avoided due to secondary carcinogenesis and orbital bone dysplasia.<sup>77,78</sup> Preferred practice patterns for treatment of retinoblastoma are even more complex and beyond the scope of this review.<sup>79</sup>

## ALTERNATIVE RADIATION THERAPY TECHNIQUES

Proton therapy was pioneered at the Harvard Cyclotron Laboratory and by researchers at the Massachusetts Eye and Ear Infirmary and Massachusetts General Hospital. <sup>80</sup> Since that time, at least 12 additional institutions around the world have embraced this technique with numerous additional centers under construction. <sup>81-83</sup> These centers typically use a proton radiobiologic effectiveness value of 1.1 compared to <sup>60</sup>Co. For uveal melanoma, doses of approximately 60 Gy are delivered in four (15 Gy) daily fractions. Though there exists no significant alpha/beta dose comparison high-dose-rate proton beam versus low dose rate plaque brachytherapy, the ABS-OOTF recognizes (Level 1 Consensus) that both the dose rates and dose volumes differ. Further, we agree (Level 1 Consensus) that all external beam radiation techniques (proton, helium-ion, gamma-knife and stereotactic radiosurgery) require an anterior ocular and/or adnexal entry dose with resultant dose-related collateral damage to those exposed normal tissues (even when

385

386

387

388

389

390

391

369 treating posterior tumors). However, we also recognize (Level 1 Consensus) that there is relative 370 dose sparing of tissues posterior and lateral to the proton beam. 371 In contrast, plaque brachytherapy places the source on the sclera beneath (adjacent to) the 372 tumor. Thus, in treatment of posterior choroidal melanomas, radiation must travel through the 373 sclera prior to entering the tumor and through the eye prior to exiting through normal anterior ocular tissues.<sup>23</sup> Primarily due to dose gradient and side-scatter effects, plaque brachytherapy 374 375 delivers comparatively more radiation to subjacent sclera and adjacent ocular structures. 376 The ABS-OOTF recognizes (Level 1 Consensus) that in treatment of posterior uveal 377 melanomas, there is less resultant radiobiologic effect on normal anterior ocular structures using low energy (<sup>103</sup>Pd, <sup>125</sup>I) plaque brachytherapy in comparison to proton beam. This relative dose 378 379 sparing may explain why clinical studies have revealed more anterior segment complications and secondary enucleations after charged-particle therapy. 81,84-88 380 381 External beam radiation techniques are also complicated by mobile target volume (eye 382 movement). Consider that eye plaques are sewn to the eye wall beneath their target volume. 383 When the eye moves, so does the plaque. In contrast, when a target volume is externally created 384

movement). Consider that eye plaques are sewn to the eye wall beneath their target volume. When the eye moves, so does the plaque. In contrast, when a target volume is externally created to extend within the eye (all EBRT techniques), the potential mobility of the eye and its target volume makes dose-deposition less predictable. During proton therapy, eye movements must be constantly monitored and the patient reminded (as needed) to fixate on a reference target. This is because eye movements cause misapplication of protons within the eye. In addition, should larger tumor-free safety margins become necessary, more normal tissues (anterior and posterior) fall within the cylindrical target volume. Lastly, proton beam facilities are vastly more expensive (Table 4). 89,90

The ABS-OOTF survey indicates that proton beam has been employed as an alternative to

enucleation for tumors considered unsuitable for brachytherapy. This includes tumors that touch or surround the optic disc, for very large tumors and where <sup>125</sup>I and <sup>103</sup>Pd plaques are not available. In addition, a novel strategy tries to prevent secondary inflammation, "vitritis" or "toxic tumor syndrome" has been described after brachytherapy for large choroidal melanoma. Here large uveal melanomas are first treated with proton beam, then removed by internal resection. <sup>75</sup> There are only a few centers using this technique (ABS-OOTF Level 3 Consensus).

#### CLINICAL RESULTS

Reporting the results of treatment is particularly challenging. Consider that when multiple centers use the same radionuclides source they often differ in plaque construction, dosimetry, dose and dose rate. Further, until acceptance of the AJCC staging system, their existed no universal method to report the size of uveal melanomas. Further, there is no uniform method of reporting with respect to follow up duration, visual acuity, local control or metastasis. Herein, we have assembled a non-inclusive table of representative case series with >100 treated patients (Table 5).

Select observations derived from Table 5 include: the radionuclides <sup>125</sup>I and <sup>106</sup>Ru are best represented and on average the data is over 10 years old. Note that a mean 341 patients were reported per center, average follow up was 4.5 years and tumor size reporting lacks AJCC or UICC standardization. With respect to treatment, the mean and median prescription dose was 83 Gy and 80 Gy respectively (range 70 Gy to 100 Gy). Similarly, reported and 5-year local control rates averaged 89.5% (range 69.9% to 97.9%). However, there exist no data to allow a meta-analysis comparing relative tumor size and location. In general, there exists no information concerning cases lost to follow up. Note that the median rates of metastasis are quite similar except for series reporting on larger tumors.<sup>35</sup> Lastly, visual acuity results varied widely.

Visual acuity outcomes are difficult to compare in that they depend on many factors including but not limited to: pre-existing exudative retinal detachments, subfoveal position, radiation dose to critical structures, cataract onset, cataract repair, secondary vitreous hemorrhage, radiation maculopathy, optic neuropathy and the availability of anti-vascular endothelial growth factor (anti-VEGF) treatment. Clearly, this outcome analysis supports the need for more uniform data collection and reporting among eye cancer specialists.

#### Radiation Complications Overview

Ophthalmic brachytherapy complications have been related to both radiation and patient specific factors. These include total dose, dose-rate, dose volume, dose to critical structures, tumor size, location and the biologically variable responses to irradiation.

#### Radiation Cataract

The ABS-OOTF survey indicates (Level 1 Consensus) that there exists no increased risk associated with radiation cataract removal. However, almost all centers recommended waiting until 6-12 months after brachytherapy.

#### Intraocular Radiation Vasculopathy

Radiation induces a progressive vasculopathy caused by loss of pericytes and endothelial cells. <sup>92</sup> Clinical findings include transudation of intravascular components (blood, serum, lipids) and small vessel closure (cotton-wool spots). First retinal findings include hemorrhages, edema and cotton-wool infarcts. However, it is the earlier onset radiation macular edema causes reversible vision loss. Later, small vessel closure leads to ischemia, neovascularization and irreversible atrophy. Variations of this process are also seen in the optic disc and iris.

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

The ABS-OOTF concur (Level 2 Consensus) that untreated radiation maculopathy and optic neuropathy typically results in poor visual acuity. The prognosis for vision diminishes with vasculopathy of the macula, optic nerve, vitreous hemorrhage and neovascular glaucoma. In that radiation maculopathy is the most common cause of radiation-associated vision loss, we present a classification for radiation retinopathy based on prognosis for vision (Table 6). The ABS-OOTF agreed (Level 2 Consensus) that intravitreal anti-VEGF therapy is useful to suppress radiation induced neovascular glaucoma, radiation maculopathy and optic neuropathy. 94-98 Therapy is used to suppress transudation, thus ameliorate edema and counter neovascularization. However, while these techniques are widely used, the ABS-OOTF recognizes that no published prospective-randomized or large-scale studies examining the effects relative to initial radiation dose, dose-rate or source. The literature also contains two alternative approaches to the treatment of radiation retinopathy. Laser photocoagulation in the form of posterior tumor demarcation resulted in sector devascularization best seen on fluorescein angiography. This technique along with sector pan retinal photocoagulation has been reported to slow or prevent radiation retinopathy by two independent centers. 99,100 Treatment converted slow ischemia within and anterior to the targeted zone to scar. In theory, laser devitalization of the ischemic tumor and treated retina may decrease intraocular production of VEGF. However, brachytherapy also affects the eyelids, eyelashes, conjunctiva, tear production, corneal surface integrity, sclera and ocular muscles. 8,101,102 Within the eye radiation can cause iritis, uveitis, synechiae, neovascular glaucoma, cataract, posterior neovascularization, hemorrhage, retinal detachment, retinopathy and optic neuropathy. The most common late, sight limiting posterior segment complication is radiation maculopathy. Unusual complications

include persistent strabismus and scleral thinning. All of the aforementioned side effects can result loss of vision and quality of life.

#### STAGING OF RADIATION SIDE EFFECTS

The ABS-OOTF recognizes that there exists no comprehensive staging system for the ophthalmic side effects of radiation therapy. Though many of these findings are fundamentally, albeit less specifically, classified by the United States National Cancer Institute (Cancer Therapy Evaluation Program, Common Terminology Criteria for Adverse Events, Version 4.0, DCTD, NCI, NIH, DHHS (<a href="http://ctep.cancer.gov">http://ctep.cancer.gov</a>), the ABS-OOTF recommends that a radiation specific ophthalmic side effect staging system be developed to improve communication for patient care, research and publication.

**DISCUSSION** 

This presentation of ABS-OOTF guidelines for ophthalmic plaque brachytherapy offers both consensus and controversy. We recommend that brachytherapy be performed by skilled plaque surgeons in experienced subspecialty centers. We agreed that the recent joint AAPM/ABS TG-129 published guidelines for plaque construction, dosimetry and quality assurance be read and widely employed at active centers. <sup>22,23</sup> We also concurred that many radionuclide sources can be employed, but only <sup>125</sup>I, <sup>103</sup>Pd and <sup>106</sup>Ru are used in 3 or more ABS-OOTF centers. While there exist tumor thickness restrictions for <sup>106</sup>Ru and <sup>90</sup>Sr, taller tumors can be treated with <sup>125</sup>I or <sup>103</sup>Pd techniques. <sup>7,11,22,49</sup>

Overall, the ABS-OOTF expanded general indications for uveal melanoma patient selection.

Lastly, we found that plaque brachytherapy is not commonly used for retinoblastoma. However, indications include small anterior tumors in unilateral cases, for salvage after chemoreduction with subsequent alternative therapies and in select cases where macular laser will likely cause

loss of vision.

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

#### **Unanswered Questions**

However, the ABS-OOTF acknowledges the myriad unanswered questions that challenge ophthalmic plaque brachytherapy researchers. Select questions offered by the ABS-OOTF include: What are the radiobiological differences between continuous low dose rate plaque brachytherapy in comparison with fractionated high dose rate proton beam irradiation? What is the "correct" apical prescription dose and dose-rate required for treatment of uveal melanoma and how do we accommodate for the steep dose-gradient within the tumor? For example, should there be a dose de-escalation study or a thickness-based sliding scale in treatment of uveal melanoma? Can there be international standards for dosimetry to determine the relative efficacy (alphas and betas) of photons, electrons and protons? Is there a role for radiation sensitizers during plaque therapy? Should the presence of intravitreal melanoma seeds affect case selection? What is the role and best timing for the use of anti-VEGF agents in treatment of radiation maculopathy and optic neuropathy? Are there differences in the efficacy of anti-VEGF agents related to radionuclide, radiation dose and dose rate? Do notched and slotted plaques address geographic miss in treatment of juxtapapillary and circumpapillary tumors? With regards to retinoblastoma, are there oncogenic risks of plaque brachytherapy? What are the optimal parameters for tumor selection and radiation dose (if used before or after chemotherapy)? The ASB-OOTF hopes future research will answer some of these questions. The ABS-OOTF recommends that the eye cancer community use universal AJCC –UICC staging to define tumor size, location and associated variables. 61,62 This would enable multicenter communication, comparative analysis and patient education. It would allow for collection of numbers large enough to reach statistical significance. The ABS-OOTF recommends the

development of a site-specific staging system for complications after ophthalmic radiation therapy. This would facilitate scientific comparisons between treatments, help predict ophthalmic side effects and improve informed consent.

508 SUMMARY

Currently, plaque brachytherapy offers an eye and vision-sparing alternative to enucleation annually for thousands of patients' worldwide. Herein, we present the current ABS guidelines for patient selection, informed consent and methods of treatment. For the first time, the ABS-OOTF includes eye cancer specialists from North American, Russia, Asia and Europe. We encourage all centers to use these guidelines to formulate their treatment patterns and reporting policies. However, we realize that such guidelines are dynamic and will need to be modified as to conform to ever evolving clinical evidence.

# **ACKNOWLEDGMENTS**

The research was supported (in part) by The Eye Cancer Foundation, Inc.

(<a href="http://eyecancerfoundation.net">http://eyecancerfoundation.net</a>) and The American Brachytherapy Society.

526	The ABS – OOTF Committee
527	
528	Canada - Princess Margaret Hospital
529	Sick Kids Hospital – Toronto, Ontario
530	E. Rand Simpson MD – Ophthalmic Oncology
531	Brenda Gallie MD – Ophthalmic Oncology
532	Norman Iaperriere MD – Radiation Oncology
533	Akbar Beiki-Ardakani MCCPM – Medical Physics
534	
535	Finland - Helsinki University Central Hospital
536	University of Helsinki, Helsinki
537	Tero Kivelä, MD – Ophthalmic Oncology
538	Virpi Raivio, MD – Ophthalmic Oncology
539	Jorma Heikkonen, PhD – Medical Physics
540	·
541	France - The Curie Institute, Paris
542	Laurence Desjardins, MD - Ophthalmic Oncology
543	Remi Dendale, MD – Radiation Oncology
544	Alexandro Mazal – Medical Physics
545	The state of the s
546	Germany - University of Duisburg-Essen, Essen
547	Norbert Bornfeld, MD - Ophthalmic Oncology
548	Wolfgang Sauerwein, MD – Radiation Oncology
549	Dirk Flü Dipl. Physics - Medical Physics
550	Lorenzo Brualla, PhD – Medical Physics
551	Lorenzo Bruana, i iib - Medicai i nysies
552	India Contro for Sight Supergrapialty Eve Hagnital Hydershad
553	India Santos anovar, MD – Ophthalmic Oncology
554	Vijay Anand P Reddy, MD, DNB – Radiation Oncology
555	Vijay Aliand P Reddy, MD, DNB – Radianon Oncology
	Janes National Consess Contas Hagnital Talma
556	Japan – National Cancer Center Hospital, Tokyo
557	Shigenobu Suzuki, MD – Ophthalmic Oncology
558	Naoya Murakami, MD – Radiation Oncology
559	
560	Russia - Helmholtz Research Institute of Eye Diseases, Moscow
561	Svetlana Saakyen MD, PhD – Ophthalmic Oncology, Radiology
562	Vladimir Walsl JD, PhD – Ophthalmic Oncology, Radiology
563	Anush Amiryan MD, PhD – Ophthalmic Oncology, Radiology
564	
565	Sweden - St. Erik's Eye Hospital, Stockholm
566	Stefan Seregard MD PhD - Ophthalmic Oncology
567	Charlotta All-Eriksson MD, PhD - Ophthalmic Oncology
568	Lars Hjelmqvist MD, PhD - Ophthalmic Oncology
569	Göran Lundell MD, PhD – Radiation Oncology
570	Georges Sinclair MD – Radiation Oncology
571	Marie Lundell MD, PhD – Medical Physics

572	<b>United Kingdom - Liverpool University Medical Center, Liverpool</b>
573	Bertil Damato, MD - Ophthalmic Oncology
574	R Doug Errington – Radiation Oncology
575	Philip Mayles – Medical Physics
576	Helen Mayles – Medical Physics
577	·
578	United States - Emory Eye Cancer
579	Emory University Medical Center Atlanta, Georgia
580	Chris Bergstrom MD – Ophthalmic Oncology
581	Hans Grossniklaus MD – Ophthalmic Oncology
582	Ian Crocker MD – Radiation Oncology
583	Elizabeth Butker M.Sc. – Medical Physics
584	
585	United States - University of Tennessee - Memphis
586	thodist University Hospital
587	wratt Wilson MD – Ophthalmic Oncology
588	Barrett Haik MD – Ophthalmic Oncology
589	Holger Geischen. MD – Radiation Oncology
590	Pradeep Patr Medical Physics
591	
592	United States - Tufts University Medical Center, Boston, Mass.
593	Jay Duker MD – Ophthalmic Oncology
594	John Mignano MD – Radiation Oncology
595	Mark Rivard Ph.D. – Medical Physics
596	
597	<b>United States - The New York Eye Cancer Center, New York City</b>
598	Beth Israel Comprehensive Cancer Center
599	The New York Eye and Ear Infirmary
600	Paul T. Finger, MD – Ophthalmic Oncology
601	Ekaterina Semenova, MD – Ophthalmic Oncology
602	Walter Choi, MD – Radiation Oncology
603	Nina I. Kalach MS – Medical Physics
604	
605	
606	
607	
608	
609	
610	
611	
612	
613	
614	
615	
616	

617		REFERENCES
618	1.	Moore R. Choroidal sarcoma treated by the intraocular insertion of radon seeds. $BrJ$
619		Ophthalmol. 1930;14:145-156.
620	2.	Stallard HB. Radiotherapy for malignant melanoma of the choroid. Br J Ophthalmol.
621		1966;50:147-155.
622	3.	Lommatzsch PK. Results after beta-irradiation (106Ru/106Rh) of choroidal
623		melanomas. Twenty years' experience. American journal of clinical oncology.
624		1987;10:146-151.
625	4.	Packer S, Rotman M. Radiotherapy of choroidal melanoma with iodine-125.
626		Ophthalmology. 1980;87:582-590.
627	5.	Sealy R, le Roux PL, Rapley F, et al. The treatment of ophthalmic tumours with low-
628		energy sources. <i>Br J Radiol.</i> 1976;49:551-554.
629	6.	Finger PT, Chin KJ, Duvall G, Palladium-103 for Choroidal Melanoma Study Group.
630		Palladium-103 ophthalmic plaque radiation therapy for choroidal melanoma: 400
631		treated patients. Ophthalmology. 2009;116:790-796.
632	7.	Rivard MJ, Melhus CS, Sioshansi S, et al. The impact of prescription depth, dose rate,
633		plaque size, and source loading on the central axis using 103Pd, 125I, and 131Cs.
634		Brachytherapy. 2008;7:327-335.
635	8.	Finger PT. Radiation therapy for choroidal melanoma. Surv Ophthalmol.
636		1997;42:215-232.
637	9.	Leonard KL, Gagne NL, Mignano JE, et al. A 17-year retrospective study of
638		institutional results for eye plaque brachytherapy of uveal melanoma using (125)I,
639		(103)Pd and (131)Cs and historical perspective <i>Brachytherany</i> 2011:10:331-339

640 **10.** Vakulenko MP, Dedenkov AN, Brovkina AF, et al. [Results of beta-therapy of 641 choroidal melanoma]. Med Radiol (Mosk). 1980;25:73-74. 642 11. Brovkina AF, Zarubei GD, Val'skii VV. Criteria for assessing the efficacy of 643 brachytherapy of uveal melanomas, complications of therapy and there prevention. 644 Vestn oftalmol. 1997;113:14-16. Murakami N, Suzuki S, Ito Y, et al. 106-Ruthenium plaque therapy (RPT) for 645 **12.** 646 retinoblastoma. Int I Radiat Oncol Biol Phys. 2012;84:59-65. 647 Collaborative Ocular Melanoma Study Group, The COMS randomized trial of **13.** 648 iodine- 125 brachytherapy for choroidal melanoma: V. Twelve-year mortality rates 649 and prognostic factors; COMS report No. 28. Arch Ophthalmol. 2006:124:1684-1693. 650 **14.** Collaborative Ocular Melanoma Study Group. Ch 12: Radiation Therapy. In: Service 651 NTI, ed. *COMS manual of procedures PB95-179693*. Springfield, Virginia, 1995. 652 **15.** Fontanesi J. Meyer D. Xu S. et al. Treatment of choroidal melanoma with I-125 653 plaque. Int J Radiat Oncol Biol Phys. 1993;26:619-623. 654 **16.** Bergman L, Nilsson B, Lundell G, et al. Ruthenium brachytherapy for uveal 655 melanoma, 1979-2003; survival and functional outcomes in the Swedish population. 656 Ophthalmology. 2005;112:834-840. 657 **17.** Summanen P, Immonen I, Kivelä T, et al. Visual outcome of eyes with malignant 658 melanoma of the uvea after ruthenium plaque radiotherapy. *Ophthal Surg lasers*. 659 1995;26:449-460. 660 **18.** Damato B, Patel I, Campbell IR, Mayles HM, et al. Visual acuity after Ruthenium (106) 661 brachytherapy of choroidal melanomas. Int I Radiat Oncol Biol Phys. 2005;63:392-662 400.

19. 663 Foerster MH. Bornfeld N. Wessing A. et al. Treatment of malignant melanomas of the 664 uvea with 106-ruthenium applicators. Report on the first 100 Essen cases. Klin 665 Monbl Augenheilkd. 1984;185:490-494. 666 **20**. Schueler AO, Fluhs D, Anastassiou G, et al. Beta-ray brachytherapy of 667 retinoblastoma: feasibility of a new small-sized ruthenium-106 plaque. *Ophthalmic* 668 Res. 2006;38:8-12. 669 21. Seregard S, aft Trampe E, Lax I, et al. Results following episcleral ruthenium plaque radiotherapy for posterior uveal melanoma. The Swedish experience. Acta 670 671 ophthalmol Scand. 1997;75:11-16. 672 **22**. Chiu-Tsao ST. Astrahan MA. Finger PT. et al. Dosimetry of (125) I and (103) Pd COMS 673 eve plagues for intraocular tumors: report of Task Group 129 by the AAPM and ABS. 674 Med Phys. 2012;39:6161-6184. Rivard MJ, Chiu-Tsao S-T, Finger PT, et al. Comparison of dose calculation methods 675 **23.** 676 for brachytherapy of intraocular tumors. *Med Phys.* 2011;38:306-316. 677 **24.** Nag S, Ouivey IM, Earle ID, et al. The American Brachytherapy Society 678 recommendations for brachytherapy of uveal melanomas Int I Radiat Oncol Biol Phys 679 2003;56:544-555. 680 **25**. Finger PT. Chin KI. Tena LB. A five-year study of slotted plague radiation therapy for 681 choroidal melanoma: Near, touching or surrounding the optic nerve. *Ophthalmology*. 682 2012;119:415-422. 683 **26**. Finger PT. Plaque radiation therapy for malignant melanoma of the iris and ciliary 684 body. Am J Ophthalmol. 2001;132:328-335.

685 **27**. Yousef YA. Finger PT. Lack of radiation maculopathy after palladium-103 plaque 686 radiotherapy for iris melanoma. *Int J Radiat Oncol Biol Phys* 2012;83:1107-1112. 687 **28**. Shields C, Naseripour M, Shields I, et al. Custom-designed plaque radiotherapy for 688 nonresectable iris melanoma in 38 patients: tumor control and ocular complications. 689 Am J Ophthalmol. 2003;135:648-656. 690 29. Petousis V, Finger PT, Milman T. Multifocal iris melanoma treated with total anterior 691 segment palladium-103 plaque radiation therapy. *Graefe's Arch Clin Exp Ophthalmol* 692 2011;249:937-940. 693 **30**. Fernandes BF, Krema H, Fulda E, et al. Management of iris melanomas with iodine-694 125 plaque radiotherapy. *Am I Ophthalmol.* 2010:149:70-76. 695 **31**. Krema H, Simpson ER, Pavlin CI, et al. Management of ciliary body melanoma with 696 iodine-125 plaque brachytherapy. Can J Ophthalmol. 2009;44:395-400. 697 32. Lumbroso-Le Rouic L, Charif Chefchaouni M, Levy C, et al. 125I plaque brachytherapy 698 for anterior uveal melanomas. Eye. 2004;18:911-916. 699 33. Brovkina AF, Zarubei GD, Fishkin IG. Validation of the use of brachytherapy in uveal 700 melanomas of juxtapapillary localization. Vestn oftalmol. 1991:107:41-44. 701 **34**. Gray ME, Correa ZM, Augsburger II, et al. Ciliary body melanoma with limited 702 nodular extrascleral extension and diffuse iris-angle infiltration treated by whole 703 anterior segment plaque radiotherapy. Int ophthalmol. 2007;27:273-276. 704 Shields CL, Naseripour M, Cater J, et al. Plaque radiotherapy for large posterior uveal **35.** 705 melanomas (> or =8-mm thick) in 354 consecutive patients. *Ophthalmology*.

706

2002;109:1838-1849.

- 707 **36.** Puusaari I, Heikkonen J, Summanen P, *et al.* Iodine brachytherapy as an alternative to enucleation for large uveal melanomas. *Ophthalmology.* 2003;110:2223-2234.
- 709 37. Puusaari I, Heikkonen J, Kivelä T. Ocular complications after iodine brachytherapy
   710 for large uveal melanomas. *Ophthalmology*. 2004;111:1768-1777.
- Puusaari I, Heikkonen J, Kivelä T. Effect of radiation dose on ocular complications
   after iodine brachytherapy for large uveal melanoma: empirical data and simulation
   of collimating plaques. *Invest Ophthalmol Vis Sci.* 2004;45:3425-3434.
- Augsburger JJ, Vrabec TR. Impact of delayed treatment in growing posterior uveal
   melanomas. *Arch Ophthalmol.* 1993;111:1382-1386.
- Sobrin L, Schiffman JC, Markoe AM, et al. Outcomes of iodine-125 plaque
   radiotherapy after initial observation of suspected small choroidal melanomas: a
   pilot study. *Ophthalmology*. 2005;112:1777-1783.
- Murray TG, Sobrin L. The case for observational management of suspected small
   choroidal melanoma. *Arch Ophthalmol.* 2006;124:1342-1344.
- 721 **42.** Semenova E, Finger PT. Palladium-103 Radiation Therapy for Small Choroidal
   722 Melanoma. *Ophthalmology*. Jun 14 2013 [Epub]
- Newman H, Chin KJ, Finger PT. Subfoveal choroidal melanoma: pretreatment characteristics and response to plaque radiation therapy. *Arch Ophthalmol.*2011;129:892-898.
- Finger PT, Chin KJ, Yu GP. Risk factors for radiation maculopathy after ophthalmic
   plaque radiation for choroidal melanoma. *Am J Ophthalmol.* 2010;149:608-615.
- 728 **45.** Finger PT. Tumour location affects the incidence of cataract and retinopathy after ophthalmic plaque radiation therapy. *Br J Ophthalmol.* 2000;84:1068-1070.

- 730 46. Onken MD, Worley LA, Char DH, et al. Collaborative Ocular Oncology Group report 731 number 1: prospective validation of a multi-gene prognostic assay in uveal 732 melanoma. Ophthalmology. 2012;119:1596-1603. 733 **47.** Harbour IW. The genetics of uveal melanoma: an emerging framework for targeted therapy. Pigment Cell Melanoma Res. 2012;25:171-181. 734 McCannel TA, Chang MY, Burgess BL. Multi-year follow-up of fine-needle aspiration 735 **48**. 736 biopsy in choroidal melanoma. *Ophthalmology*. 2012;119:606-610. 737 Brualla L. Sempau I. Zaragoza FI. et al. Accurate estimation of dose distributions 49. 738 inside an eye irradiated with 106Ru plagues. *Strahlenther Onkol* 2013;189:68-73. 739 **50**. Lommatzsch PK, Lommatzsch R. Treatment of juxtapapillary melanomas. *Br J* 740 Ophthalmol. 1991;75:715-717. 741 Freton A, Pavlick A, Finger PT. Systemic Evaluation and Management of Patients **51**.
- 742 with Uveal Melanoma. In: Book editor (Ryan S), Schachat A (Section editor) Retina. 743 Volume III, Tumors of the Retina, Choroid and Vitreous. London, New York, Oxford, 744 St. Louis, Sydney, Toronto: Elsevier; 2013, Chapter 149, pp. 2313-2315.
- 745 **52**. Kivelä T, Eskelin S, Kujala E. Metastatic uveal melanoma. Int Ophthalmol Clin 746 2006;46:133-149.
- 747 **53.** Merchant TE. Gould CI. Wilson MW. et al. Episcleral plaque brachytherapy for 748 retinoblastoma. Pediatr Blood Caner. 2004;43:134-139.
- 749 Shields JA, Shields CL, De Potter P, et al. Plaque radiotherapy for residual or **54**. 750 recurrent retinoblastoma in 91 cases. J Pediatr Ophthalmol Strab. 1994;31:242-245.
- 751 Stannard C, Maree G, Munro R, et al. Iodine-125 orbital brachytherapy with a **55.** 752 prosthetic implant in situ. Strahlenther Onkol. 2011;187:322-327.

753 **56.** Sealy R. Stannard C. Shackleton D. Improved cosmesis in retinoblastoma patients 754 treated with iodine-125 orbital irradiation. Ophthalmic Paediatr Genet. 1987;8:95-755 99. 756 **57.** Finger PT, Harbour JW, Karcioglu ZA. Risk factors for metastasis in retinoblastoma. 757 Surv Ophthalmol. 2002;47:1-16. Sastre X, Chantada GL, Doz F, et al. Proceedings of the consensus meetings from the 758 **58.** 759 International Retinoblastoma Staging Working Group on the pathology guidelines 760 for the examination of enucleated eves and evaluation of prognostic risk factors in 761 retinoblastoma. Arch Pathol Lab Med. 2009;133:1199-1202. 762 **59**. Abramson DH, Dunkel IJ, Brodie SE, et al. Superselective ophthalmic artery 763 chemotherapy as primary treatment for retinoblastoma (chemosurgery). 764 Ophthalmology. 2010;117:1623-1629. 765 **60**. Dimaras H, Kimani K, Dimba EA, et al. Retinoblastoma. Lancet. 2012;379:1436-1446. 766 **61**. The American Joint Committee on Cancer Ophthalmic Oncology Task Force. 767 Retinoblastoma. In: Edge S, Byrd, DR, Compton, CC, Fritz, AG, Greene, FL, Trotti A., ed. The AICC Cancer Staging Manual, 7th ed. New York: Springer: 2009, pp. 561-568. 768 769 **62**. The American Joint Committee on Cancer Ophthalmic Oncology Task Force. Uveal 770 Melanoma. In: Edge SE BD, Compton CA, Fritz AG, Greene FL, Troti A., ed. AJCC 771 Cancer Staging Manual 7th edition. 7th ed. New York; London: Springer; 2009, pp. 772 547-559. 773 63. Damato B, Patel I, Campbell IR, et al. Local tumor control after <sup>106</sup>Ru brachytherapy 774 of choroidal melanoma. Int J Radiat Oncol Biol Phys 2005;63:385-391.

Ouivev IM, Augsburger I, Snelling L, et al. 125I plaque therapy for uveal melanoma. 775 64. 776 Analysis of the impact of time and dose factors on local control. *Cancer*. 777 1996;77:2356-2362. 778 **65**. Finger P. Murphree A. Ophthalmic Brachytherapy: Treatment of Choroidal 779 Melanoma and Retinoblastoma. In: Peyman G, Meffert S, Conway M, Chou F, eds. 780 Vitreoretinal Surgical Techniques London: Martin Dunitz; 2006, pp. 452-468. 781 66. Kiratli H, Bilgic S, Atahan IL. Plaque radiotherapy in the management of retinoblastoma. Turk J Pediatr. 1998;40:393-397. 782 783 **67.** Temming P, Lohmann D, Bornfeld N, et al. Current concepts for diagnosis and 784 treatment of retinoblastoma in Germany: aiming for safe tumor control and vision 785 preservation. Klin Pediatr. 2012;224:339-347. 786 **68**. Harbour JW, Murray TG, Byrne SF, et al. Intraoperative echographic localization of 787 iodine 125 episcleral radioactive plaques for posterior uveal melanoma. Retina. 788 1996;16:129-134. 789 69. Chang MY, Kamrava M, Demanes DJ, et al. Intraoperative ultrasonography-guided 790 positioning of iodine-125 plaque brachytherapy in the treatment of choroidal 791 melanoma. Ophthalmology. 2012;119:1073-1077. 792 **70**. Finger PT. Reddy S. Chin K. High-frequency ultrasound characteristics of 24 iris and 793 iridociliary melanomas: before and after plaque brachytherapy. *Arch Ophthalmol.* 794 2007;125:1051-1058. 795 **71.** Finger PT. Finger's amniotic membrane buffer technique: protecting the cornea

during radiation plaque therapy. *Arch Ophthalmol.* 2008;126:531-534.

796

797 72. Finger PT. Chin KI. Yu GP. et al. Risk factors for cataract after palladium-103 798 ophthalmic plague radiation therapy. *Int J Radiat Oncol Biol Phys* 2010;80:800-806. 799 **73.** Kujala E, Mäkitie T, Kivelä T. Very long-term prognosis of patients with malignant 800 uveal melanoma. *Invest Ophthalmol Vis Sci.* 2003;44:4651-4659. 801 **74.** Hawkins BS, Collaborative Ocular Melanoma Study Group. The Collaborative Ocular 802 Melanoma Study (COMS) randomized trial of pre-enucleation radiation of large 803 choroidal melanoma: IV. Ten-vear mortality findings and prognostic factors. COMS 804 report number 24. Am I Ophthalmol. 2004:138:936-951. 805 **75.** Bechrakis NE, Hocht S, Martus P, et al. Endoresection following proton beam 806 irradiation of large uveal melanomas. Ophthalmologe, 2004:101:370-376. 807 **76.** Damato BE, Paul I, Foulds WS. Risk factors for residual and recurrent uveal 808 melanoma after trans-scleral local resection. Br J Ophthalmol. 1996;80:102-108. 809 77. Chan HS, DeBoer G, Thiessen JJ, et al. Combining cyclosporin with chemotherapy 810 controls intraocular retinoblastoma without requiring radiation. *Clin Cancer Res* 811 1996;2:1499-1508. 812 **78.** Bunin GR. Felice MA. Davidson W. et al. Medical radiation exposure and risk of 813 retinoblastoma resulting from new germline RB1 mutation. *Int I Cancer*. 814 2011:128:2393-2404. 815 79. Desiardins L, Chefchaouni MC, Lumbroso L, et al. [Functional results of 816 retinoblastoma treatment with local treatment used in isolation or associated with 817 chemotherapy]. I Fr Ophtalmol. 2005;28:725-731. 818 Gragoudas ES, Goitein M, Verhey L, et al. Proton beam irradiation of uveal 80. 819 melanomas. Results of 5 1/2-year study. Arch. Ophthalmol. 1982;100:928-934.

820 81. D'Hermies F. Meyer A. Morel X. et al. Neovascular glaucoma following proton-beam 821 therapy. Case report. I Fr Ophtalmol 2001;24:95-101. 822 **82.** Egger E, Zografos L, Schalenbourg A, et al. Eve retention after proton beam 823 radiotherapy for uveal melanoma. *Int J Radiat Oncol Biol Phys.* 2003;55:867-880. 824 83. Damato B, Kacperek A, Chopra M, et al. Proton beam radiotherapy of choroidal 825 melanoma: the Liverpool-Clatterbridge experience. *Int J Radiat Oncol Biol Phys* 826 2005;62:1405-1411. 827 **84**. Kim MK, Char DH, Castro IL, et al. Neovascular glaucoma after helium ion irradiation 828 for uveal melanoma. Ophthalmology. 1986;93:189-193. 829 **85**. Hungerford IL. Foss AI. Whelahan I. et al. Side effects of photon and proton 830 radiotherapy for ocular melanoma. Front Radiat Ther Oncol. 1997;30:287-293. 831 Cassoux N, Cayette S, Plancher C, et al. Choroidal Melanoma: Does Endoresection 86. 832 Prevent Neovascular Glaucoma in Patient Treated with Proton Beam Irradiation? 833 Retina. 2013;33:1441-1447. 834 **87.** Wilson MW, Hungerford JL. Comparison of episcleral plaque and proton beam 835 radiation therapy for the treatment of choroidal melanoma, Ophthalmology, 836 1999;106:1579-1587. 837 88. Boyd SR, Gittos A, Richter M, et al. Proton beam therapy and iris neovascularisation 838 in uveal melanoma. Eve. 2006;20:832-836. 839 Nakagawa Y, Yoshihara H, Kageji T, et al. Cost analysis of radiotherapy, carbon ion 89. 840 therapy, proton therapy and BNCT in Japan. *Appl Radiat Isot.* 2009;67:S80-83. 841 90. Anonymous. Proton therapy appears to be less cost-effective. *Cancer Discov.* 2013, 842 [ePub]

91. 843 Anzai K. Mori K. Yoneva S. Clinicopathological correlation of a retinal angiomatous 844 proliferation-like lesion in a case of radiation retinopathy. *Ipn J Ophthalmol*. 845 2006;50:291-293. 846 92. Archer DB, Amoaku WM, Gardiner TA. Radiation retinopathy--clinical, 847 histopathological, ultrastructural and experimental correlations. Eye. 1991;5:239-848 251. 849 93. Osman IM, Abouzeid H, Balmer A, et al. Modern cataract surgery for radiation-850 induced cataracts in retinoblastoma. *Br I Ophthalmol.* 2011;95:227-230. 851 94. Finger PT. Anti-VEGF bevacizumab (Avastin) for radiation optic neuropathy. Am I 852 Ophthalmol. 2007:143:335-338. 853 95. Arriola-Villalobos P, Donate-Lopez I, Calvo-Gonzalez C, et al. Intravitreal 854 bevacizumab (Avastin) for radiation retinopathy neovascularization. *Acta* 855 ophthalmol. 2008;86:115-116. 856 96. Finger PT. Radiation retinopathy is treatable with anti-vascular endothelial growth 857 factor bevacizumab (Avastin). Int J Radiat Oncol Biol Phys. 2008;70:974-977. 858 97. Gupta A. Muecke IS. Treatment of radiation maculopathy with intravitreal injection 859 of bevacizumab (Avastin). Retina. 2008;28:964-968. 860 98. Finger PT, Chin KJ. Intravitreous ranibizumab (lucentis) for radiation maculopathy. 861 Arch Ophthalmol. 2010;128:249-252. 862 Finger PT, Kurli M. Laser photocoagulation for radiation retinopathy after 99. 863 ophthalmic plaque radiation therapy. *Br J Ophthalmol.* 2005;89:730-738.

864	100.	Materin MA, Bianciotto CG, Wu C, et al. Sector laser photocoagulation for the
865		prevention of macular edema after plaque radiotherapy for uveal melanoma: a pilot
866		study. Retina. 2012;32:1601-1607.
867	101.	Finger PT. Radiation therapy for orbital tumors: concepts, current use, and
868		ophthalmic radiation side effects. Surv Ophthalmol. 2009;54:545-568.
869	102.	Radin PP, Lumbroso-Le Rouic L, Levy-Gabriel C, et al. Scleral necrosis after radiation
870		therapy for uveal melanomas: report of 23 cases. Graefe's Arch Clin Exp Ophthalmol
871		2008;246:1731-1736.
872	103.	Bechrakis NE, Bornfeld N, Zoller I, et al. Iodine-125 plaque brachytherapy versus
873		transcleral tumor resection in the treatment of large uveal melanomas.
874		Ophthalmology. 2002;109:1855-1861.
875		
876		
877		
077		
878		
879		
880		
881		
882		

883 884 885	Table Legends
886	Table 1: ABS-OOTF Levels of Consensus
887	ABS – OOTF = American Brachytherapy Society–Ophthalmic Oncology Task Force
888	Table 2: Changes in General Guidelines for Treatment of Uveal Melanoma
889	AJCC =7 <sup>th</sup> edition, American Joint Commission on Cancer <sup>62</sup>
890	* $^{106}$ Ru and $^{90}$ Sr plaques are less accommodating for nodular extrascleral extension.
891	Table 3: Radiobiological characteristics of radionuclides used for episcleral brachytherapy.
892	Photon emissions less than 5 KeV were removed from calculations of mean energy and tenth
893	value layers (TVLs)
894	Table 4: Comparison of Plaque and Proton Therapy
895	$^{103}$ Pd = palladium-103, $^{125}$ I = iodine-125, $^{106}$ Ru = ruthenium-106
896	Table 5: Review of Uveal Melanoma Clinical Case Series
897	Author = first author, Year = year of publication, Pts. = numbers of patients, Follow-up =
898	months, Dose = Gy, Tumor dimensions = mm, Local control and metastasis = % References
899	used.
900	Table 6: Classification for Radiation Retinopathy
901	DA = disc areas, OCT = optical coherence tomography, ICG = indocyanine green
902	angiography, FAG = fluorescein angiography, vision loss must be related to associated
903	sign(s). This table is modified from an original classification. <sup>99</sup>
904	
905	
906	

# The 2013 American Brachytherapy Society (ABS) Guidelines for Plaque Brachytherapy of Uveal Melanoma and Retinoblastoma

## The American Brachytherapy Society Ophthalmic Oncology Task Force

Running Title: ABS-00TF- ABS Plaque Brachytherapy Guidelines
Text Pages = 26
Reference Pages = 14
Tables = 6
Figures = 0
Abstract = 250 words
Key Words: brachytherapy, plaque, eye, melanoma, retinoblastoma, choroid
Corresponding author: Paul T. Finger MD, The New York Eye Cancer Center, Suite 5B, 115 East 61 <sup>st</sup> Street, New York City, New York, USA 10065. Tel: 1212-832-8170, E-Mail: <a href="mailto:pfinger@eyecancer.com">pfinger@eyecancer.com</a>

Table(s)

Table 1. ABS-OOTF Levels of Consensus

Level 1: Uniform panel consensus, evidence primarily from the published literature.

Level 2: Uniform panel consensus, based on clinical experience.

Level 3: No uniform panel consensus nor specific recommendation.

ABS – OOTF = American Brachytherapy Society–Ophthalmic Oncology Task Force

Table 2: Changes in General Guidelines for Treatment of Uveal Melanoma

2003 ABS Recommendations	2013 ABS Recommendations
Clinical diagnosis of uveal melanoma is adequate for treatment.  Histopathologic verification is not required.  Small melanomas may be treated if there is	Clinical diagnosis of uveal melanoma is adequate for treatment.  Histopathologic verification is not required.  Small melanomas can be treated at the eye cancer
evidence of growth.	specialist's discretion.
COMS medium and large uveal melanomas can be treated, after counseling about likely vision outcomes.	AJCC T1, T2, T3, T4a-d uveal melanoma patients can be treated, after counseling about likely vision, eye retention and local control outcomes.
Patients with peripapillary melanomas have poorer vision and local control outcomes and should be accordingly counseled.	Patients with peripapillary, subfoveal and those with exudative retinal detachments typically have poorer resultant vision and local control outcomes. They should be accordingly counseled.
Patients with gross extrascleral extension, ring melanoma and tumor involvement of half of the ciliary body are not suitable for plaque therapy.	Tumors with T4e extraocular extension,* basal diameters that exceed the limits of brachytherapy, blind painful eyes and those with no light perception vision are not suitable for plaque therapy.

AJCC = 7<sup>th</sup> edition, American Joint Commission on Cancer

<sup>\*</sup>  $^{106}\mathrm{Ru}$  and  $^{90}\mathrm{Sr}$  plaques are less accommodating for nodular extrascleral extension.

Table 3. Radiological characteristics of radionuclides used for episcleral brachytherapy. Photon emissions less than 5 keV were removed from calculations of mean energy and tenth-value layers (TVLs).

Photon emitters	<sup>1</sup> Half-life	<sup>1</sup> Mean photon energy (keV)	<sup>2</sup> Water TVL (mm)	<sup>3</sup> Pb TVL (mm)	
125 <b>I</b>	59.4 days	28.4	55	0.059	
<sup>103</sup> Pd	16.99 days	20.7	30	0.026	
131Cs	9.69 days	30.4	62	0.070	
			CSDA range in		
Beta emitters	Half-life	<sup>1</sup> Endpoint beta energy (MeV)	water (mm) 5		
106Ru/106Rh	371.8 days	3.541 4	17		
90 Sr	28.8 years	0.546 6	1.9		

<sup>1</sup> http://www.nndc.bnl.gov/chart/

http://www.alpharubicon.com/basicnbc/article16radiological71.htm

<sup>&</sup>lt;sup>2</sup> http://physics.nist.gov/PhysRefData/XrayMassCoef/ComTab/water.html

<sup>3</sup> http://physics.nist.gov/PhysRefData/XrayMassCoef/ElemTab/z82.html

<sup>4</sup> http://www.nndc.bnl.gov/chart/decaysearchdirect.jsp?nuc=106Rh&unc=nds

<sup>&</sup>lt;sup>5</sup> Handbook of Radioactivity Analysis, edited by M. F. L'Annunziata (2003) http://books.google.com/books?id=OfqdTC6deZkC&pg=PA19&lpg=PA19&dq=beta+particle+range+in+air&sou rce=bl&ots=D7gm8TeI3a&sig=zmcdrOUSI5NVqqfDl oPfOvhRCA&hl=en&ei=yN7MSfvZDprNlQfnqtXQCQ& sa=X&oi=book result&resnum=8&ct=result#v=onepage&q&f=false

<sup>6</sup> http://www.nndc.bnl.gov/chart/decaysearchdirect.jsp?nuc=106Rh&unc=nds

Table 4: Comparison of Plaque and Proton Therapy

PLAQUE	PROTON
Surgical Insertion and Removal	Surgical Clip Implantation
Continuous Low Dose Rate Treatment	4 Daily High Dose Rate Fractions
5 – 7 Day ( <sup>125</sup> I, <sup>103</sup> Pd)	
3-7 Day ( <sup>106</sup> Ru)	
Mobile Radiation Field	Static Radiation Field
Fewer Anterior Segment Complications	More Anterior Segment Complications
Posterior Segment Complications	Posterior Segment Complications
Less Expensive	More Expensive

 $^{103}\mathrm{Pd}$  = palladium-103,  $^{125}\mathrm{I}$  = iodine-125,  $^{106}\mathrm{Ru}$  = ruthenium-106

Table 5: Review of Uveal Melanoma Clinical Case Series

Author	Year	Pts.	Radionuclide	Follow up	Thickness	Basal Diameter	Radiation Dose	Local Control	Local Control	Metastasis	Metastasis	Visual Acuity
		No.		Mean or Median	Mean or Median	Mean or Median	Apex	Overall	5-year	Overall	5-year	Final %
				months	(range)	(range)	Mean	%				>20/200
Lommatzsch <sup>3</sup>	1987	309	ruthenium-106	80	3.7 (1.2-11.8)	9.7 (4.5-21.5)	100	69.9	84	12.9	NA	NA
Quivey <sup>64</sup>	1993	239	iodine-125	36	5.5 (1.9-11.0)	10.9 (4-18)	70	91.7	82	7.5	12	NA
Fontenesi <sup>15</sup>	1993	144	iodine-125	46	Small n=15, Medium n=84, Large n=45	NA	75	97.7	94.4	2.7	2	71.3
Seregard <sup>21</sup>	1997	266	ruthenium-106	43	4.4 (1.0-13.1)	10.0 (3-23)	100	83	82	11	14	NA
COMS 13	2001	657	iodine-125	96	4.8 (2.5-10.0)	11.4 (up to 16)	85	NA	NA	9	9	63
Bechrakis <sup>103</sup>	2002	152	iodine-125	30.1	9.0 ± 1.1	14.6 ± 2.4	98 ± 18	88.8	NA	11.1	NA	5.6
Shields 35	2002	354	iodine-125	60	9.0 (9.8-16)	14.0 (5-21)	80	91	91	24	24	43
Puusaari <sup>36</sup>	2003	97	iodine-125	43.2	10.7 (4.5-16.8)	16.1 (7.3-25)	87	94.8	94	28.9	35	42 at 1 year
Damato 83	2005	458	ruthenium-106	47	3.2 (0.7-7.0)	10.6 (5-16.6)	80	97	97.9	8.1	NA	57
Finger <sup>6</sup>	2009	400	palladium-103	51	3.8 (1.5-12.3)	10.5 (5-19.9)	73	97	NA	6	7.3	79
Mean	2003	341		52.9			84	91.9	91.2	14	17.9	48.3
Median	2002	354		47			82.5	92.9	92.5	11	14	48

Tumor measurements = milimeters , Followup = months, Pts. No. = number of patients, %<20/200 = those with better than 20/200 vision, NA = not available

Table 6: Classification for Radiation Retinopathy

Stage	Sign	Symptom	Location	Best viewed by	Risk of Vision Loss
1	Cotton-wool spots	None	Extramacular	Ophthalmoscopy	mild
	Retinal hemorrhages	None	Extramacular	Ophthalmoscopy	mild
	Retinal micro-aneurysms	None	Extramacular	Ophthalmoscopy / FAG	mild
	Exudate	None	Extramacular	Ophthalmoscopy	mild
	Uveal effusion	None	Extramacular	Ophthalmoscopy / OCT	mild
	Chorioretinal atrophy	None	Extramacular	Ophthalmoscopy	mild
	Choroidopathy	None	Extramacular	ICG	mild
	Retinal ischemia (< 5 DA)	None	Extramacular	FAG	mild
2	Above findings	None	Macular	Both	moderate
3	Any combination of the above plus				
	Retinal neovascularization	Vision loss	Extramacular	FAG	severe
	Macular edema - New onset	Vision loss	Macular	FAG / OCT	severe
4	Any combination of the above plus				
	Vitreous hemorrhage	Vision loss	Vitreous	Ophthalmoscopy	severe
	Retinal ischemia (>or = 5DA)	Vision loss	Both	FAG	severe

DA = disc areas, OCT = optical coherence tomography, ICG = indocyanine green angiography, FAG = fluorescein angiography,

vision loss must be related to associated sign(s). This table is modified and updated from an original classification. 99