

Adenoid cystic carcinoma of the lacrimal gland: role of nuclear survivin (BIRC5) as a prognostic marker

Kaustubh Mulay, Fairouz M Puthyapurayil,¹ Javed Ali Mohammad,¹ Mohammad Hasnat Ali,² Santosh G Honavar¹ & Vijay Anand P Reddy¹

L.V. Prasad Eye Institute, Ocular Pathology Service, ¹L.V. Prasad Eye Institute, Ocular Oncology Service, Hyderabad, India, and ²L.V. Prasad Eye Institute, Biostatistics Service, Hyderabad, India

Date of submission 25 August 2012
Accepted for publication 26 November 2012

Mulay K, Puthyapurayil F M, Mohammad J A, Ali M H, Honavar S G & Reddy V A P
(2013) *Histopathology* **62**, 840–846

Adenoid cystic carcinoma of the lacrimal gland: role of nuclear survivin (BIRC5) as a prognostic marker

Aims: This study aimed to evaluate the expression of nuclear survivin in adenoid cystic carcinoma (ACC) of the lacrimal gland and to determine if this expression is associated with histopathological features, markers of apoptosis and proliferation or clinical outcomes.

Methods and results: Immunohistochemical staining for survivin, p53, Ki-67 and Bcl-2 was analyzed in 55 cases of ACC of lacrimal gland. Thirty-one cases (56.3%) expressed nuclear survivin. All cases

expressed p53, Ki-67 and Bcl-2. Eleven cases (35.5%) had a high nuclear survivin score (NS-SCORE) and 20 cases (64.5%) had a low NS-SCORE. Cases with a high NS-SCORE had a shorter progression-free survival (PFS) ($P < 0.0001$), higher expression of Ki-67 ($P < 0.005$) and a solid tumour pattern $>30\%$ ($P < 0.005$).

Conclusion: Nuclear expression of survivin impacts prognosis significantly and is thus a promising prognostic marker in ACC of the lacrimal gland.

Keywords: adenoid cystic carcinoma, lacrimal gland, survivin (BIRC5)

Introduction

Tumours of the lacrimal gland are uncommon in clinical practice. They constitute approximately 7–9% of orbital tumours.^{1–3} Traditionally, it has been said that 50% of lacrimal masses are epithelial and 50% are non-epithelial.⁴ Epithelial tumours of the lacrimal gland account for 5–8% of all tumours at this site.^{5,6} Adenoid cystic carcinoma (ACC) is the most common malignant epithelial tumour of the lacrimal gland, with a deceptively benign histological appearance characterized by indolent, locally invasive growth with a high propensity for local recurrence and distant metastases. ACC accounts for 1.6% of all orbital tumours and 3.8% of all primary orbital tumours.⁵

Several clinicopathological features, such as advanced stage, solid histological subtype, high grade, the presence of perineurial invasion or a positive surgical margin, have been associated with recurrence or poor survival in ACC of the salivary gland.^{7–10} The primary treatment for ACC of the lacrimal gland has been *en-bloc* surgical excision, which is usually followed by postoperative radiotherapy.^{11,12} Neoadjuvant chemotherapy carries a risk of substantial toxicity.¹³

Cytotoxic agents have been associated with dismal antitumour activity in metastatic ACC of the salivary glands; this uniform lack of efficacy has prompted the evaluation of molecularly targeted therapies in these tumours. Unfortunately, their theoretical promise has yet to be fulfilled.¹³ Several biomarkers have been studied for their possible role as prognostic indicators of ACC in the lacrimal and salivary glands and as potential targets of molecular therapy.^{14–24} No validated protocols exist for ACC of the lacrimal glands.

Address for correspondence: Dr K Mulay MD, L.V. Prasad Eye Institute, Road No. 2, Banjara Hills, Hyderabad-500034, India.
e-mail: kaustubh.m@hotmail.com

Survivin, also called baculoviral inhibitor of apoptosis repeat-containing 5 (BIRC5), is a member of the inhibitor of apoptosis (IAP) family. Survivin has been studied extensively in various human cancers for its potential role as a prognostic marker. Studies have associated survivin expression with poor clinical outcome.^{25–28} It has also been evaluated for its prognostic role in ACC of the head and neck region.¹² However, its role in ACC of the lacrimal glands remains largely unknown. The purpose of this study was to determine whether nuclear survivin is expressed in ACC of the lacrimal gland and whether such expression has any association with various histopathological features, clinical outcomes of the disease, or markers of proliferation and apoptosis.

Materials and methods

We retrieved 65 cases of ACC involving the lacrimal gland diagnosed in the Ocular Pathology Service from 1995 to 2011. Ten cases were excluded from the study due to lack of follow-up information. Medical records were reviewed to evaluate clinical outcomes. All patients were assigned a pathological grade and a TNM stage according to the 7th edition of the American Joint Committee on Cancer (AJCC) staging system. A trained pathologist reviewed tumour tissue slides for histological pattern, histological grade, volume of solid component, perineurial invasion and status of the resected surgical margins. ACC was classified into three types based on the morphological growth pattern: cribriform, tubular and solid. The surgical resection margin was considered positive when residual tumour cells were found at the margin. The follow-up period ranged from 3 to 196 months (mean 51.8).

IMMUNOHISTOCHEMISTRY AND ANALYSIS

Serial 5-µm-thick tissue sections cut from paraffin blocks were mounted individually onto gelatine-coated glass slides. Sections were deparaffinized and rehydrated. Endogenous peroxidase activity was blocked by incubating the slides in 0.3% solution of hydrogen peroxide in methanol for 30 min. Microwave antigen retrieval was performed, for which tissues were immersed in Coplin jars containing 0.01 M phosphate-buffered saline, pH 7.4, and subjected to intermittent heating for three cycles of 5 min each in a 625W microwave oven to maintain the temperature of the buffer at approximately 95°C. After cooling the slides to room temperature, non-specific

immunoreactivity was blocked by incubation with normal horse serum for 15 min (three cycles of 5 min each) at room temperature. Sections were rinsed in running tap water and phosphate-buffered saline (2 min each) at room temperature, and then stained immunohistochemically using commercially available monoclonal antibodies, with diaminobenzidine as the chromogen.

The following primary antibodies were used in the study: Ki-67 (clone Ki88), p53 (clone BP53-12-1), survivin (clone EP2880Y) and Bcl-2 (clone 100) (all from BioGenex, Fremont, CA, USA). Positive and negative controls were run simultaneously with every batch. Positive immunostaining for Ki-67, survivin and p53 was defined as brown staining confined to the nucleus, and for Bcl-2 as cytoplasmic/membranous staining (Figure 1).

NUCLEAR SURVIVIN SCORE (NS-SCORE)

Nuclear expression for survivin was scored in terms of intensity and percentage expression. Intensity was scored as zero (no staining), 1 (weak), 2 (mild), 3 (moderate) and 4 (strong). Percentage of staining was scored as 0 (0%), 1 (1–25%), 2 (26–50%), 3 (51–75%) and 4 (76–100%). Intensity and percentage scores were then summed to calculate the NS-SCORE. Tumours with NS-SCORES of 0–4 were grouped as low-score tumours, while those with NS-SCORES of 5–8 were scored as high-score tumours. NS-SCORES were correlated with histopathological features, clinical outcomes and expression of other markers.

STATISTICAL ANALYSIS

R (version 2.14.1) software was used for the statistical analysis. Progression-free survival (PFS) was calculated from the date of diagnosis to the date when the tumour recurred or the patient died. Survival was derived using the Kaplan–Meier method. The log-rank test was used to evaluate the statistical differences in the cumulative survival scores. The Cox model was used for the multivariate analysis. A 95% confidence interval (CI) was used for survival rates and hazard ratio.

ETHICS STATEMENT

This study was reviewed by the ethics committee of L.V.Prasad Eye Institute, Hyderabad, and was performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki. Informed consent was obtained from the patients.




Figure 1. Immunohistochemical analysis of adenoid cystic carcinoma. Nuclear immunoreactivity for p53 (A), nuclear immunoreactivity for survivin (B), membranous/cytoplasmic immunoreactivity for Bcl-2 (C) and nuclear immunoreactivity for Ki-67 (D).

Results

CLINICOPATHOLOGICAL FEATURES

Table 1 depicts the clinicopathological features of the 55 cases of ACC of the lacrimal gland.

EXPRESSION OF KI-67, P53 AND BCL-2

All cases expressed p53, Ki-67 and Bcl-2. The proportion of tumour cells expressing Bcl-2, Ki-67 and p53 varied from 54% to 99%, 3.2% to 32% and 51% to 88%, respectively.

EXPRESSION OF SURVIVIN

We categorized the tumours into two groups: the 'nuclear' group, which included all tumours where survivin was expressed in the nucleus of tumour cells

with/without cytoplasmic immunoreactivity for surviving, and the 'non-nuclear' group, where survivin expression was limited only to the cytoplasm or the tumour cells failed to express survivin either in the cytoplasm or the nucleus. Of the 55 cases, 31 cases (56.4%) expressed survivin in the nucleus of tumour cells. Table 2 shows the comparative analysis of the nuclear and non-nuclear groups. Tumours expressing nuclear survivin had a higher Ki-67 index ($P < 0.0001$), greater p53 expression ($P < 0.0001$), a greater volume showing a solid pattern ($P = 0.0015$) and were larger ($P < 0.0001$). Also, the recurrence rate was significantly higher ($P < 0.001$) and the PFS shorter ($P < 0.0001$) for patients whose tumours expressed nuclear survivin compared to those whose tumours did not. No significant association was seen between Bcl-2 expression and nuclear expression of survivin.

Table 1. Clinicopathological features of Adenoid Cystic Carcinoma of the lacrimal gland

Feature	No. of patients	Percentage (%)
Total	55	
Age in years, median (range)	35 (11–77)	
Gender		
Male	24	43.63
Female	31	56.36
Lymph node involvement	01	1.81
Tumour size		
≤2cm	31	56.36
>2cm	24	43.63
Histological growth pattern (Predominant)		
Cribriform	40	72.7
Tubular	09	16.36
Solid	06	10.9
Solid component		
≤30%	42	76.36
>30%	13	23.63
Histological grade		
Gx	0	0
G1	33	60.0
G2	9	16.36
G3	7	12.72
G4	6	10.9
Perineurial invasion		
Present	47	85.45
Absent	08	14.54
Resected surgical margins		
Involved	04	7.27
Not involved	51	92.72

EVALUATION OF THE NS-SCORES

Eleven cases (35.4%) had a high NS-SCORE, while 20 cases (65.4%) had a low NS-SCORE. Table 3 shows the comparative analysis of the high and low NS-SCORE groups. Cases with a high NS-SCORE had a shorter PFS ($P < 0.0001$), higher expression of Ki-67

($P < 0.005$) and a greater proportion of solid tumour pattern ($P < 0.005$).

Figure 2 shows the Kaplan–Meier plots for survival analysis between the nuclear/non-nuclear and the high/low NS-SCORE groups. The reproducibility of the study was acceptable, with least-square regression analysis resulting in an R^2 -value of 0.93.

Discussion

Adenoid cystic carcinoma (ACC) is a rare epithelial tumour of the lacrimal gland, most common in the 5th and 6th decades of life, but can appear at virtually any age ranging from 6.5 to 79 years.^{29,30} In the present study, the age at presentation ranged from 11 to 77 years. Gender predisposition is an inconsistent feature in the literature, with authors variedly reporting male preponderance, female preponderance, and no gender predilection.^{31,32} In the present study, the male:female ratio was 1:1.29. However, this difference was not statistically significant. Studies in the literature have correlated tumour size with prognosis.^{12,33} A study on ACC of the head and neck found that all tumours >4 cm had an unfavourable prognosis.³³ In the present study, tumours >2 cm showed a trend for a shorter PFS but this difference was not statistically significant.

ACC can be classified morphologically into three prognostically significant patterns: cribriform, tubular and solid.³⁴ The presence of a solid component has been a consistent predictor of poor prognosis in several series.^{7,35–37} In the present study, we found that cases with a solid pattern involving >30% of the tumour had a shorter PFS ($P < 0.0001$). The AJCC tumour stage is more predictive of prognosis and distant metastasis.^{7,31} Wright *et al.*²⁹ have reported a greater frequency of recurrence in patients with ACC with a solid pattern accounting for more than half the tumour, compared to those where this pattern was not evident. Although we did not find a statistically significant association between solid tumour pattern and recurrence, such tumours showed a trend for a greater recurrence rate. In a recent review of ACC, 74.3% of patients showed advanced disease at the time of initial evaluation.³⁸ Despite this, cervical node metastases are a rare event. Lung, bone and liver are common metastatic sites.^{32,39} Two patients in the present study had lung metastases in the follow-up period.

Survivin, or BIRC5, is a protein encoded by the *BIRC5* gene⁴⁰ and is a member of the IAP family. The survivin protein functions to inhibit caspase

Table 2. Comparative analysis between nuclear and non-nuclear groups for survivin

Feature	Nuclear group	Non-nuclear group	P-value
Age at presentation (years)	11–77 (Median 35)	11–72 (Median 35.5)	0.388
Predominant solid morphologic pattern	16.1% (5/31)	4.16% (1/24)	0.216
Solid component >30%	41.9% (13/31)	4.16% (1/24)	0.0015
p53 (% expression)	65–87 (Median 82)	51–88 (Median 67)	<0.0001
Bcl-2 (% expression)	54–99 (Median 92)	72–98 (Median 95)	0.353
Ki 67 index	5.2–32.0 (Median 25.2)	3.2–26.3 (Median 5.75)	<0.0001
Recurrence	74.2 (23/31)	4.2 (1/24)	<0.001
PFS (Months)	3–120 (31)	40–196 (Median 91.5)	<0.0001
Size (>2 cm)	87.09% (27/31)	16.6% (4/24)	<0.0001

Table 3. Comparative analysis of low and high NS-SCORE groups

Feature	High SSCORE	Low SSCORE	P-value
Age at presentation (years)	11–55 (Median 32)	11–77 (Median 36.5)	0.342
Predominant solid morphologic pattern	9.09% (1/11)	20% (4/20)	0.624
Solid component >30%	90.9% (10/11)	15% (3/20)	<0.005
p53 (% expression)	73–87 (Median 83)	85–87 (Median 82)	0.451
Bcl-2 (% expression)	54–99 (Median 93)	73–99 (Median 92)	0.408
Ki 67 index	23.4–32.0 (Median 29.1)	5.2–26.7 (Median 24.05)	<0.005
Recurrence	100% (11/11)	60% (12/20)	0.265
PFS (Months)	3–20 (Median 13)	12–120 (Median 40)	<0.0001
Size (>2 cm)	100% (11/11)	80% (16/20)	0.269

activation, thereby leading to negative regulation of apoptosis or programmed cell death. It is expressed highly in most human tumours and fetal tissue, but is completely absent in terminally differentiated cells.⁴¹ The single survivin gene can give rise to four different alternatively spliced transcripts: survivin-2A, survivin-2B, survivin-delta-ex-3 and survivin-3B.⁴² Survivin-2A and survivin-3B are predicted to be truncated forms of 120 and 128 amino acids, respectively, while survivin-2B and survivin-delta-ex-3 result from alternative splicing at the interface between exons 2 and 3.⁴³ The presence of transcripts of survivin has been correlated with cancer progression with differing conclusions.^{42,44–46} Reports of the localization of survivin and its relationship with prognosis are also conflicting. Survivin-2B by itself is localized to both nuclear and cytoplasmic compartments,

whereas survivin-delta-ex-3 is localized only to the nucleus. The localization of these three variants (survivin-2A, survivin-2B, and survivin-delta-ex-3) differ, however, when co-transfected together rather than individually.⁴² A Western blot and immunocytochemistry analysis has confirmed that survivin could be detected both in the nucleus and the cytoplasm. Ponnelle *et al.*⁴⁷ detected survivin both in the cytoplasm and the nucleus of colorectal carcinoma, and also found a significant association between cytoplasmic expression and better prognosis. In a recent study, PSORT II analysis predicted a preferential cytoplasmic localization of survivin-2A and survivin-2B, but a preferential nuclear localization of survivin-delta-ex-3.⁴⁴ Liao *et al.*⁴⁸ found survivin expression in ACC of the lacrimal gland to be localized to the cytoplasm exclusively, and its expression was related to




Figure 2. Kaplan–Meier survival plots for progression-free survival of patients with adenoid cystic carcinoma according to location of survivin expression (left) and nuclear survivin score (NS-SCORE) (right).

the T stage and proliferative index. They also documented that, of all the IAP proteins included in their study, survivin was the only factor that could predict prognosis independently. Ko *et al.*¹² concluded that expression of nuclear survivin has clinicopathological implications in patients with ACC of the head and neck region. In the present study, nuclear expression of survivin was associated with an increased recurrence rate, shorter PFS, higher Ki-67 index and higher p53 expression (Table 2). Also, tumours expressing nuclear survivin were larger, and had a greater proportion showing a solid tumour pattern, thus relating nuclear survivin to histological grade (G) and stage (T) of the tumour.

In the present study we scored the expression of nuclear survivin. This, to our knowledge, is the first time that a scoring system has been used to evaluate the expression of survivin in ACC of the lacrimal glands. Patients with tumours having a high NS-SCORE had a shorter PFS, and these tumours were more likely to show higher Ki-67 expression and a solid pattern involving >30% (Table 3).

Conclusion

Nuclear expression of survivin in ACC of the lacrimal gland correlates with histopathological indicators of poor outcome, such as the proportion of the solid component and tumour stage. It also correlates with high expression of markers of apoptosis and proliferation. Overall, nuclear expression of survivin impacts prognosis significantly and is thus a promising prognostic marker in ACC of the lacrimal gland. The novel concepts incorporated within our study may

provide newer directions for further research into the management of ACC of the lacrimal gland.

Conflicts of interest

None.

References

1. Goder GJ. Tumours of lacrimal gland. *Orbit* 1982; 12: 91–96.
2. Shields JA, Shields CL, Scartozzi R. Survey of 1264 patients with orbital tumors and simulating lesions: The 2002 Montgomery Lecture, part 1. *Ophthalmology* 2004; 111: 997–1008.
3. Santos RR, Damasceno RW, de Pontes FS *et al.* Ten-year follow-up of a case series of primary epithelial neoplasms of the lacrimal gland: clinical features, surgical treatment and histopathological findings. *Arq. Bras. Ophthalmol.* 2010; 73: 33–39.
4. Reese AB. Expanding lesions of the orbit. *Trans. Ophthalmol. Soc. UK* 1971; 91: 85–104.
5. Henderson JW. *Orbital tumours*. 3rd ed. New York: Raven Press, 1994; 323–342.
6. Shinder R, Al-Zubidi N, Esmali B. Survey of orbital tumors at a comprehensive cancer center in the United States. *Head Neck* 2011; 33: 610–614.
7. Khan AJ, DiGiovanna MP, Ross DA *et al.* Adenoid cystic carcinoma: a retrospective clinical review. *Int. J. Cancer* 2001; 96: 149–158.
8. Spiro RH, Huvos AG. Stage means more than grade in adenoid cystic carcinoma. *Am. J. Surg.* 1992; 164: 623–628.
9. Le QT, Birdwell S, Terris DJ *et al.* Post operative irradiation of minor salivary gland malignancies of the head and neck. *Radiother. Oncol.* 1999; 52: 165–171.
10. Prokopakis EP, Snyderman CH, Hanna EY, Carrau RL, Jhonson JT, D'Amico F. Risk factors for local recurrence of adenoid cystic carcinoma: the role of post operative radiation therapy. *Am. J. Otolaryngol.* 1999; 20: 281–286.
11. Garden AS, Weber RS, Morrison WH *et al.* The influence of positive margins and nerve invasion in adenoid cystic carcinoma.

- noma of the head and neck treated with surgery and radiation. *Int. J. Radiat. Oncol. Biol. Phys.* 1995; **32**: 619–626.
12. Ko YH, Roh SY, Won HS *et al.* Prognostic significance of nuclear survivin expression in resected adenoid cystic carcinoma of the head and neck. *Head Neck Oncol.* 2010; **2**: 30–38.
 13. Le Tourneau C, Razak AR, Levy C *et al.* Role of chemotherapy and molecularly targeted agents in the treatment of adenoid cystic carcinoma of the lacrimal gland. *Br. J. Ophthalmol.* 2011; **95**: 1483–1489.
 14. Costa AF, Altemani A, Vékony H *et al.* Genetic profile of adenoid cystic carcinomas with high-grade transformation versus solid type. *Cell Oncol.* 2011; **34**: 369–379.
 15. Falbo V, Floridia G, Censi F, Marra M, Foschini MP, Taruscio D. Three cases of rare salivary gland tumours: a molecular study of TP53, CDKN2A/ARF, RAS, BRAF, PTEN, MAPK2 and EGFR genes. *Oncol. Rep.* 2011; **26**: 3–11.
 16. Seethala RR, Cieply K, Barnes EL, Dacic S. Progressive genetic alterations of adenoid cystic carcinoma with high-grade transformation. *Arch. Pathol. Lab. Med.* 2011; **135**: 123–130.
 17. Al-Rawi NH, Omer H, Al Kawas S. Immunohistochemical analysis of p53 and bcl-2 in benign and malignant salivary glands tumors. *J. Oral Pathol. Med.* 2010; **39**: 48–55.
 18. de Lima MD, Marques YM, Alves SM Jr *et al.* MDM2, P53, P21WAF1 and pAKT protein levels in genesis and behaviour of adenoid cystic carcinoma. *Cancer Epidemiol.* 2009; **33**: 142–146.
 19. Yamazaki M, Fujii S, Murata Y, Hayashi R, Ochiai A. High expression level of geminin predicts a poor clinical outcome in salivary gland carcinomas. *Histopathology* 2010; **56**: 883–892.
 20. Ettl T, Schwarz S, Kleinsasser N, Hartmann A, Reichert TE, Driemel O. Overexpression of EGFR and absence of C-KIT expression correlate with poor prognosis in salivary gland carcinomas. *Histopathology* 2008; **53**: 567–577.
 21. Suzzi MV, Alessi A, Bertarelli C *et al.* Prognostic relevance of cell proliferation in major salivary gland carcinomas. *Acta Otorhinolaryngol. Ital.* 2005; **25**: 161–168.
 22. Marques YM, Giudice FS, Freitas VM *et al.* Oestrogen receptor β in adenoid cystic carcinoma of salivary glands. *Histopathology* 2012; **60**: 609–616.
 23. Dori S, Trougouboff P, David R, Buchner A. Immunohistochemical evaluation of estrogen and progesterone receptors in adenoid cystic carcinoma of salivary gland origin. *Oral Oncol.* 2000; **36**: 450–453.
 24. Defaud-Hénon F, Tunon-de-Lara C, Fournier M *et al.* Adenoid cystic carcinoma of the breast: clinical, histological and immunohistochemical characterization. *Ann. Pathol.* 2010; **30**: 7–16.
 25. Marioni G, Agostini M, Bedin C *et al.* Survivin and laryngeal carcinoma prognosis: nuclear localization and expression of splice variants. *Histopathology* 2012; **61**: 247–256.
 26. Hui MK, Lai KK, Chan KW *et al.* Clinical correlation of nuclear survivin in esophageal squamous cell carcinoma. *Med. Oncol.* 2012; **29**: 3009–3016.
 27. Adamkov M, Kajo K, Vybohova D. Correlations of survivin expression with clinicomorphological parameters and hormonal receptor status in breast ductal carcinoma. *Neoplasm* 2012; **59**: 30–37.
 28. Ekeblad S, Lejonklou MH, Stålberg P, Skogsied B. Prognostic relevance of survivin in pancreatic endocrine tumors. *World J. Surg.* 2012; **36**: 1411–1418.
 29. Wright JE, Rose GE, Garner A. Primary malignant neoplasms of the lacrimal gland. *Br. J. Ophthalmol.* 1992; **76**: 401–407.
 30. Tellado MV, McLean IW, Specht CS, Varga J. Adenoid cystic carcinomas of the lacrimal gland in childhood and adolescence. *Ophthalmology* 1997; **104**: 1622–1625.
 31. Kokemueller H, Eckardt A, Brachvogel P, Hausamen JE. Adenoid cystic carcinoma of the head and neck – a 20 years experience. *Int. J. Oral Maxillofac. Surg.* 2004; **33**: 25–31.
 32. Da Cruz Perez DE, de Abreu Alves F, Nobuko Nishimoto I, de Almeida OP, Kowalski LP. Prognostic factors in head and neck adenoid cystic carcinoma. *Oral Oncol.* 2006; **42**: 139–146.
 33. Hamper K, Lazar F, Dietel M *et al.* Prognostic factors for adenoid cystic carcinoma of the head and neck: a retrospective evaluation of 96 cases. *J. Oral Pathol. Med.* 1990; **19**: 101–107.
 34. Jaso J, Malhotra R. Adenoid cystic carcinoma. *Arch. Pathol. Lab. Med.* 2011; **135**: 511–515.
 35. Fordice J, Kershaw C, El-Naggar A, Goepfert H. Adenoid cystic carcinoma of the head and neck: predictors of morbidity and mortality. *Arch. Otolaryngol. Head Neck Surg.* 1999; **125**: 149–152.
 36. Seethala RR. Histologic grading and prognostic biomarkers in salivary gland carcinomas. *Adv. Anat. Pathol.* 2011; **18**: 29–45.
 37. Zushi Y, Noguchi K, Hashitani S *et al.* Relations among expression of CXCR4, histological patterns, and metastatic potential in adenoid cystic carcinoma of the head and neck. *Int. J. Oncol.* 2008; **33**: 1133–1139.
 38. Wiseman SM, Popat SR, Rigual NR *et al.* Adenoid cystic carcinomas of the paranasal sinuses or nasal cavity: a 40 year review of 35 cases. *Ear Nose Throat J.* 2002; **81**: 510–514.
 39. Matsuba HM, Thawley SE, Simpson JR, Levine LA, Mauney M. Adenoid cystic carcinoma of major and minor salivary gland origin. *Laryngoscope* 1984; **94**: 1316–1318.
 40. Altieri DC. Molecular cloning of effector cell protease receptor-1, a novel cell surface receptor for the protease factor Xa. *J. Biol. Chem.* 1994; **269**: 3139–3142.
 41. Kuo P-C, Lui H-F, Chao J-I. Survivin and p53 modulate Quercetin-induced cell growth inhibition and apoptosis in human lung carcinoma cells. *J. Biol. Chem.* 2004; **279**: 55875–55885.
 42. Caldas H, Jiang Y, Holloway MP *et al.* Survivin splice variants regulate the balance between proliferation and cell death. *Oncogene* 2005; **24**: 1994–2007.
 43. Noton EA, Colnaghi R, Tate S *et al.* Molecular analysis of survivin isoforms. Evidence that alternatively spliced variants do not play a role in mitosis. *J. Biol. Chem.* 2006; **281**: 1286–12895.
 44. Mahotka C, Liebmann J, Wenzel M *et al.* Differential subcellular localization of functionally divergent survivin splice variants. *Cell Death Differ.* 2002; **9**: 1334–1342.
 45. Krieg A, Mahotka C, Krieg T *et al.* Expression of different survivin variants in gastric carcinomas: first clues to a role of survivin-2B in tumour progression. *Br. J. Cancer* 2002; **86**: 737–743.
 46. Li F. Role of survivin and its splice variants in tumorigenesis. *Br. J. Cancer* 2005; **92**: 212–216.
 47. Ponnelle T, Chapusot C, Martin L *et al.* Cellular localization of survivin: impact on the prognosis in colorectal cancer. *J. Cancer Res. Clin. Oncol.* 2005; **131**: 504–510.
 48. Liao Y, Zeng H, Wang X *et al.* Expression patterns and prognostic significance of inhibitor of apoptosis proteins in adenoid cystic carcinoma and pleomorphic adenoma of the lachrymal gland. *Exp. Eye Res.* 2009; **88**: 4–11.