



 [Print this Page for Your Records](#)

[Close Window](#)

Control/Tracking Number: 12-A-545-SNO

Activity: Abstract

Current Date/Time: 5/15/2012 12:28:51 PM

Stereotactic body radiotherapy of spinal metastases using Novalis Tx system: plan comparison of intensity modulated radiotherapy (IMRT) and volumetric modulated arc therapy (VMAT).

Author Block: Vijay Anand Reddy Palkonda, Niteen More, Padma Ganesan, Revathy Kesavan, Muthulingam Shunmugavel, Tamilselvan Kasirajan, Vinitha Reddy Maram, Sajal Kakkar, Prashant Upadhyay, Suman Das, Shantling Nigudgi, Apollo Hospitals, Hyderabad, India

Abstract:

INTRODUCTION: Intensity modulated radiotherapy is routinely used for stereotactic body radiotherapy of spinal metastases. Volumetric modulated arc therapy is the new technique that needs to be compared with IMRT. **METHODS:** Five patients of spinal metastases treated with stereotactic body radiotherapy were the study population. CT scans were obtained at 1mm slice thickness after immobilizing patients in thermoplastic masks. Volumetric MRI was performed and registered with CT images using BrainLab iPlan RT Image software. Vertebral bodies and pedicles were drawn as clinical target volume (CTV) and planning target volume (PTV) was generated after expanding CTV by 1mm. Spinal cord was drawn at the level of involved vertebra with cranial and caudal extension of 6mm. Three patients were planned for 30Gy and two for 35Gy divided in 5 daily fractions. Planning was performed on BrainLab iPlan RT dose version 4.1.2 workstation with static IMRT beams. Treatment was delivered on Novalis Tx system with ExacTrac image verification. For dosimetric comparison images with contours were transferred to Eclipse planning workstation version 8.6 and single arc VMAT plans were generated. **RESULTS:** Mean volume of the PTV was 33.1cc (21.5-49.5cc). Mean dose to PTV was 99.96 \pm 1.29 with IMRT and 100.4% \pm 0.89 with VMAT plans. Mean of conformity index was 1.31 \pm 0.12 in IMRT and 1.18 \pm 0.1 VMAT plans. Mean of maximum dose (0.035cc) to spinal cord with IMRT and VMAT plans was 27.33Gy and 26.43Gy, respectively. Ten percentage of spinal cord was receiving 26.36Gy in IMRT plans and 25.48Gy in VMAT plans. **CONCLUSION:** Conformity index was better with high-definition VMAT compared to IMRT plans. There was no significant dosimetric advantage of single arc VMAT over IMRT in sparing spinal cord. Treatment delivery time would be shorter with VMAT technique possibly reducing intrafraction variations.

:

Disclosures & Release (Complete):

*: My presentation will **NOT** include discussion of investigational or off-label use of a product.

*: I agree my presentation will follow the guidelines stated above.

*: I agree to statement above

Educational Objectives (Complete):*** Educational Objective 1**

: 7. Use advances in pharmacology, experimental therapeutics, biologic therapies, and radiobiology to improve future therapies for patients with CNS tumors

Category (Complete): CLIN-Radiation Therapy

Status: Complete

*****To log out, simply close your browser window. All information will be saved if you have clicked the "Save and Continue" button after each step****

Society for Neuro-Oncology

4617 Birch Street

Bellaire, TX 77401-5509

Technical Support: OASIS Helpdesk

Phone: (217)398-1792 (Mon. - Fri. 9 am - 5 pm CST).

Powered by OASIS, The Online Abstract Submission and Invitation System SM

© 1996 - 2012 Coe-Truman Technologies, Inc. All rights reserved.