

Disrupted development and altered hormone signaling in male *Padi2/Padi4* double knockout mice

Kelly L. Sams¹, Chinatsu Mukai¹, Brooke A. Marks¹, Chitvan Mittal¹, Elena Alina Demeter², Sophie Nelissen², Jennifer K. Grenier^{2,3}, Ann E. Tate^{2,3}, Faraz Ahmed^{2,3}, Scott A. Coonrod^{1,2*}

¹ Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University

² Department of Biomedical Sciences, College of Veterinary Medicine, Cornell University

³ Transcriptional Regulation and Expression Facility, Department of Biomedical Sciences, Cornell University

*Correspondence: sac269@cornell.edu

ABSTRACT

Background: Peptidylarginine deiminase enzymes (PADs) convert arginine residues to citrulline in a process called citrullination or deimination. Recently, two PADs, PAD2 and PAD4, have been linked to hormone signaling *in vitro* and the goal of this study was to test for links between PAD2/PAD4 and hormone signaling *in vivo*.

Methods: Preliminary analysis of *Padi2* and *Padi4* single knockout (SKO) mice did not find any overt reproductive defects and we predicted that this was likely due to genetic compensation. To test this hypothesis, we created a *Padi2/Padi4* double knockout (DKO) mouse model and tested these mice for a range of reproductive defects.

Results: Controlled breeding trials found that DKO breeding pairs, particularly males, appeared to take longer to have their first litter than wild-type FVB controls (WT), and that pups and DKO male weanlings weighed significantly less than their WT

counterparts. Additionally, DKO males took significantly longer than WT males to reach puberty and had lower serum testosterone levels. Furthermore, DKO males had smaller testes than WT males with increased rates of germ cell apoptosis.

Conclusions: The *Padi2/Padi4* DKO mouse model provides a new tool for investigating PAD function and outcomes from our studies provide the first *in vivo* evidence linking PADs with hormone signaling.

KEY WORDS: PAD2, PAD4, citrullination, hormone signaling, estrogen receptor, androgen receptor, delayed puberty

BACKGROUND

Peptidylarginine deiminases (PADs or PADIs) are a family of calcium-dependent enzymes that post-translationally convert positively-charged arginine residues to neutrally-charged citrulline in a process called citrullination or deimination. The loss of charge on target arginine residues following citrullination can significantly alter the target protein's tertiary structure and affect protein-protein and protein-DNA/RNA interactions (1). There are five PAD family members (PAD1-4 and 6), with each of the isoforms showing some degree of substrate and tissue specificity (2). Functional roles for PADs in mammalian physiology and pathology are diverse and include cellular differentiation, nerve growth, apoptosis, inflammation, early embryonic development, and gene regulation (3).

While PAD activity is most closely associated with autoimmune disease (4), previous studies have found that PAD2 and PAD4 are expressed in reproductive epithelial

tissues and that their expression is regulated by steroid hormone signaling (5,6). Steroid hormones are essential mediators of many physiological events including reproduction and development (7). These hormones carry out their effects by binding to their cognate nuclear receptors which, in turn, bind to hormone response DNA elements and regulate transcription via interaction with a wide range of co-regulators and the basal transcriptional machinery. Given the potential links between PADs and steroid hormones and that histones were found to be targeted by PAD4 for citrullination (8), we previously tested whether PAD2 and PAD4 may play a role in estrogen signaling in breast cancer cells by facilitating ER target gene expression via histone citrullination at ER binding sites. Results from our studies found that PAD4 is recruited to the TFF1 promoter (a canonical ER α binding site) in MCF7 cells following E2 treatment leading to citrullination of histone H4 arginine 3 (9). Additionally, we found that PAD2 interacts with ER in an estrogen-dependent manner and that estrogen (E2) treatment activates PAD2 leading to citrullination of histone H3 arginine 26 (H3Cit26) at ER binding sites (10,11). We then demonstrated by ChIP-Seq that ~95% of ER binding sites throughout the genome are citrullinated at H3Cit26 within 5-10 minutes of E2 treatment (12). We also found that PAD inhibitors potently blocked ER binding at Estrogen Response Elements (EREs) and prevented ER target gene activation (11).

Regarding a potential role for PADs in hormone signaling in males, another group recently found that PAD2 plays a role in androgen receptor (AR) signaling by facilitating nuclear import of AR and promoting AR target gene activation via histone citrullination at AR binding sites (10). This finding suggests that the role of PADs in regulating hormone-mediated transcription may be broader than we had previously realized.

Importantly, to date, there have not been any published reports linking PADs to hormone signaling in animal models and this lack of *in vivo* data has likely limited research into this area.

To address this gap in knowledge, we began testing for links between PADs and hormone signaling *in vivo* by determining whether *Padi2* and *Padi4* single knockout (SKO) mouse lines displayed overt reproductive defects. Initial studies with these mice found that both PAD2 and PAD4 mice are largely normal with no clear reproductive defects (data not shown). Given the greater than 59% amino acid homology between PAD family members and that *Padi* genes are closely clustered together at chromosome 1p35-36 in humans and chromosome 4E1 in mice (3,13), we predicted that the lack of a phenotype in the SKO lines was due to genetic compensation by other *Padis*. In order to test this hypothesis, we generated a *Padi2/Padi4* double knockout mouse (DKO) line by deleting *Padi4* from our existing *Padi2* knockout (P2KO) mouse line using CRISPR/CAS9 technology. We demonstrate that, as opposed to the SKO lines, the *Padi2/Padi4* DKO mice display a range of reproductive defects, particularly in males. Outcomes from our studies show that DKO males displayed a reduced pup and weanling weight, delayed puberty, reduced testis size, increased rates of apoptosis during spermatogenesis, and altered circulating hormone levels. Our results provide the first *in vivo* evidence that PADs play a role in male reproduction, potentially via their role in hormone signaling.

MATERIALS AND METHODS

Animals

All mice used in this work were on an FVB/NJ background and were maintained as inbred homozygous lines. The mice were housed in an AAALAC certified facility, with 12 hours on/off light cycle, temperature control and food and water ad libitum. All procedures were approved by Cornell University's Animal Care and Use Committee (IACUC) and in compliance with the US National Research Council's Guide for the Care and Use of Laboratory Animals (14). Breedings were recorded and lineages tracked with Softmouse Colony Management Software (15). Euthanasia was performed by CO₂ inhalation in accordance with American Veterinary Medical Association approved practice.

Generation of transgenic single knockout mice

Padi2-null mice (FVB/NJ background) were generated using gene trap technology at the Texas A&M Institute for Genomic Medicine (16). For the gene trap, 163 nucleotides of exon 1 in *Padi2* (from the ATG initial codon to intron 1-2) were replaced by the LacZ and Neomycin coding sequence. These mice are genotyped in a multiplex mixture using *Padi2* WT primers: Forward 5' CTTATTCCTGGGAGCAGCAC 3', Reverse 5' CTCTCTCTCGGCTCACCTGT 3' and KO primers: LacZ-Neo Forward 5' GATCGGCCATTGAACAAGAT 3', Reverse 5' GTAGCCGGATCAAGCGTATG 3'. *Padi4*-null mice (FVB/NJ background) were generated using Regeneron's VelociGene technology. The genomic sequence of *Padi4* from the ATG initial codon to approximately 100bp downstream of the TGA stop codon was replaced in frame (with respect to the *Padi4* initiation codon) with a Lox-flanked LacZ and neomycin coding

sequence. These mice are genotyped using *Padi4* primers: Forward 5' CTCGCCTGGCACTATAAAGG 3' and Reverse 5' AGGATCTCTGCCCCCTCTCAC 3'.

Generation of *Padi2/Padi4* double knockout mice

Padi2-null mice were used for IVF and CRISPR/Cas9 RNA injection was performed at Cornell's Stem Cell & Transgenic Core Facility. Cas9 mRNA and *Padi4* sgRNA targeting Exon 1 of the *Padi4* gene was injected into the pronucleus of *Padi2* KO embryos and the embryos were cultured to the two-cell stage and transferred into pseudopregnant recipient FVB/NJ female mice (17,18). Thirty-nine founders were produced and 13 edited founders were then identified by heteroduplex PCR (19). Briefly, genomic DNA was extracted from punched ear tissue and evaluated by PCR




Figure 1. Design of *Padi2/Padi4* double knockout mice. (a) Schematic of the CRISPR protocol using *Padi2* KO zygotes. (b) Design and sequence of the guide RNA targeting Exon 1 of the *Padi4* gene. (c) Sanger sequencing confirmed a four-nucleotide deletion in the DKO *Padi4* Exon 1, resulting in (d) a frame-shift mutation and premature stop codon in the predicted amino acid sequence.

using the *Padi4* primer set noted above. Standard PCR conditions for GoTaq (Promega) were used as follows: 95°C for 5 min; 94°C for 30 s, 59°C for 30 s, 72°C for 30s for 35 cycles; 72°C for 5 min followed by 5 minutes of denaturation at 95°C. PCR products were loaded onto a 12% polyacrylamide gel. Positive heteroduplex bands were purified and subjected to TA cloning for Sanger sequencing. All 13 edited founders were found to be heterozygous mutants, with 7 founders being identified as having a frameshift mutation either by insertion or deletion. One heterozygous founder with a four-nucleotide deletion was bred back to *Padi2* knockout mice, and their heterozygous offspring were crossed to create a homozygous DKO line (Figure 1).

Authentication of Padi2/Padi4 double knockout mice

RT-PCR of Padi2 transcripts. Total RNA was extracted from mouse salivary glands using Tri-Reagent (Molecular Research Center #TR118) and cDNA was created with a High Capacity RNA-to-cDNA Kit (Applied Biosystems #4387406) according to manufacturers' instructions. PCR amplification of *Padi2* was accomplished under standard cycling conditions with GoTaq (Promega #M3001) using the following *Padi2* primers: Forward 5' TCCCCTCGAGATGGAAACCT 3' and Reverse 5' CTCCCAGGCCCTTGAACATAA 3'. PCR products were run on a 2% Agarose gel to confirm the presence or absence of *Padi2* transcripts (Figure 2a).

PCR of Padi4 locus. Genomic DNA was isolated from ear-punch tissue samples and the *Padi4* locus was amplified under standard GoTaq conditions using the following *Padi4* primers: Forward 5' CTCGCCTGGCACTATAAAGG 3' and Reverse 5'

AGGATCTCTGCCCCTCTCAC 3'. PCR products were run on a 2% Agarose gel to confirm the presence or absence of the *Padi4* gene (Figure 2b).

Western blot. Tissue lysate was prepared from splenic tissue and resolved by SDS-PAGE followed by transfer to a PVDF membrane. The membranes were incubated overnight with primary antibodies diluted in 1% BSA-TBST at 4°C using the following concentrations: anti-PAD4 (1:1000; Abcam; ab214810), anti-PAD2 (1:1000, Proteintech, 12110-1-AP) and anti-actin (1:1000, Abcam, ab-8227). The primary antibodies were detected with HRP-conjugated secondary antibodies and were exposed to ECL reagents to determine the presence or absence of PAD2 and PAD4 proteins (Figure 2c).

Breeding Trials

Seven-week old virgin mice were paired in the following groups: DKOxDKO, DKOxWT, WTxDKO or WTxWT (n=5-6). Pairs were monitored daily for 63 days and birth dates, litter sizes, sex ratios, 2-day old pup weights, and weanling weights were recorded. Statistical analyses were carried out using Microsoft Excel and Social Science Statistics calculators for unpaired Student's t-test, ANOVA, χ^2 and Goodness of Fit tests with significance set at $p < 0.05$ (20).




Figure 2. Authentication of the *Padi2/Padi4* double knockout. (a) RT-PCR of salivary gland tissue shows that the DKO mice lack *Padi2* transcripts. (b) PCR analysis of genomic DNA (15% polyacrylamide gel) confirms a 4bp deletion within the *Padi4* gene of DKO mice. (c) Western blot analysis of spleen lysate confirms that DKO mice lack both PAD2 and PAD4 mature proteins.

Growth and development

Male weanlings were housed singly (to avoid intra-cage dominance issues), weighed and monitored daily for preputial separation as previously described (21). Mice were euthanized for tissue harvest at ages 46, 48, 50 and 90 days old. At each of these time points, five mice were euthanized by CO₂ inhalation and blood was collected by cardiac puncture, allowed to clot at room temperature for 90 minutes, spun at 2000g for 15 minutes, and the serum was snap frozen and saved for hormone analysis. Serum testosterone, Follicle Stimulating Hormone (FSH), and Lutenizing Hormone (LH) levels were quantified in duplicate by ELISA at the University of Virginia Ligand Assay Core. Testes were collected and weighed individually. Results were analyzed with unpaired t-tests with significance set at $p < 0.05$.

Histological evaluation

Testis, epididymis, vas deferens, seminal vesicle and splenic tissues were collected at 90 days from 5 animals from each of the following strains: Wild-type (WT, FVB/NJ), P2KO, P4KO, and DKO. Collected tissues were fixed in 10% formalin solution, embedded in paraffin, sectioned at 5 μ m, and stained with hematoxylin and eosin (H&E). Initial evaluation revealed epididymis, vas deferens, seminal vesicles, and spleen within normal limits based on previously published observations (22,23), therefore, only testis sections were evaluated further. The noted changes were scored on a scale of 0 to 3 according to the following criteria: 0 = absent to affecting less than 5% of the tissue; 1 = mild change, affecting 5-25% of the tissue; 2 = moderate change, affecting 25-50% of the tissue; 3 = severe change, affecting >50% of the tissue (Supplemental 2).

RNA sequencing

Mouse testes were dissected from two age-matched adult mice per strain (WT, P2KO, and DKO) and snap-frozen immediately following euthanasia. Total RNA extraction was performed using TRI reagent (Molecular Research Center, Inc. OH) according to the manufacturer's protocol. The RNA concentration and purity were measured by Nanodrop and Qubit (Thermo Scientific, USA).

RNA-seq experiments were performed in the Transcriptional Regulation & Expression Facility (TReX) at Cornell University. RNA integrity was confirmed on a Fragment Analyzer (Agilent) and RNAseq libraries were prepared using the NEB Ultra II Directional RNA library Prep kit (New England Biolabs), with 1 ug total RNA input and initial poly(A) selection. Illumina sequencing was performed on a NovaSeq to generate at least 20M 2x150 paired end reads per sample. The raw fastq reads were first processed with the Trim Galore package to trim for low quality reads, noisy short fragments, and adapter sequence (24). The filtered reads were then aligned to the GRCm38 reference genome (mm10) with ENSEMBL annotations via STAR using parameters [--outSAMstrandField intronMotif , --outFilterIntronMotifs RemoveNoncanonical, --outSAMtype BAM SortedByCoordinate, --quantMode GeneCounts] (25). Differential gene expression analysis was performed by DESeq2 (26,27). A heatmap and hierarchical clustering dendrogram of differentially expressed genes was produced using Morpheus (28). Venn diagrams were produced with BioVenn (29).

Gene ontology classification analysis was performed using DAVID v6.8, a web-based bioinformatics resource for functional genomics analysis to identify biological processes,

molecular functions, and cellular components (30,31). 246 uniquely upregulated genes were identified when comparing DKO to WT using an FDR < 0.05. 238 of these genes were identified in the DAVID database and used for subsequent analysis.

RESULTS and DISCUSSION

Generation of Padi2/Padi4 double knockout mice

Our strategy for generating the *Padi2/Padi4* DKO mouse line is shown in Figure 1a. *Padi2*-null males and females were bred to generate *Padi2*-null zygotes and these embryos were then injected with a *Padi4* sgRNA construct and transferred to wild type (FVB/NJ) recipient females to generate founder pups. The *Padi4* sgRNA construct was designed to target Exon 1 of the *Padi4* gene (Figure 1b). In total, 39 founders were born, with 13 CRISPR-edited founders being identified by heteroduplex PCR, subcloning, and Sanger sequencing. All of the edited founders were heterozygous mutants, with seven of these founders having a frame-shift mutation. Following back breeding to P2KO, crossing the heterozygous F1 offspring, and genotyping successive generations, one homozygous DKO strain was established and selected for further characterization. This mutant mouse line contained a four-nucleotide deletion in Exon 1 of *Padi4* (Figure 1c) and a predicted premature stop codon (Figure 1d). Validation of the DKO mutant line is shown in Figure 2. We first confirmed that P2KO and DKO mice lacked *Padi2* transcripts. RT-PCR analysis of salivary gland RNA showed that the P2KO and DKO mice lacked a *Padi2* PCR product, while the appropriately sized product was seen in the WT sample (Figure 2a). We next tested whether the *Padi4* sgRNA targeted site in DKO mice contained a deleted sequence by

performing PCR analysis of genomic DNA and then separating the DNA fragments on high percentage polyacrylamide gels. Results showed that the DKO amplicon displayed a reduced molecular weight when compared to the P2KO and WT amplicons (Figure 2b), suggesting that DKO mice contained the targeted deletion. As expected, no amplicon was observed in P4KO mice. The above results suggested that DKO mice lacked both *Padi2* and *Padi4* transcripts. We then performed Western blot analysis of WT, P2KO, P4KO and DKO spleen lysates to test if the DKO mice also lacked PAD2 and PAD4 protein. We probed the protein lysates with anti-PADI2 antibodies and found that, while the antibody was reactive with appropriately sized ~75 kDa bands in the WT and P4KO lysates, the antibody did not react with similarly sized bands in the P2KO and DKO mice (Figure 2c, Supplemental 1). Likewise, the anti-PADI4 antibody was reactive with ~75 kDa bands in the WT and P2KO lysates but was not reactive with similarly sized bands in the P4KO and DKO lysates (Figure 2c, Supplemental 1). These results indicate that the *Padi2/Padi4* DKO strain does not express mature PAD2 and PAD4 proteins.

Effect of Padi2/Padi4 DKO on fertility parameters

In order to begin testing for links between PADs and reproductive function, we first performed a controlled breeding trial to test whether a range of reproductive parameters were affected by DKO. Seven-week old mice were paired in 4 mating categories (WTxWT, DKOxDKO, DKOxWT and WTxDKO) and monitored daily for 63 days. Results showed that, although not statistically significant, DKOxDKO pairs tend to lag behind WTxWT pairs in the time taken to have their first litter. To identify which sex may

be responsible for this reproductive delay, we bred DKO males with WT females and DKO females with WT males, and the trend towards decreased fertility is still readily apparent for DKO males, but not with females (Figure 3a). Another larger fertility trial is currently being planned to test whether increasing the sample size will yield statistically significant differences in the fertility rate.

With respect to the offspring produced, we found that the sex ratio of homozygous DKO offspring was skewed (0.85 male:female) but did not deviate significantly from the expected equal ratio (χ^2 test, $n=243$, $p=0.22$), and that the average DKO litter size was similar to that of WT mice ($n=4$, $p=0.5$). We also observed that 2-day old DKO pups




Figure 3. Breeding phenotype of *Padi2/Padi4* double knockout mice. (a) In controlled breeding trials comparing DKO males to DKO females when paired with WT mates, DKO males ($n=12$) took longer than DKO females ($n=10$) to have their first litter and had fewer pup-producing pairs (data trending, but not significant, one-tailed Fisher's Exact Test, $p=0.19$). (b) Two days postpartum, DKO pups were significantly smaller than WT pups (one-tailed t-test, $*p=0.004$). (c) Male DKO are significantly smaller than both SKO and WT mice at weaning (ANOVA, $p=0.02$, post-hoc one-tailed t-test with WT, $**p=0.003$). (d) Male DKO weanlings are visibly smaller and less developed than WT and SKO weanlings.

weighed significantly less than WT pups (Figure 3b, $n=4$ litters, $p=0.004$) and that DKO male weanlings weighed significantly less than WT male weanlings (Figure 3c-d, $n=21,45$, $p=0.003$). Taken together, results from these studies suggest that loss of PAD2 and PAD4 suppresses fertility and affects offspring weight. Previous studies have shown that pup weight is also commonly reduced in sub-fertile mouse strains, including androgen receptor knockout (ARKO) mice (32) which is relevant to the research reported here. Given the growing links between PADs and hormone signaling, we predicted that the effect of DKO on both fertility and pup weight was due to disrupted hormone signaling in these mice.

Effect of Padi2/Padi4 DKO on pubertal onset

Pubertal onset is thought to be primarily driven by increased testosterone production in males and one well-established external marker of pubertal development in rodents is preputial separation (PS) (21,33,34).

Therefore, as a further test of the hypothesis that hormone signaling is disrupted in DKO males, we next documented the timing of PS in WT and DKO males. Results show that DKO males ($n=46$, $m=31$) took an average of three days longer than WT males ($n=49$, $m=28$) to undergo PS ($p<0.00001$, Figure 4).

Given that PS is known to be an androgen-




Figure 4. Pubertal onset is delayed in *Padi2/Padi4* DKO male mice. DKO males took an average of 31 days to undergo preputial separation (PS) compared to 28 days for WT males ($n=46$, two-tailed t-test, $*p<0.00001$).

dependent process (34,35) and to be strongly linked to hormone signaling along the hypothalamic-pituitary-gonadal (HPG) axis (36,37), these results further support the hypothesis that hormone signaling is altered in DKO males. Additionally, these observations suggest that PAD2 and PAD4 play an important role in androgen-driven sexual development in males.

Effect of Padi2/Padi4 DKO on serum hormone levels

As a more direct test of the hypothesis that hormone signaling is defective in DKO mice, we next measured serum testosterone (T), luteinizing hormone (LH), and follicle stimulating hormone (FSH) in WT and DKO males. Results showed that levels of T, LH, and FSH differed significantly between DKO and WT at various times (n=5 per time point). Although highly variable, serum testosterone levels were lower in DKO males than WT males at all time-points, most significantly at day 48 (p=0.05, Figure 5). In contrast, FSH and LH levels were statistically similar between DKO and WT at the earlier time points, but by




Figure 5. *Padi2/Padi4* DKO male mice show altered serum hormone levels. (a) Serum Testosterone levels were lower in DKO males when compared to WT males at all timepoints, reaching significance at day 48. (b,c) LH and FSH levels were significantly higher in DKO on day 50, then dropped to values that were significantly lower than WT by day 90 (n=5, one-tailed t-test, *p=0.05, **p=0.02).

day 50 both hormones were significantly higher in DKO sera ($p < 0.03$). Serum testosterone levels are difficult to accurately measure in mice, likely due to the fact that mice do not express significant levels of sex-hormone binding globulin (38). Therefore, we were encouraged by the observation that serum T levels in DKO mice were lower than WT mice at each time point evaluated, reaching significance at day 48. Given that T is known to negatively regulate FSH and LH secretion (39), we were also encouraged by the observation that, on the day when T levels appeared to be lowest in DKO males (day 50), FSH and LH levels were correspondingly elevated. Importantly, studies with ARKO mice also found that serum T levels are lower in mutant mice compared to WT mice (32). Given this observation, our findings lend support to the prediction that PAD enzymes play a role in androgen signaling *in vivo*.

Effect of Padi2/Padi4 DKO on testis size and histology

Testis Size. Hormone signaling within the HPG axis not only regulates pubertal onset, but also regulates body growth (40) and testis size (41,42). In support of this observation, previous studies with ARKO mice found that both pup size and testis weight are reduced in mutant mice (32). As a further test for potential associations between PADs and hormone signaling, we next investigated whether the total body weight and testis weight was altered in 90-day old adult mice. Results showed that, while there was no difference in total body weight between 90-day old WT ($m = 27.4g$) and DKO ($m = 27.4g$) mice ($n = 5$, $p = 1$), testis weight was significantly lower in DKO males when compared to WT males ($n = 5$, $p = 0.002$, Figure 6a). The finding that DKO testes are significantly smaller than WT testes in adults fits well with the hypothesis that PADs

play a role in androgen signaling in males. Given our hypothesis, it is currently unclear why the body weight of adult DKO mice is similar to that of adult WT mice. It is possible that the effect of PAD2/4 DKO on adult male body weight is transient in nature and that other PADs may be able to compensate for PAD2/4 in adults.




Figure 6. *Padi2/Padi4* DKO male mice have disrupted testicular development. (a) Total testes weight was significantly lower in 90-day old DKO mice compared to WT (n=5, one-tailed t-test, *p=0.0005). (b) Hematoxylin and Eosin stained sections of WT testis showing normal spermatogenic cell structures. (c-e) Hematoxylin and Eosin stained sections of DKO testis. Black arrows indicate: (c) apoptotic spermatocytes, (d) apoptotic spermatogonia and (e) atypical residual bodies. (scale bars = 50 μ m).

Testis Histology. The observation that DKO testes are smaller than WT testes raised the possibility that testicular histology may also be defective in DKO males. To test this hypothesis, FFPE sections of adult testes were stained with H&E and scored for cellular morphology by a board-certified pathologist (Supplemental 2). Results of this analysis found that there were variable degrees of germ cell apoptosis in the DKO testes when compared to WT testes (Figure 6b), with spermatogonia and spermatocytes being the most significantly affected. More specifically, apoptosis (mild/moderate grade) in spermatocytes (Figure 6c) was noted in 5 out of 5 DKO mice compared to only 2 out of 5 WT samples. Apoptosis (mild grade) of spermatogonia (Figure 6d) was noted in 4 out

of 5 DKO mice testes compared to only 1 out of 4 WT mice. All five DKO testes also displayed moderate grade residual bodies (Figure 6e) when compared with WT testes, which only displayed mild grade residual bodies. Sertoli, Leydig cells and interstitium were overall unremarkable. The increased rate of apoptosis and residual bodies did not appear to result in a decrease in sperm production with the lumen of most tubules being lined with histomorphologically unremarkable elongated spermatids and spermatozoa. Interestingly, in ARKO mice, spermatogenesis was arrested at the pachytene stage of meiosis and there was an increase in apoptotic-like bodies within the tubules (32) suggesting that androgen signaling is critical for normal spermatogenesis. Taken together, these results support the hypothesis that PAD2/4 signaling may play a role in androgen signaling-mediated spermatogenesis. The relatively modest effect of DKO on testicular histology and the other observed hormone-mediated phenotypic traits could potentially be due to compensatory effects of other PADs in the DKO males and subsequent studies are planned to test this hypothesis. As an alternate hypothesis, it may be that PAD2/4 is only involved in regulating a portion of androgen-mediated signaling events.

Effect of Padi2/Padi4 DKO on gene expression in the testis

In order to begin investigating whether PADs are associated with specific signaling pathways in the testis, and to more directly test the hypothesis that PAD2/4 play a role in regulating AR target gene expression, we next carried out RNA-Seq analysis of adult WT, P2KO, and DKO testes. Results show that 22 genes were upregulated and 19 genes were downregulated in P2KO samples when compared to WT samples ($p < 0.05$).

Additionally, we found that 263 genes were upregulated and 140 genes were down-regulated in the DKO samples compared to WT samples ($p < 0.05$). Seventeen of the 22 genes that were significantly upregulated in the P2KO samples were also upregulated in the DKO samples. A comparison of upregulated genes is shown in Figure 7a. The complete lists of differentially expressed genes (DEGs) are shown in Supplementals 3 and 4. Two hundred and forty-six genes that were uniquely upregulated in DKO testes were then subjected to DAVID gene ontology analysis. Of the 246 genes, 238 were identified in the mouse database and these genes were categorized into three domains: Cellular Components, Molecular Functions, and Biological Processes (Supplemental 5). Analysis of the Biological Processes domain finds that a majority of the DE genes are involved in reproductive functions (Figure 7b), thus supporting the phenotypic traits we observed. In addition, analysis of the




Figure 7. RNA-seq analysis of gene expression in WT and DKO testes. (a) Venn diagram of differential gene expression shows that 263 genes are significantly upregulated in DKO in comparison to WT, with only 17 genes overlapping the P2KO profile. Gene lists are provided in Supplemental 3 and 4. (b-c) Gene ontology analysis with DAVID 6.8 divides these upregulated genes into (b) Biological Processes with reproductive pathways heavily represented and (c) Molecular Functions almost entirely represented by genes involved in nucleic acid regulation.

Molecular Function domain finds that nearly all of the DEGs appear to be involved in either nucleic acid binding or protein phosphorylation (Figure 7c), thus further




Figure 8. Heatmap and unsupervised hierarchical dendrogram illustrating the top 100 differentially expressed genes between WT, P2KO and DKO. Red=upregulated, Blue=down-regulated. The list of DEGs used to create the heatmap is provided in Supplement 6 (Table S5).

supporting the hypothesis that PADs play a role in regulating gene transcription. We next generated a hierarchical dendrogram and heatmap showing the top 100 significantly DEGs using summary counts over all pairwise tests of WT, P2KO, and DKO samples (FDR cut-off 0.05). Results show that there is a good correlation in gene expression levels between replicates for each experimental group. Additionally, the gene expression profile for the DKO samples is distinctly different from that of the WT samples, while there are some overlapping expression patterns when comparing the DKO and P2KO samples (Figure 8). Analysis of the list of genes that are most highly overexpressed in the DKO samples compared to WT and P2KO samples, finds that 3 of the top 10 genes (Klk1b27, Klk1b21, and Klk1b24) are members of the kallikrein gene family

(Supplemental 6). Perhaps the best-known member of this family is prostate-specific antigen (KLK3), whose expression is mainly induced by androgen and is transcriptionally regulated by the androgen receptor (43). Additionally, the kallikrein gene locus is highly responsive to steroid hormones, having at least 14 functional HREs in this region. In fact, many researchers use kallikreins as markers of hormone receptor activity (44). Importantly, Klk1b27, Klk1b21, and Klk1b24 are all expressed in Leydig cells in rodents and their expression is regulated by androgen (44). In addition to kallikreins, a number of other genes that were overexpressed in the DKO gene set also appear to be regulated by androgens. These findings further support our hypothesis that PADs play an important role in regulating gene expression in the testis, and more specifically, in regulating AR-mediated gene expression.

CONCLUSIONS

Our previous work with PAD2 and PAD4 in breast cancer cell lines has found that both of these PAD isozymes appear to regulate ER target gene expression via citrullination of histone residues at ER binding sites. More recent studies have shown that PAD2 also appears to facilitate androgen receptor target gene expression in prostate cancer cells via similar mechanisms. Together, these observations suggest that PAD2 and PAD4 may represent key mediators of hormone signaling in mammals. While these *in vitro* findings are intriguing, an important next step in linking PADs with hormone signaling is demonstrating that the phenomenon also occurs at the organismal level. While our *Padi2* and *Padi4* SKO mouse models have failed to show a reproductive phenotype, outcomes from our *Padi2/Padi4* DKO studies provide a series of clear associations

between PADs and hormone signaling and, to our knowledge, this study is the first to make such genetic links using animal models. Subsequent studies are now being designed to further investigate the direct mechanisms by which PADs are regulating hormone signaling in males. An important part of these future studies will be to localize the expression of both PAD2 and PAD4 in the testis. Recent studies have documented the presence of PAD2 in the testis. For example, Hosokawa et al. showed that PAD2 appears to play an important role in testis development and is specifically expressed in fetal Sertoli cells with its expression being regulated by SOX9, a critical mediator of sex determination (45). Another recent study found by PCR, western blotting, and IHC that PAD2 is expressed in sperm during the first wave of spermatogenesis and appears to localize to the acrosomal region of the sperm head (46). There are currently no reports of PAD4 localization to the testes, however, IHC analysis of PAD4 staining in the testis from the Human Protein Atlas website (47) does show anti-PAD4 staining (moderate intensity) specifically in Leydig cells. Based on these reports, we predict that the phenotypic effects of DKO that we have observed in this study are due to the lack of PAD2 and PAD4 activity in the testis.

Our DKO mice described here exhibit several phenotypic traits characteristic of other sub-fertile strains, including delayed puberty, small offspring and decreased serum testosterone. Since PAD2 and PAD4 have previously been implicated in interactions with estrogen and androgen receptors, it is not unexpected to see reproductive deficits in these mice. However, our further finding of dysregulated serum Testosterone, LH and FSH levels indicates a wider reaching disruption of the HPG axis with implications for more than just reproductive fitness.

LIST OF ABBREVIATIONS

Padi or PAD = Peptidylarginine deiminase

DKO = Double knockout

SKO = Single knockout

WT = Wildtype (FVB/NJ)

E2 = Estrogen

ER = Estrogen receptor

ERE = Estrogen response element

AR = Androgen receptor

ARKO = Androgen receptor knockout

PS = Preputial separation

HPG = Hypothalamic-pituitary-gonadal axis

T = Testosterone

FSH = Follicle stimulating hormone

LH = Lutenizing hormone

H&E = Hematoxylin and Eosin stain

DEGs = Differentially expressed genes

DECLARATIONS

Ethics approval and consent to participate: All animal procedures were approved by Cornell University's Animal Care and Use Committee (IACUC protocol #2007-0115) and are in compliance with the US National Research Council's Guide for the Care and Use

of Laboratory Animals, the US Public Health Service's Policy on Humane Care and Use of Laboratory Animals, and Guide for the Care and Use of Laboratory Animals.

Consent for publication: Not applicable.

Availability of data and materials: The datasets supporting the conclusions of this article will be available upon publication in NCBI's Gene Expression Omnibus (GEO) repository (GSE194034, <https://www.ncbi.nlm.nih.gov/geo/>). *Padi2/Padi4* double knockout mice are available from the corresponding author upon reasonable request, and in accordance with Cornell University's Material Transfer Agreement.

Competing interests: All authors declare that they have no competing interests.

Funding: This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Authors' contributions: Experimentation was conceived and conducted by KLS, CM, BAM and SAC. Breeding and phenotype analysis by KLS and SAC. Histological scoring by EAD and SN. RNA sequencing and analysis by JKG, AET, and FA, with visualization by CM and KLS. All authors read and approved the final manuscript.

Acknowledgements: The authors acknowledge the excellent skills and technology provided by Cornell's Stem Cell & Transgenic Core Facility to create our double

knockout mice. We also thank the Baker Institute husbandry technicians for the outstanding care of the animals used in this research.

REFERENCES

1. Wang S, Wang Y. Peptidylarginine deiminases in citrullination, gene regulation, health and pathogenesis. *Biochim Biophys Acta*. 2013 Oct;1829(10):1126–35.
2. Witalison EE, Thompson PR, Hofseth LJ. Protein Arginine Deiminases and Associated Citrullination: Physiological Functions and Diseases Associated with Dysregulation. *Curr Drug Targets*. 2015;16(7):700–10.
3. Vossenaar ER, Zendman AJW, van Venrooij WJ, Pruijn GJM. PAD, a growing family of citrullinating enzymes: genes, features and involvement in disease. *BioEssays*. 2003;25(11):1106–18.
4. Valesini G, Gerardi MC, Iannuccelli C, Pacucci VA, Pendolino M, Shoenfeld Y. Citrullination and autoimmunity. *Autoimmun Rev*. 2015 Jun 1;14(6):490–7.
5. Takahara H, Kusubata M, Tsuchida M, Kohsaka T, Tagami S, Sugawara K. Expression of peptidylarginine deiminase in the uterine epithelial cells of mouse is dependent on estrogen. *J Biol Chem*. 1992 Jan 5;267(1):520–5.
6. Dong S, Zhang Z, Takahara H. Estrogen-Enhanced Peptidylarginine Deiminase Type IV Gene (PADI4) Expression in MCF-7 Cells Is Mediated by Estrogen Receptor- α -Promoted Transfactors Activator Protein-1, Nuclear Factor-Y, and Sp1. *Mol Endocrinol*. 2007 Jul 1;21(7):1617–29.
7. Mani SK, Mermelstein PG, Tetel MJ, Anesetti G. Convergence of Multiple Mechanisms of Steroid Hormone Action. *Horm Metab Res*. 2012 Jul;44(8):569–76.

8. Hagiwara T, Nakashima K, Hirano H, Senshu T, Yamada M. Deimination of Arginine Residues in Nucleophosmin/B23 and Histones in HL-60 Granulocytes. *Biochem Biophys Res Commun*. 2002 Jan 25;290(3):979–83.
9. Wang Y, Wysocka J, Sayegh J, Lee YH, Perlin JR, Leonelli L, et al. Human PAD4 Regulates Histone Arginine Methylation Levels via Demethylination. *Science*. 2004 Oct 8;306(5694):279–83.
10. Wang L, Song G, Zhang X, Feng T, Pan J, Chen W, et al. PADI2-Mediated Citrullination Promotes Prostate Cancer Progression. *Cancer Res*. 2017 Nov 1;77(21):5755–68.
11. Zhang X, Bolt M, Guertin MJ, Chen W, Zhang S, Cherrington BD, et al. Peptidylarginine deiminase 2-catalyzed histone H3 arginine 26 citrullination facilitates estrogen receptor α target gene activation. *Proc Natl Acad Sci*. 2012 Aug 14;109(33):13331–6.
12. Guertin MJ, Zhang X, Anguish L, Kim S, Varticovski L, Lis JT, et al. Targeted H3R26 Deimination Specifically Facilitates Estrogen Receptor Binding by Modifying Nucleosome Structure. *PLOS Genet*. 2014 Sep 11;10(9):e1004613.
13. Briot J, Simon M, Méchin MC. Deimination, Intermediate Filaments and Associated Proteins. *Int J Mol Sci*. 2020 Nov 19;21(22):8746.
14. Guide-for-the-Care-and-use-of-laboratory-animals.pdf [Internet]. [cited 2022 Apr 18]. Available from: <https://grants.nih.gov/grants/olaw/Guide-for-the-Care-and-use-of-laboratory-animals.pdf>

15. Mouse Colony Management Software: Access the SoftMouse.NET app on your mobile Android and iPad tablets [Internet]. [cited 2022 Apr 18]. Available from: <https://www.softmouse.net/>
16. Hansen GM, Markesich DC, Burnett MB, Zhu Q, Dionne KM, Richter LJ, et al. Large-scale gene trapping in C57BL/6N mouse embryonic stem cells. *Genome Res.* 2008 Oct 1;18(10):1670–9.
17. Singh P, Schimenti JC, Bolcun-Filas E. A Mouse Geneticist's Practical Guide to CRISPR Applications. *Genetics.* 2015 Jan 1;199(1):1–15.
18. Singh P, Schimenti JC. The genetics of human infertility by functional interrogation of SNPs in mice. *Proc Natl Acad Sci.* 2015 Aug 18;112(33):10431–6.
19. Zhu X, Xu Y, Yu S, Lu L, Ding M, Cheng J, et al. An Efficient Genotyping Method for Genome-modified Animals and Human Cells Generated with CRISPR/Cas9 System. *Sci Rep.* 2014 Sep 19;4(1):6420.
20. Social Science Statistics [Internet]. [cited 2022 Apr 18]. Available from: <https://www.socscistatistics.com/>
21. Hoffmann HM. Determination of Reproductive Competence by Confirming Pubertal Onset and Performing a Fertility Assay in Mice and Rats. *JoVE J Vis Exp.* 2018 Oct 13;(140):e58352.
22. Creasy D, Bube A, Rijk E de, Kandori H, Kuwahara M, Masson R, et al. Proliferative and Nonproliferative Lesions of the Rat and Mouse Male Reproductive System. *Toxicol Pathol.* 2012 Aug 1;40(6_suppl):40S-121S.
23. Meistrich ML, Hess RA. Assessment of Spermatogenesis Through Staging of Seminiferous Tubules. In: Carrell DT, Aston KI, editors. *Spermatogenesis* [Internet].

Totowa, NJ: Humana Press; 2013 [cited 2021 Nov 18]. p. 299–307. (Methods in Molecular Biology; vol. 927). Available from: http://link.springer.com/10.1007/978-1-62703-038-0_27

24. Babraham Bioinformatics - Trim Galore! [Internet]. [cited 2022 Apr 18]. Available from: https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/

25. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 2013 Jan;29(1):15–21.

26. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*. 2014 Dec 5;15(12):550.

27. Varet H, Brillet-Guéguen L, Coppée JY, Dillies MA. SARTools: A DESeq2- and EdgeR-Based R Pipeline for Comprehensive Differential Analysis of RNA-Seq Data. *PLoS ONE*. 2016 Jun 9;11(6):e0157022.

28. Morpheus [Internet]. [cited 2022 Apr 18]. Available from: <https://software.broadinstitute.org/morpheus/>

29. Hulsen T, de Vlieg J, Alkema W. BioVenn – a web application for the comparison and visualization of biological lists using area-proportional Venn diagrams. *BMC Genomics*. 2008 Oct 16;9(1):488.

30. Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res*. 2009 Jan 1;37(1):1–13.

31. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc*. 2009 Jan;4(1):44–57.

32. Yeh S, Tsai MY, Xu Q, Mu XM, Lardy H, Huang KE, et al. Generation and characterization of androgen receptor knockout (ARKO) mice: an in vivo model for the study of androgen functions in selective tissues. *Proc Natl Acad Sci U S A*. 2002 Oct 15;99(21):13498–503.
33. Gaytan F, Bellido C, Aguilar R, Aguilar E. Balano-Preputial Separation as an External Sign of Puberty in the Rat: Correlation with Histologic Testicular Data/Balano-
praeputial-Trennung als ein äußeres Zeichen der Pubertät für die Ratte: Korrelation mit histologischen Daten. *Andrologia*. 1988;20(5):450–3.
34. Korenbrot CC, Huhtaniemi IT, Weiner RI. Preputial Separation as an External Sign of Pubertal Development in the Male Rat¹. *Biol Reprod*. 1977 Sep 1;17(2):298–303.
35. Yoshimura S, Yamaguchi H, Konno K, Ohsawa N, Noguchi S, Chisaka A. Observation of Preputial Separation is a Useful Tool for Evaluating Endocrine Active Chemicals. *J Toxicol Pathol*. 2005;18(3):141–57.
36. Handelsman DJ. Testosterone, Spermatogenesis, and Unravelling the Mysteries of Puberty. *Endocrinology*. 2020 Sep 1;161(9):bqaa120.
37. Keene DE, Suescun MO, Bostwick MG, Chandrashekar V, Bartke A, Kopchick JJ. Puberty Is Delayed in Male Growth Hormone Receptor Gene—Disrupted Mice. *J Androl*. 2002;23(5):661–8.
38. Jänne M, Deol HK, Power SG, Yee SP, Hammond GL. Human sex hormone-binding globulin gene expression in transgenic mice. *Mol Endocrinol Baltim Md*. 1998 Jan;12(1):123–36.

39. Hayes FJ, DeCruz S, Seminara SB, Boepple PA, Crowley WF. Differential regulation of gonadotropin secretion by testosterone in the human male: absence of a negative feedback effect of testosterone on follicle-stimulating hormone secretion. *J Clin Endocrinol Metab.* 2001 Jan;86(1):53–8.
40. Corre C, Shinoda G, Zhu H, Cousminer DL, Crossman C, Bellissimo C, et al. Sex-specific regulation of weight and puberty by the Lin28/let-7 axis. *J Endocrinol.* 2016 Mar;228(3):179–91.
41. Huh K, Nah WH, Xu Y, Park MJ, Gye MC. Effects of Recombinant Human Growth Hormone on the Onset of Puberty, Leydig Cell Differentiation, Spermatogenesis and Hypothalamic KISS1 Expression in Immature Male Rats. *World J Mens Health.* 2021 Apr;39(2):381–8.
42. Miyaso H, Nagahori K, Takano K, Omotehara T, Kawata S, Li ZL, et al. Neonatal maternal separation causes decreased numbers of sertoli cell, spermatogenic cells, and sperm in mice. *Toxicol Mech Methods.* 2021 Feb 12;31(2):116–25.
43. Kim J, Coetzee GA. Prostate specific antigen gene regulation by androgen receptor. *J Cell Biochem.* 2004;93(2):233–41.
44. Lawrence MG, Lai J, Clements JA. Kallikreins on Steroids: Structure, Function, and Hormonal Regulation of Prostate-Specific Antigen and the Extended Kallikrein Locus. *Endocr Rev.* 2010 Aug 1;31(4):407–46.
45. Tsuji-Hosokawa A, Kashimada K, Kato T, Ogawa Y, Nomura R, Takasawa K, et al. Peptidyl arginine deiminase 2 (Padi2) is expressed in Sertoli cells in a specific manner and regulated by SOX9 during testicular development. *Sci Rep.* 2018 Sep 5;8:13263.

46. Sahadevan M, Kumar PG. Peptidyl Arginine Deiminase 2 (PADI2) is Expressed in Post-Meiotic Germ Cells in the Mouse Testis and is Localized Heavily on the Acrosomal Region of Spermatozoa. J Endocrinol Reprod. 2021 Jul 20;25(1):37–49.
47. Pontén F, Jirström K, Uhlen M. The Human Protein Atlas--a tool for pathology. J Pathol. 2008 Dec;216(4):387–93.