Palatine Tonsils vs Attached/Keratinized gingiva

PERMANOVA (Bray−Curtis, permutations within subject): R2 = 0.379, F = 8.56, p = 0.008, q = 0.011
PERMDISP p = 0.270 | n = 8 + 8 samples



Palatine Tonsils vs Buccal mucosa

PERMANOVA (Bray−Curtis, permutations within subject): R2 = 0.278, F = 15.41, p = < 0.001, q = < 0.001
PERMDISP p = 0.013 | n = 21 + 21 samples




Palatine Tonsils vs Saliva
PERMANOVA (Bray−Curtis, permutations within subject): R2 = 0.148, F = 1.39, p = 0.062, q = 0.073
PERMDISP p = 0.719 | n = 5 + 5 samples
PCoA2 (16.6%)
0.4
0.0
−0.4
−0.8
−1.0
−0.5
0.0
0.5
PCoA1 (33.4%)
Palatine Tonsils
Saliva

Palatine Tonsils vs Subgingival plaque

PERMANOVA (Bray−Curtis, permutations within subject): R2 = 0.343, F = 10.43, p = < 0.001, q = 0.002
PERMDISP p = 0.510 | n = 11 + 11 samples



Palatine Tonsils vs Supragingival plaque

PERMANOVA (Bray−Curtis, permutations within subject): R2 = 0.319, F = 18.74, p = < 0.001, q = < 0.001
PERMDISP p = 0.104 | n = 21 + 21 samples



Palatine Tonsils vs Throat

PERMANOVA (Bray−Curtis, permutations within subject): R2 = 0.040, F = 0.59, p = 0.500, q = 0.500
PERMDISP p = 0.801 | n = 8 + 8 samples



Palatine Tonsils vs Tongue dorsum

PERMANOVA (Bray−Curtis, permutations within subject): R2 = 0.036, F = 1.57, p = < 0.001, q = < 0.001
PERMDISP p = 0.371 | n = 22 + 22 samples