MAT1575, Classwork11, Fall2026

3.7 Improper Integrals (p.288-300)

1.(Review) **Arrow Notation**: Given a constant a and we have

$x \to a^+$:	x approaches a from the **right** (x is very closed to a but $x \neq a$ and x __>__ a)
$x \to a^-$:	x approaches a from the **left** (x is very closed to a but $x \neq a$ and x __<__ a)
$x \to \infty$:	x approaches infinity (x increases without bound)
$x \to -\infty$:	x approaches negative infinity (x decreases without bound)

2. Evaluate each of the following integrals:

$(a) \int_1^\infty \frac{1}{x}dx$ $\ln|x|\Big|_1^\infty$

$(b) \int_0^4 \frac{1}{\sqrt{4-x}}dx$ ← $[0,4]$ when $x \to 4^-$, $\frac{1}{\sqrt{4-x}} \to \infty$

$(c) \int_{-1}^1 \frac{1}{x^3}dx$ $x \to 0^+$, $\frac{1}{x^3} \to \infty$ $x \to 0^-$, $\frac{1}{x^3} \to -\infty$

3. *Improper integrals*: we define integrals over an infinite interval as well as integrals of functions containing a discontinuity on the interval. Integrals of these types are called Improper Integrals. We examine several techniques for evaluating improper integrals, all of which involve taking limits.

4. Case1: Integrating over an Infinite Interval

Evaluate $(a) \int_1^\infty \frac{1}{x}dx = \lim_{t\to\infty} \int_1^t \frac{dx}{x}$

$= \lim_{t\to\infty} [\ln|x|]_1^t$

$= \lim_{t\to\infty} [\ln|t| - \ln 1]$

$= \lim_{t\to\infty} \ln|t| \to \infty$ or it diverges

$(b) \int_1^\infty \frac{1}{x^2}dx = \lim_{t\to\infty} \int_1^t \frac{dx}{x^2} = \lim_{t\to\infty} \left[-\frac{1}{x}\right]_1^t$

$= \lim_{t\to\infty} \left[-\frac{1}{t} - \left(-\frac{1}{1}\right)\right] = \lim_{t\to\infty} \left[\left(-\frac{1}{t}\right) + 1\right]$

$= 1$

5. Definition for Case1:

(a) Let $f(x)$ be continuous on interval $[a,\infty)$: $\int_a^\infty f(x)\,dx = \lim_{t\to\infty} \int_a^t f(x)dx$ provides this limit exists

(b) Let $f(x)$ be continuous on interval $(-\infty,b]$: $\int_{-\infty}^b f(x)\,dx = \lim_{t\to-\infty} \int_t^b f(x)dx$ provides this limit exists

In each items, if limit exists, then the improper integral is said to converge. If the limit does not exist, then the improper integral is said to diverge.

(c) Let $f(x)$ be continuous on interval $(-\infty,\infty)$: $\int_{-\infty}^\infty f(x)\,dx =$

Then this improper integral is said to converge if

6. Evaluate $\int_{-\infty}^\infty xe^x\,dx$

7. Case2: Integrating a Discontinuous Integrand

Evaluate $(a) \int_0^4 \frac{1}{\sqrt{4-x}}dx$

$(b) \int_{-1}^1 \frac{1}{x^3}dx$

8. Definition for Case2:

(a) Let $f(x)$ be continuous on interval $[a, b)$: $\int_a^b f(x)\,dx =$

(b) Let $f(x)$ be continuous on interval $(a, b]$: $\int_a^b f(x)\,dx =$

In each items, if _____, then _____. If the limit does not exist, then the improper integral is said to _____.

(c) Let $f(x)$ be continuous on interval $[a, b]$ except at a point $c \in (a, b)$: $\int_a^b f(x)\,dx =$

Then this improper integral is said to converge if

9. A Comparison Theorem: Alternative Way to Evaluate Improper Integral.

Consider two continuous functions $f(x)$ and $g(x)$ satisfying $0 \le f(x) \le g(x)$ for $x \ge a$. Then we have

for $t \ge a$

(a) If $\int_a^\infty f(x)\,dx = \lim_{t\to\infty} \int_a^t f(x)\,dx \to \infty$, then $\int_a^\infty g(x)\,dx =$

(b) If $\int_a^\infty g(x)\,dx = \lim_{t\to\infty} \int_a^t g(x)\,dx = L$, where L is a real number, then $\int_a^\infty f(x)\,dx =$

10. Applying the Comparison Theorem.

Use a comparison to show that $\int_1^\infty \frac{1}{xe^x}\,dx$ converges

11. Check if each given integral is an improper integral:

$(a) \int_0^\infty e^{-x}\cos x\,dx$ $(b) \int_0^1 \frac{\ln x}{\sqrt{x}}\,dx$ $(c) \int_1^5 \frac{1}{\sqrt{x-1}}\,dx$ $(d) \int_{-1}^2 \frac{dx}{x^3}$ $(e) \int_3^5 \frac{5}{(x-4)^2}\,dx$