

Mathematics Notebook

November 1, 2022

Preface

I used to blog about mathematics and is planning to revive the work on Medium. Sadly, Medium lack support for LaTeX and it is terribly hard to type equations without scratching my head looking for Unicode characters. The easiest solution is not use Medium but Wordpress with LaTeX plug-in, like Prof. Tao's blog. However, I was also advised to use Medium as it is increasingly popular platform. So I will just type my piece here and put them on medium later, with equations added as images.

Contents

Preface	i
1 Little Gauss	1
2 How to prove it?	3

Chapter 1

Little Gauss

There is this story that is often told in mathematics classes. While the story itself is likely apocryphal, it likely have some pedagogical value. The story goes this way:

There was once a German school where a boy Carl Friedrich made mischief during mathematics lesson. Instead of corporal punishment that was common in that time, the teacher instead decided to give him mathematics assignment to keep him busy. He was asked to add up the numbers from one to a hundred. Most students would diligently start adding and be busy for a while. The young Carl Friedrich, on the other hand, answered after a few minutes. The teacher was surprised at the request to speak, since he had just kept the boy busy. He was all the more astonished when Carl Friedrich said that he had finished the task and was even able to say the correct result (5050).

How had he solved it?

How he did it so fast? Carl Friedrich discovered discovered the following - unfortunately I do not know what coincidence was behind it. He wrote the numbers down like this:

$123\text{latex}\dots991001009998\text{latex}\dots21$

This still doesn't look interesting yet. He would then add up the numbers.

$123\text{latex}\dots991001009998\text{latex}\dots21101101101\text{latex}\dots101101$

Each of them have the sum 101. This looks rather promising.

To sum it up, we write down the numbers from one to one hundred twice, once in increasing order and once in decreasing order, we would then sum them up and we can clearly see that we obtain the sum of $\text{latex}100 \cdot 101$. But we are not finished yet because we counted each numbers twice so we still have to divide the results by two. Then, we would have the sum of numbers from

one to a hundred. And that's exactly how Carl Friedrich proceeded. Do we know Carl Friedrich? Hopefully that's the case, because Carl Friedrich was none other than Carl Friedrich Gauss. One of the most important German mathematicians (if not the most important German mathematician).

Let us talk about the formula

Mathematicians love formulas or should I say the general solution of a problem. The sum of the first n of natural numbers follows the formula:

$$\Sigma = \frac{n \cdot (n+1)}{2}$$

This is not as complicated as it looks. We could for example count the sum of 1 to 150, then we set n equals to 150.

$$\Sigma = \frac{150 \cdot (150+1)}{2} = \frac{150 \cdot (151)}{2} = \frac{22650}{2} = 11325$$

This formula is today is still affectionately referred as "Der Kleine Gauss", German for "Little Gauss". Anyone studying higher mathematics would have to prove the validity of the formula.

Chapter 2

How to prove it?




Figure 2.1:

One of the largest stumbling block in studying mathematics is learning how to prove theorems. In this post, I would share with you 3 of the most commonly used technique with at least one step by step example.

1. Direct proof

Perhaps the most intuitive and straightforward way to write proofs. It goes by "If A , then B " or " A implies B " or mathematically $A \Rightarrow B$.

Example 1 : *The sum of two even numbers is also even.*

Proof. Let x and y be even numbers. Since they are even, by definition they can be rewritten as $2n$ and $2m$ respectively. Thus, the sum $x+y = 2n+2m = 2(n+m)$, which is even number by definition. \square

Example 2 : *Third Binomial Formula*

Proof.

$$(a - b) \cdot (a + b) = a \cdot a + a \cdot b - b \cdot a - b \cdot b \quad (2.1)$$

$$= a^2 + a \cdot b - b \cdot a - b^2 \quad (2.2)$$

$$= a^2 - b^2 \quad (2.3)$$

\square

Example 3 : *Square of odd number is also odd*

2. Indirect proof or proof by contradiction

Example 4 : *Square root of two is irrational*

Bibliography

- [1] M. Artin, *Algebra*. Pearson, 2010.