

Module 4 : Fisherian inference

Marie-Abèle Bind

STAT 140


Spring 2026

Quote #1

"Science must begin with myths, and with the criticism of myths."


Quote #1

"Science must begin with myths, and with the criticism of myths."



Quote #1

"Science must begin with myths, and with the criticism of myths."
Popper, 1962




Quote #2

"All models are wrong but some are useful."


Quote #2

"All models are wrong but some are useful."



Quote #2

"All models are wrong but some are useful."
Box, 1979



Quote #3

"Truth is much too complicated to allow anything but approximations."

Quote #3

"Truth is much too complicated to allow anything but approximations."



Quote #3

"Truth is much too complicated to allow anything but approximations."
von Neumann, 1947




A Popperian perspective in defense of ...

- "Science must begin with myths, and with the criticism of myths."

A Popperian perspective in defense of ...

- "Science must begin with myths, and with the criticism of myths."
- A p-value, as described by Fisher (1925) in the context of a randomized experiment is a mathematical formalization of the Popperian perspective : a **stochastic proof by contradiction** of the plausibility of an empirical hypothesis when confronted with observed data.



- Desires to prohibit the reporting of p-values to help assess the compatibility of proposed models with observed data (Traffimow and Marks, 2015).



Potential outcomes (simplest situation)

- Consider N units, indexed by i , in an experiment randomized to a binary treatment W , either active (i.e., $W_i=1$) or control (i.e., $W_i=0$).

Potential outcomes (simplest situation)

- Consider N units, indexed by i , in an experiment randomized to a binary treatment W , either active (i.e., $W_i=1$) or control (i.e., $W_i=0$).
- An outcome variable Y is measured after exposure to the treatment.

Potential outcomes (simplest situation)

- Consider N units, indexed by i , in an experiment randomized to a binary treatment W , either active (i.e., $W_i=1$) or control (i.e., $W_i=0$).
- An outcome variable Y is measured after exposure to the treatment.
- No hidden treatment and no interference between units (SUTVA).
 $Y_i(W_i=w)$ is a function of unit i and treatment w .

Potential outcomes (simplest situation)

- Consider N units, indexed by i , in an experiment randomized to a binary treatment W , either active (i.e., $W_i=1$) or control (i.e., $W_i=0$).
- An outcome variable Y is measured after exposure to the treatment.
- No hidden treatment and no interference between units (SUTVA).
 $Y_i(W_i=w)$ is a function of unit i and treatment w .
- For each unit $i=1, \dots, N$:
 - $Y_i(W_i=1)$ =value of Y when i is exposed to the active treatment
 - $Y_i(W_i=0)$ =value of Y when i is exposed to the control treatment.

Potential outcomes (simplest situation)

- Consider N units, indexed by i , in an experiment randomized to a binary treatment W , either active (i.e., $W_i=1$) or control (i.e., $W_i=0$).
- An outcome variable Y is measured after exposure to the treatment.
- No hidden treatment and no interference between units (SUTVA).
 $Y_i(W_i=w)$ is a function of unit i and treatment w .
- For each unit $i=1, \dots, N$:
 - $Y_i(W_i=1)$ =value of Y when i is exposed to the active treatment
 - $Y_i(W_i=0)$ =value of Y when i is exposed to the control treatment.
- Because of SUTVA, the science table simplifies to 2 columns and N rows.

Science and observed tables

- Science table :

i	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	$Y_1(W_1=0)$	$Y_1(W_1=1)$
...
N	$Y_N(W_N=0)$	$Y_N(W_N=1)$

Science and observed tables

- Science table :

i	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	$Y_1(W_1=0)$	$Y_1(W_1=1)$
...
N	$Y_N(W_N=0)$	$Y_N(W_N=1)$

- Observed data table :

i	W_i^{obs}	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	0	Y_1^{obs}	?
...
N	1	?	Y_N^{obs}

Science and observed tables

- Science table :

i	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	$Y_1(W_1=0)$	$Y_1(W_1=1)$
...
N	$Y_N(W_N=0)$	$Y_N(W_N=1)$

- Observed data table :

i	W_i^{obs}	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	0	Y_1^{obs}	?
...
N	1	?	Y_N^{obs}

$$Y_i^{obs} = W_i^{obs} Y_i(W_i = 1) + (1 - W_i^{obs}) Y_i(W_i = 0)$$

Stochastic proof by contradiction

- We follow three steps to assess the plausibility of a Fisher sharp null hypothesis (H_0), *the myth*.

Stochastic proof by contradiction

- We follow three steps to assess the plausibility of a **Fisher sharp null hypothesis** (H_0), *the myth*.

$$\underline{H_0} : \forall i \in \{1, \dots, N\} \quad Y_i(W_i=1) = Y_i(W_i=0)$$

Stochastic proof by contradiction

- We follow three steps to assess the plausibility of a Fisher sharp null hypothesis (H_0), *the myth*.
 - 1 Guided by scientific considerations, choose an appropriate test statistic, T , and define more extreme (e.g., which group is expected to have a better outcome).

Stochastic proof by contradiction

- We follow three steps to assess the plausibility of a **Fisher sharp null hypothesis** (H_0), *the myth*.
 - 1 Guided by scientific considerations, choose an appropriate **test statistic**, T , and define more extreme (e.g., which group is expected to have a better outcome).

Example of T :
$$T(W, Y^{obs}) = \frac{1}{N_1} \sum_{i: W_i=1} Y_i^{obs} - \frac{1}{N_0} \sum_{i: W_i=0} Y_i^{obs}$$


- Statistic : function of the observed data.
- Sensitive to expected departures from the null hypothesis.
- Example of more extreme : $T(W, Y^{obs}) > T^{obs}$

Stochastic proof by contradiction

- We follow three steps to assess the plausibility of a Fisher sharp null hypothesis (H_0), *the myth*.
 - 1 Guided by scientific considerations, choose an appropriate test statistic, T , and define more extreme (e.g., which group is expected to have a better outcome).
 - 2 Assuming H_0 , calculate the value of T for all possible randomized allocations to obtain the null randomization distribution of T .

Stochastic proof by contradiction


- We follow three steps to assess the plausibility of a **Fisher sharp null hypothesis** (H_0), *the myth*.
 - 1 Guided by scientific considerations, choose an appropriate **test statistic**, T , and define more extreme (e.g., which group is expected to have a better outcome).
 - 2 **Assuming H_0** , calculate the value of T for all possible randomized allocations to obtain the null randomization distribution of T .



Example of a null randomization distribution

Stochastic proof by contradiction

- We follow three steps to assess the plausibility of a **Fisher sharp null hypothesis** (H_0), *the myth*.
 - ① Guided by scientific considerations, choose an appropriate **test statistic**, T , and define more extreme (e.g., which group is expected to have a better outcome).
 - ② **Assuming H_0** , calculate the value of T for all possible randomized allocations to obtain the null randomization distribution of T .



- ③ **Locate** the observed value of the test statistic, T^{obs} , in the null randomization distribution constructed in Step 2.


Stochastic proof by contradiction

- We follow three steps to assess the plausibility of a **Fisher sharp null hypothesis** (H_0), *the myth*.
 - ① Guided by scientific considerations, choose an appropriate **test statistic**, T , and define more extreme (e.g., which group is expected to have a better outcome).
 - ② **Assuming H_0** , calculate the value of T for all possible randomized allocations to obtain the null randomization distribution of T .
 - ③ **Locate** the observed value of the test statistic, T^{obs} , in the null randomization distribution constructed in Step 2.


The Fisher-exact p-value corresponds to the proportion of values of the test statistic that are as extreme or more extreme than the observed value of that test statistic.

If computationally impossible, approximate?

For computational reasons, when N is large, the null randomization distribution of the test statistic may be approximated by randomly drawing fewer permutations than $N_{\text{randomizations}}$.





Approximated p-value (left) = $\frac{1}{1,000} \sum_{r=1}^{1,000} \mathbb{1}(T^{(r)} \geq T^{obs}) = 0.20$



Fisher-exact p-value (right) = $\frac{1}{N_{\text{randomizations}}} \sum_{r=1}^{N_{\text{randomizations}}} \mathbb{1}(T^{(r)} \geq T^{obs}) = 0.07$

Smallest and largest attainable p-values?




$$\text{Fisher-exact p-value} = \frac{1}{N_{\text{randomizations}}} \sum_{r=1}^{N_{\text{randomizations}}} \mathbb{1}(T^{(r)} \geq T^{\text{obs}})$$

Smallest p-value is important

B vitamins attenuate the epigenetic effects of ambient fine particles in a pilot human intervention trial


PNAS March 28, 2017 114 (13) 3503-3508; first published March 13, 2017



Smallest p-value is important

B vitamins attenuate the epigenetic effects of ambient fine particles in a pilot human intervention trial

PNAS March 28, 2017 114 (13) 3503-3508; first published March 13, 2017



$$\frac{1}{\sqrt{10}} \approx 0.00098$$