

Module 3 : Assignment mechanism

Marie-Abèle Bind

STAT 140

Spring 2026

Single-factor randomized experiment

- Let i index participants, $i \in \{1, \dots, N\}$.

Single-factor randomized experiment

- Let i index participants, $i \in \{1, \dots, N\}$.
- Let W_i denote the unit-level treatment assignment :

Single-factor randomized experiment

- Let i index participants, $i \in \{1, \dots, N\}$.
- Let W_i denote the unit-level treatment assignment :

$$W_i = \begin{cases} 0 & \text{if unit } i \text{ is assigned to placebo} \\ 1 & \text{if unit } i \text{ is assigned to active treatment} \end{cases}$$

Potential outcomes framework (Rubin, 1974)

- Potential outcomes :

- $Y_i(W_i = 0)$ denotes the outcome Y under the control treatment
- $Y_i(W_i = 1)$ denotes the outcome Y under the active treatment

Potential outcomes framework (Rubin, 1974)

- Potential outcomes :

- $Y_i(W_i = 0)$ denotes the outcome Y under the control treatment
- $Y_i(W_i = 1)$ denotes the outcome Y under the active treatment

- Observed and missing outcomes :

- $$Y_i^{obs} = \begin{cases} Y_i(W_i = 0) & \text{if } W_i^{obs} = 0 \\ Y_i(W_i = 1) & \text{if } W_i^{obs} = 1 \end{cases}$$

Potential outcomes framework (Rubin, 1974)

- Potential outcomes :

- $Y_i(W_i = 0)$ denotes the outcome Y under the control treatment
- $Y_i(W_i = 1)$ denotes the outcome Y under the active treatment

- Observed and missing outcomes :

- $Y_i^{obs} = \begin{cases} Y_i(W_i = 0) & \text{if } W_i^{obs} = 0 \\ Y_i(W_i = 1) & \text{if } W_i^{obs} = 1 \end{cases}$
- $Y_i^{mis} = \begin{cases} Y_i(W_i = 1) & \text{if } W_i^{obs} = 0 \\ Y_i(W_i = 0) & \text{if } W_i^{obs} = 1 \end{cases}$

Potential outcomes framework (Rubin, 1974)

- Potential outcomes :

- $Y_i(W_i = 0)$ denotes the outcome Y under the control treatment
- $Y_i(W_i = 1)$ denotes the outcome Y under the active treatment

- Observed and missing outcomes :

- $Y_i^{obs} = \begin{cases} Y_i(W_i = 0) & \text{if } W_i^{obs} = 0 \\ Y_i(W_i = 1) & \text{if } W_i^{obs} = 1 \end{cases}$
- $Y_i^{mis} = \begin{cases} Y_i(W_i = 1) & \text{if } W_i^{obs} = 0 \\ Y_i(W_i = 0) & \text{if } W_i^{obs} = 1 \end{cases}$

- One-to-one correspondence :

- $Y_i(W_i = 0) = \begin{cases} Y_i^{obs} & \text{if } W_i^{obs} = 0 \\ Y_i^{mis} & \text{if } W_i^{obs} = 1 \end{cases}$

Potential outcomes framework (Rubin, 1974)

- Potential outcomes :

- $Y_i(W_i = 0)$ denotes the outcome Y under the control treatment
- $Y_i(W_i = 1)$ denotes the outcome Y under the active treatment

- Observed and missing outcomes :

- $Y_i^{obs} = \begin{cases} Y_i(W_i = 0) & \text{if } W_i^{obs} = 0 \\ Y_i(W_i = 1) & \text{if } W_i^{obs} = 1 \end{cases}$
- $Y_i^{mis} = \begin{cases} Y_i(W_i = 1) & \text{if } W_i^{obs} = 0 \\ Y_i(W_i = 0) & \text{if } W_i^{obs} = 1 \end{cases}$

- One-to-one correspondence :

- $Y_i(W_i = 0) = \begin{cases} Y_i^{obs} & \text{if } W_i^{obs} = 0 \\ Y_i^{mis} & \text{if } W_i^{obs} = 1 \end{cases}$
- $Y_i(W_i = 1) = \begin{cases} Y_i^{mis} & \text{if } W_i^{obs} = 0 \\ Y_i^{obs} & \text{if } W_i^{obs} = 1 \end{cases}$

Causal inference as a missing data problem (1976)

- Science table provides observable outcomes :

i	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	$Y_1(W_1=0)$	$Y_1(W_1=1)$
...
N	$Y_N(W_N=0)$	$Y_N(W_N=1)$

Causal inference as a missing data problem (1976)

- Science table provides observable outcomes :

i	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	$Y_1(W_1=0)$	$Y_1(W_1=1)$
...
N	$Y_N(W_N=0)$	$Y_N(W_N=1)$

- Observed data table :

i	W_i^{obs}	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	0	Y_1^{obs}	?
...
N	1	?	Y_N^{obs}

Causal inference as a missing data problem (1976)

- Science table provides observable outcomes :

i	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	$Y_1(W_1=0)$	$Y_1(W_1=1)$
...
N	$Y_N(W_N=0)$	$Y_N(W_N=1)$

- Observed data table :

i	W_i^{obs}	$Y_i(W_i=0)$	$Y_i(W_i=1)$
1	0	Y_1^{obs}	?
...
N	1	?	Y_N^{obs}

$$Y_i^{obs} = W_i^{obs} Y_i(W_i = 1) + (1 - W_i^{obs}) Y_i(W_i = 0)$$

Assignment mechanism

- The fundamental problem of causal inference is the presence of missing data :
 - Recall, for each unit, we can observe at most one of the potential outcomes.
- A key component in a causal analysis is the **assignment mechanism** :
 - The process that determines which units receive which treatments, hence which potential outcomes are observed, and which potential outcomes are missing.
- Formally, the assignment mechanism describes the probability of any vector of assignments as a function of all covariates and all potential outcomes.
- Taxonomy of assignment mechanisms (Rubin, 1974).

Assignment mechanism

- The assignment mechanism is the function that assigns probabilities to all 2^N possible values for the N-vector of assignments \mathbf{W} (i.e., each unit can be assigned to treatment or control) given the N-vector of potential outcomes $\mathbf{Y}(0)$ and $\mathbf{Y}(1)$, the $N \times K$ matrix of covariates \mathbf{X} .
- Given a population of N units, the assignment mechanism is a row-exchangeable function $P(\mathbf{W}|\mathbf{X}, \mathbf{Y}(0), \mathbf{Y}(1))$, taking values in $[0,1]$, satisfying :

$$\sum_{\mathbf{W} \in \{0,1\}^N} P(\mathbf{W}|\mathbf{X}, \mathbf{Y}(0), \mathbf{Y}(1)) = 1$$

- for all $\mathbf{X}, \mathbf{Y}(0)$, and $\mathbf{Y}(1)$
- $\{0,1\}^N$ is the set of all N-vectors with all elements equal to 0 or 1.

Unit assignment probability

- The unit-level assignment probability for unit i is :

$$p_i(\mathbf{X}, \mathbf{Y}(0), \mathbf{Y}(1)) = \sum_{\mathbf{W}: W_i=1} P(\mathbf{W}|\mathbf{X}, \mathbf{Y}(0), \mathbf{Y}(1))$$

Taxonomy

Classical randomized experiments : the assignment mechanism satisfies all three restrictions below :

- **Individualistic** assignment :
 - This limits the dependence of a particular unit's assignment probability on the values of covariates and potential outcomes for other units.

Taxonomy

Classical randomized experiments : the assignment mechanism satisfies all three restrictions below :

- **Individualistic** assignment :
 - This limits the dependence of a particular unit's assignment probability on the values of covariates and potential outcomes for other units.
- **Probabilistic** assignment :
 - This requires the assignment mechanism to imply a non-zero probability for each treatment value, for every unit.
 - For all units $i = 1, \dots, N$:

$$0 < p_i(\mathbf{X}, \mathbf{Y}(0), \mathbf{Y}(1)) < 1$$

Taxonomy

Classical randomized experiments : the assignment mechanism satisfies all three restrictions below :

- **Individualistic** assignment :
 - This limits the dependence of a particular unit's assignment probability on the values of covariates and potential outcomes for other units.
- **Probabilistic** assignment :
 - This requires the assignment mechanism to imply a non-zero probability for each treatment value, for every unit.
 - For all units $i = 1, \dots, N$:

$$0 < p_i(\mathbf{X}, \mathbf{Y}(0), \mathbf{Y}(1)) < 1$$

- **Unconfounded** assignment :
 - This disallows dependence of the assignment mechanisms on the potential outcomes.

Assignment mechanism

- The assignment mechanism differentiates randomized experiments from observational studies.

Assignment mechanism

- The assignment mechanism differentiates randomized experiments from observational studies.
- The assignment, for example, of active treatment vs. control for each unit is **controlled** by the experimenter in randomized experiments, in contrast to observational studies.

Assignment mechanism

- The assignment mechanism differentiates randomized experiments from observational studies.
- The assignment, for example, of active treatment vs. control for each unit is **controlled** by the experimenter in randomized experiments, in contrast to observational studies.
- Classical randomized experiments :
 - The assignment mechanism is individualistic, probabilistic, and unconfounded.
 - The researcher **knows and controls** the functional form of the assignment mechanism.

Completely randomized treatment assignment

- In randomized experiments, scientists sometimes choose a **completely randomized mechanism** to assign participants to placebo vs. active treatment.

Completely randomized treatment assignment

- In randomized experiments, scientists sometimes choose a **completely randomized mechanism** to assign participants to placebo vs. active treatment.
- Here, out of N participants, N_T are randomly assigned to the active treatment and N_C ($N_C = N - N_T$) are assigned to placebo.

Completely randomized treatment assignment

- In randomized experiments, scientists sometimes choose a **completely randomized mechanism** to assign participants to placebo vs. active treatment.
- Here, out of N participants, N_T are randomly assigned to the active treatment and N_C ($N_C = N - N_T$) are assigned to placebo.
- There are " N choose N_T ", i.e., $\binom{N}{N_T}$ number of ways to do this.

Completely randomized treatment assignment

- In randomized experiments, scientists sometimes choose a **completely randomized mechanism** to assign participants to placebo vs. active treatment.
- Here, out of N participants, N_T are randomly assigned to the active treatment and N_C ($N_C = N - N_T$) are assigned to placebo.
- There are " N choose N_T ", i.e., $\binom{N}{N_T}$ number of ways to do this.
- $$\binom{N}{N_T} = \frac{N!}{N_T!(N-N_T)!} = \frac{N(N-1)\dots(N-N_T+1)}{N_T(N_T-1)\dots\times 1}$$

Completely randomized treatment assignment

- In randomized experiments, scientists sometimes choose a **completely randomized mechanism** to assign participants to placebo vs. active treatment.
- Here, out of N participants, N_T are randomly assigned to the active treatment and N_C ($N_C = N - N_T$) are assigned to placebo.
- There are " N choose N_T ", i.e., $\binom{N}{N_T}$ number of ways to do this.
- Each participant has the same chance of receiving the active treatment, i.e., $\frac{N_T}{N}$.

Completely randomized experiment

- Assignment mechanism :

$$P(\mathbf{W}|\mathbf{X}, \mathbf{Y}(0), \mathbf{Y}(1)) = \begin{cases} \binom{N}{N_t}^{-1} & \text{if } \sum_{i=1}^N W_i = N_t \\ 0 & \text{otherwise.} \end{cases}$$

The "sample" function in R

- The "`sample`" function in R creates a vector of 0's and 1's with the same numbers of 0's and 1's, in random order.

The "sample" function in R

- The "sample" function in R creates a vector of 0's and 1's with the same numbers of 0's and 1's, in random order.

```
> sample(c(0,0,0,1,1))  
[1] 0 1 1 0 0
```

The "sample" function in R

- The "sample" function in R creates a vector of 0's and 1's with the same numbers of 0's and 1's, in random order.

```
> sample(c(0,0,0,1,1))  
[1] 1 0 0 0 1
```

Bernoulli assignment mechanism

- The Bernoulli assignment is an alternative assignment mechanism using independent coin tosses.

Bernoulli assignment mechanism

- The Bernoulli assignment is an alternative assignment mechanism using independent coin tosses.
- Each study participant is assigned to treatment with probability p_i (e.g., for each participant i , $p_i = 0.5$).

Bernoulli assignment mechanism

- The Bernoulli assignment is an alternative assignment mechanism using independent coin tosses.
- Each study participant is assigned to treatment with probability p_i (e.g., for each participant i , $p_i = 0.5$).
- There are 2^N number of ways to randomize the N study participants to the active treatment.

Bernoulli assignment mechanism

- The Bernoulli assignment is an alternative assignment mechanism using independent coin tosses.
- Each study participant is assigned to treatment with probability p_i (e.g., for each participant i , $p_i = 0.5$).
- There are 2^N number of ways to randomize the N study participants to the active treatment.
- Possible assignment vectors are $W = (W_1 = 0, \dots, W_N = 0)$ and $W = (W_1 = 1, \dots, W_N = 1)$.

Bernoulli trials

- Bernoulli experiment with a fair coin :

For each unit,

$$P(W_i = 1 | X_i, Y_i(0), Y_i(1)) = 0.5$$

- Assignment mechanism :

$$P(\mathbf{W} | \mathbf{X}, \mathbf{Y}(0), \mathbf{Y}(1)) = \prod_{i=1}^N P(W_i = 1 | X_i)^{W_i} [1 - P(W_i = 1 | X_i)]^{1-W_i}$$

Paired randomized experiment

- Each unit has probability $\frac{1}{2}$ of being assigned to the treatment group.
- Assignment mechanism :

$$P(\mathbf{W}|\mathbf{X},\mathbf{Y}(0),\mathbf{Y}(1)) = \begin{cases} 2^{-N/2} & \text{if } \mathbf{W} \in W^+ \\ 0 & \text{otherwise.} \end{cases}$$