



IMDELLTRA®
(tarlatamab-dlle)



Extensive-Stage Small Cell Lung Cancer

**with disease progression on or after platinum-based chemotherapy*

Mechanism of Action:

IMDELLTRA® (Tarlatamab-dlle) is a bispecific T-cell engager (BiTE) therapy that helps the immune system recognize and attack small cell lung cancer (SCLC) cells. It works by binding to delta-like ligand 3 (DLL3), a protein that is normally found inside healthy cells but is abnormally present on the surface of SCLC cells. This abnormal surface expression makes DLL3 an important target in SCLC.

The drug is built with two single-chain variable fragment (scFv) binding domains joined by a flexible linker. One domain attaches to DLL3 on tumor cells, while the other binds to CD3 receptors on T cells. This connection brings T cells into direct contact with DLL3-positive cancer cells, leading to T-cell activation, release of inflammatory cytokines, and subsequent killing (lysis) of the tumor cells.

Pre & Post-Infusion

Labs:

Pre-Infusion only: hCG

Pre-Infusion and prior to each dose: CBC, ALT/AST, Bilirubin

Other affected levels: lymphocytes (L), sodium (L), neutrophils (L), hemoglobin (L), potassium (L), white blood cells (L), platelets (L), alanine aminotransferase (L), uric acid (H), activated partial thromboplastin time (H), aspartate aminotransferase (H)

Pre-Medications (Cycle 1): *reduces risk of cytokine release syndrome (CRS)*

Dexamethasone 8 mg IV within 1 hour prior to infusion; Day 1 and Day 8

Compounding Information

Step 1: Reconstituting

Reconstitute IMDELLTRA® with Sterile Water for injection. *Do not use IV Solution Stabilizer (IVSS) to reconstitute IMDELLTRA®.*

The IV Solution Stabilizer (IVSS) is used to coat the intravenous bag prior to addition of reconstituted IMDELLTRA® to prevent adsorption of IMDELLTRA® to IV bags and IV tubing.

- For 1 mg dose, use 1.3 mL Sterile Water to reconstitute.
- For 10 mg dose, use 4.4 mL Sterile Water to reconstitute.

Reconstitution:

- Using a needle and syringe filled with the required amount of sterile water, inject the sterile water against the glass vial. Avoid injecting the water directly onto the powder to prevent foaming.
- Gently swirl the contents to mix. Do not shake.
- Inspect for particulate matter and discoloration prior to administration. Inspect that the solution is clear to opalescent, colorless to slightly yellow. *Do not use if the solution is cloudy or has particulates.*
- Further dilute reconstituted IMDELLTRA® within 4 hours of reconstitution or discarded.

Step 2: Preparing the infusion bag

Using a 250 mL prefilled bag of 0.9% Sodium Chloride for Injection, withdraw the appropriate amount of sodium chloride and discard.

- For 1 mg dose, withdraw 14 mL 0.9% NaCl from IV bag
- For 10 mg dose, withdraw 17 mL 0.9% NaCl from IV bag

Inject IV Solution Stabilizer (IVSS) into the 250 mL 0.9% Sodium Chloride infusion bag. Gently mix the contents of the infusion bag to avoid foaming. Do not shake.

- For both 1 mg and 10 mg dose, add 13 mL IVSS to IV bag

Dilute the reconstituted IMDELLTRA® into the infusion bag. Transfer the required volume of reconstituted IMDELLTRA® to the infusion bag (containing IVSS). Gently mix the contents of the bag. Do not shake.

- For 1 mg dose, add 1.1 mL reconstituted IMDELLTRA® to IV bag
- For 10 mg dose, add 4.2 mL reconstituted IMDELLTRA® to IV bag

Finally, remove air from the prepared IV bag using an empty syringe to avoid foaming.

Step 3: Priming the IV tubing

Prime intravenous tubing with either 0.9% NaCl for injection or with the final prepared product.


Step 3: Administration & Storage

Administer reconstituted and diluted IMDELLTRA® immediately.

Maximum storage time: *includes total duration from the time of reconstitution of the vial of IMDELLTRA to the end of the infusion*

- For room temperature (20°C-25°C / 68°F-77°F): 8 hours
- For refrigeration (2°C-8°C / 36°F-46°F): 7 days

Administration



The IV catheter for concomitant medications administration can be used to administer the IMDELLTRA® infusion. To ensure patency, flush the IV catheter over 3-5 mins using 0.9% Sodium Chloride for Injection. Administer the reconstituted and diluted IMDELLTRA® as an IV infusion at a constant flow rate of 250 mL/hour for 1 hour using an infusion pump. The pump should be programmable, lockable, non-elastomeric, and have an alarm.

Dosing

Step-up Dosing Schedule	Day 1 ^a : 1 mg
	Day 8 ^a : 10 mg
Cycle 1	Day 15: 10 mg
	Monitoring (Day 1 ^a and 8 ^a): Monitor patients from the start of the infusion for 22-24 hours on Cycle 1, Day 1 and Cycle 1, Day 8 in a healthcare setting. Recommend that patients remain within 1-hour of an appropriate healthcare setting for a total of 48 hours from start of the infusion, accompanied by a caregiver.
	Monitoring (Day 15): Observe patients for 6-8 hours post-infusion ^b .
Cycle 2	Day 1 and Day 15: 10 mg
	Monitoring: Observe patients for 6-8 hours post-infusion ^b .
Cycles 3 and 4	Day 1 and Day 15: 10 mg
	Monitoring: Observe patients for 3-4 hours post-infusion ^b .
Cycle 5 and subsequent infusions	Day 1 and Day 15: 10 mg
	Monitoring: Observe patients for 2 hours post-infusion ^b .

a= Administer recommended concomitant medications before and after Cycle 1 IMDELLTRA[®] infusions as described.

b= Extended monitoring in a healthcare setting is not required unless the patient experiences Grade ≥ 2 CRS, ICANS or neurological toxicity during prior treatments. See *Monitoring* below.

Restarting IMDELLTRA[®] After Dosage Delay

1 mg on Cycle 1, Day 1	2 weeks or less (≤ 14 days): • Administer IMDELLTRA® 10 mg, then resume with the planned dosage schedule. Greater than 2 weeks (> 14 days): • Administer IMDELLTRA® step-up dose 1 mg. If tolerated, increase to 10 mg 1 week later. Then resume with the planned dosage schedule.
10 mg on Cycle 1, Day 8	3 weeks or less (≤ 21 days): • Administer IMDELLTRA® 10 mg, then resume with the planned dosage schedule. Greater than 3 weeks (> 21 days): • Administer IMDELLTRA® step-up dose 1 mg. If tolerated, increase to 10 mg 1 week later. Then resume with the planned dosage schedule
10 mg on Cycle 1, Day 15 and subsequent Cycles every 2 weeks thereafter	4 weeks or less (≤ 28 days): • Administer IMDELLTRA® 10 mg, then resume with the planned dosage schedule. Greater than 4 weeks (> 28 days): • Administer IMDELLTRA® step-up dose 1 mg. If tolerated, increase to 10 mg 1 week later. Then resume with the planned dosage schedule.



Adverse Reactions

Cytokine release syndrome (CRS)

Fatigue

Pyrexia

Dysgeusia

Decreased appetite

Musculoskeletal pain

Constipation

Anemia

Warnings

Neurologic Toxicity; ICANS

Cytopenias

Infections

Hepatotoxicity

Hypersensitivity

Embryo-Fetal Toxicity



IMDELLTRA® Package Insert (PI) Link

www.imdelltrahcp.com
