


1. Most proteins that are secreted from a cell must be transported to the endoplasmic reticulum (ER) either during translation or after translation.

A. Describe the function of ribosomes.

For proteins transported during translation, this process begins in the cytosol and pauses when a specific sequence of amino acids is translated. The translation complex is then transported to the surface of the ER where translation continues. Proteins that are transported after translation are translated entirely in the cytosol and then transported to the ER. In both instances, the translated proteins enter the ER through a protein channel in the membrane of the ER.

Researchers studying the two types of protein transport identified that the ER membrane protein SR is necessary for transport during translation, while the ER membrane protein Sec62 is necessary for transport after translation. To investigate which transport mechanism is used for different proteins, researchers first created small interfering RNAs (siRNAs) that reduce expression of either SR or Sec62. They then treated groups of cells with either the SR siRNA or the Sec62 siRNA and determined the relative amount of SR and Sec62 protein in each group of cells compared with cells treated with a control siRNA (Figure 1).

Figure 1. Average relative amounts of Sec62 and SR proteins in cells treated with control siRNA, Sec62 siRNA, or SR siRNA. Error bars represent $\pm SE_{\bar{x}}$.



B.

- Identify** the dependent variable in the experiments shown in Figure 1.
- Justify** why the researchers included the control of measuring the relative amounts of both Sec62 and SR proteins in cells that were treated with Sec62 siRNA only (data shown in Figure 1).
- Based on Figure 1, **describe** the effect on the production of SR protein when cells are treated with Sec62 siRNA.