

Solutions to R Worksheet 3.

Q1(a)

```
> chivals <- rchisq(1000,df=18)
```

Q1(b)

```
> hist(chivals)
```

Q1(c)

```
> min(chivals)
[1] 4.895851
```

```
> max(chivals)
[1] 44.4928
```

Q1(d)

```
> hist(chivals,breaks=seq(floor(min(chivals)),floor(max(chivals)+1),1))
> hist(chivals,breaks=seq(floor(min(chivals)),floor(max(chivals)+1),0.5))
> hist(chivals,breaks=seq(floor(min(chivals)),floor(max(chivals)+1),0.25))
```

Q1(e)

```
> for(i in 1:24) {
+   chivals_chivals+rchisq(1000,df=18)
+ }
> chiavg <- chivals/25
> hist(chiavg, breaks=seq(floor(min(chiavg)),floor(max(chiavg)+1),0.5))
> mean(chiavg)
[1] 17.99582
```

```
# The mean should be equal to 18, the degrees of freedom (which is the
# expectation of a chi-squared random variable)
```

```
> sqrt(var(chiavg))
[1] 1.201205
```

```
# The variance of these values should be  $36/25 = 1.44$ , twice the degrees of
# freedom (which is the variance of a chi-squared random variable) over the
# number of summands in the average. Thus, the standard deviation out to
# be 1.2, which is the square root of 1.44.
```

Q1(f)

```
> grid <- seq(floor(min(chiavg)),floor(max(chiavg)+1),0.1)
> normals <- dnorm(grid,18,1.2)
> hist(chiavg, breaks=seq(floor(min(chiavg)),floor(max(chiavg)+1),0.5))
> lines(grid,normals)
> lines(grid,1000*0.5*normals)
```

Q1(g)

```
> qqnorm(chiavg)
```

Q2

```
# The function was given in the worksheet.  
# To test it, we might look at the "worksheet2.women2" dataframe.
```

```
> worksheet2.women <- as.data.frame(worksheet2.women)  
> attach(women.df)  
> influence(height,1)  
[1] 0.4533333 -4.2446111  
> influence(height,2)  
[1] 0.1422222 -6.3290556  
> influence(height,3)  
[1] -0.9355556 3.2901111  
> influence(height,4)  
[1] 1.731111 27.156778  
> influence(height,5)  
[1] -0.7911111 0.4842778  
> influence(height,6)  
[1] 0.008888889 -6.555722222  
> influence(height,7)  
[1] 0.1311111 -6.3632222  
> influence(height,8)  
[1] -1.002222 4.743444  
> influence(height,9)  
[1] 0.2866667 -5.6321111  
> influence(height,10)  
[1] -0.02444444 -6.54988889
```

```
# Looks like point 4 is one to look out for
```

```
> influence(weight,1)  
[1] 1.303333 8.944000  
> influence(weight,2)  
[1] -0.1411111 -9.9421111  
> influence(weight,3)  
[1] -0.3855556 -8.4937778  
> influence(weight,4)  
[1] 0.5588889 -6.6521111  
> influence(weight,5)  
[1] -0.7188889 -4.3521111  
> influence(weight,6)  
[1] -0.7855556 -3.2237778  
> influence(weight,7)  
[1] -0.2966667 -9.1760000  
> influence(weight,8)  
[1] -1.141111 4.482889  
> influence(weight,9)  
[1] -0.430 -8.086  
> influence(weight,10)  
[1] 2.036667 36.499000  
> influence(weight,c(1,10))  
[1] 3.75750 60.55346  
> influence(weight,c(1,2))  
[1] 1.307500 -1.396536  
> influence(weight,c(2,10))  
[1] 2.13250 30.18918
```

```
# Looks like point 10 is the one to look out for here, and in combination  
# with point 1 they seem to have a rather large effect.
```