

STAT 6046 Tutorial Week 6

By Isaac Pan
The Australian National University



Today's plan

Brief review of course material

Go through selective tutorial questions



Equation of value

Basic relationship

General formula

$$\sum_{k=1}^{n} I_{t_k} \cdot (1+i)^{-t_k} + \int_{0}^{n} \rho_I(t) \cdot (1+i)^{-t} dt = \sum_{k=1}^{n} O_{t_k} \cdot (1+i)^{-t_k} + \int_{0}^{n} \rho_O(t) \cdot (1+i)^{-t} dt$$



Solving for unknown time

Annuity payments' value equals to a fixed value

$$A = Bs_{\overline{n|i}} \text{ or } A = Ba_{\overline{n|i}}$$

$$n = \frac{-\ln\left(1 - \frac{Ai}{B}\right)}{\ln(1+i)} = \frac{-\ln\left(1 - \frac{Ai}{B}\right)}{\delta}$$

$$n = \frac{\ln\left(\frac{Ai}{B} + 1\right)}{\ln(1+i)} = \frac{\ln\left(\frac{Ai}{B} + 1\right)}{\delta}$$



Solving for unknown time

Annuity payments' value equals to a time-varying value

$$90s_{\overline{n}|0.01} > 100n$$
 $\Rightarrow 90\frac{(1.01)^n - 1}{0.01} > 100n$

Linear Interpolation:
$$\frac{f(n_0) - f(n_1)}{f(n_2) - f(n_1)} \cong \frac{n_0 - n_1}{n_2 - n_1}$$

$$\Rightarrow n_0 \cong n_1 + \frac{f(n_0) - f(n_1)}{f(n_2) - f(n_1)} \cdot (n_2 - n_1)$$

 Note: This approximation works best if the trial values are close to the true value.



Solving for unknown interest rate

Quadratic formula:

$$ax^2 + bx + c = 0$$

• Solution:

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$



Inflation

Real interest rate:

$$i_{real} = \frac{i - r}{1 + r}$$

- Time varying inflation: CPI
- At time 0, purchasing power of \$1 at time

t:
$$\$1 * \frac{CPI_0}{CPI_1}$$