

PLEASE HAND IN

UNIVERSITY OF TORONTO
FACULTY OF ARTS AND SCIENCE

APRIL 2010 EXAMINATIONS

CSC 165H1S
DURATION — 3 HOURS

PLEASE HAND IN

ONE 8.5" × 11" HANDWRITTEN (BOTH SIDES) AID SHEET ALLOWED

STUDENT NUMBER: _____

LAST NAME: _____

FIRST NAME: _____

*Do NOT turn this page until you have received the signal to start.
(In the meantime, please fill out the identification section above,
and read the instructions below.)*

This exam consists of 9 questions on 14 pages (including this one).
*When you receive the signal to start, please make sure that your copy
of the exam is complete.*

Please answer questions in the space provided. You will earn 20% for
any question you leave blank or write "I cannot answer this question,"
on. You will earn substantial part marks for writing down the outline of
a solution and indicating which steps are missing.

Write your student number at the bottom of pages 2-14 of this exam.

1: _____/12

2: _____/10

3: _____/10

4: _____/10

5: _____/10

6: _____/10

7: _____/10

8: _____/10

9: _____/10

TOTAL: _____/92

Good Luck!

QUESTION 1. [12 MARKS]

Consider the three sentences below:

$$S1: \quad \forall x \in X, P(x) \Rightarrow Q(x)$$

$$S2: \quad \exists x \in X, P(x) \Rightarrow Q(x)$$

$$S3: \quad \forall x \in X, Q(x) \Rightarrow P(x)$$

In each of the subquestions below “devise an example of set X and predicates P and Q ” means you must suggest elements for X and meanings for predicates (boolean functions) P and Q that satisfy the condition in that subquestion.

PART (A) [2 MARKS]

Devise an example of set X and predicates P and Q that makes $S1$ true and $S2$ false.

PART (B) [2 MARKS]

Devise an example of set X and predicates P and Q that makes $S1$ false and $S2$ true.

PART (C) [2 MARKS]

Devise an example of set X and predicates P and Q that makes $S1$ true and $S3$ false.

PART (D) [2 MARKS]

Devise an example of set X and predicates P and Q that makes $S1$ false and $S3$ true.

PART (E) [2 MARKS]

Devise an example of set X and predicates P and Q that makes $S2$ true and $S3$ false.

PART (F) [2 MARKS]

Devise an example of set X and predicates P and Q that makes $S2$ false and $S3$ true.

QUESTION 2. [10 MARKS]

Consider the definition of $U(n)$ below:

$U(n) \Leftrightarrow n$ has remainder 2 when divided by 3. In other words $\exists i \in \mathbb{N}, n = 3i + 2$.

Use the given definition, and the proof structure from this course, to PROVE:

$$\forall n \in \mathbb{N}, U(n) \Rightarrow U(n^3)$$

HINT: You may find it helpful to recall the binomial expansion, for any natural numbers a and b ,

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3.$$

QUESTION 3. [10 MARKS]

In this question \mathbb{N}^+ denotes the positive (greater than 0) natural numbers, and $(h, i, j, k) \in \mathbb{N}^+$ denotes a quadruple of positive natural numbers. Use the proof structure from this course to prove the following statement:

$$\forall (h, i, j, k) \in \mathbb{N}^+, \frac{h}{i} < \frac{j}{k} \Rightarrow \frac{h}{i} < \frac{h+j}{i+k}.$$

QUESTION 4. [10 MARKS]

In this question \mathbb{Z} denotes the integers and \mathbb{R} denotes the real numbers. Consider the following definition of $\lfloor x \rfloor$:

$$y = \lfloor x \rfloor \Leftrightarrow y \in \mathbb{Z} \wedge y \leq x \wedge (\forall z \in \mathbb{Z}, z \leq x \Rightarrow z \leq y).$$

Use the given definition, and the proof structure from this course, to prove:

$$\forall x \in \mathbb{R}, \forall y \in \mathbb{R}, x > y \Rightarrow \lfloor x \rfloor \geq \lfloor y \rfloor.$$

QUESTION 5. [10 MARKS]

In this question \mathbb{Z} denotes the integers, \mathbb{R} denotes the real numbers, and \mathbb{R}^+ denotes the real numbers that are greater than zero. Consider the following definition of $\lfloor x \rfloor$:

$$y = \lfloor x \rfloor \Leftrightarrow y \in \mathbb{Z} \wedge y \leq x \wedge (\forall z \in \mathbb{Z}, z \leq x \Rightarrow z \leq y).$$

Use the given definition and the proof structure from this course to DISPROVE:

$$\forall x \in \mathbb{R}, \forall \varepsilon \in \mathbb{R}^+, \exists \delta \in \mathbb{R}^+, \forall w \in \mathbb{R}, |x - w| < \delta \Rightarrow |\lfloor x \rfloor - \lfloor w \rfloor| < \varepsilon$$

QUESTION 6. [10 MARKS]

In this question \mathbb{Z} denotes the integers, \mathbb{R} denotes the real numbers, and \mathbb{R}^+ denotes the real numbers that are greater than zero. Consider the following definition of $\lfloor x \rfloor$:

$$y = \lfloor x \rfloor \Leftrightarrow y \in \mathbb{Z} \wedge y \leq x \wedge (\forall z \in \mathbb{Z}, z \leq x \Rightarrow z \leq y).$$

Use the given definition and the proof structure from this course to PROVE:

$$\exists x \in \mathbb{R}, \forall \epsilon \in \mathbb{R}^+, \exists \delta \in \mathbb{R}^+, \forall w \in \mathbb{R}, |x - w| < \delta \Rightarrow |\lfloor x \rfloor - \lfloor w \rfloor| < \epsilon$$

QUESTION 7. [10 MARKS]

Consider the definition of $\mathcal{O}(g)$ below:

$$\mathcal{O}(g) = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0} : \exists c \in \mathbb{R}^+, \exists B \in \mathbb{N}, \forall n \in \mathbb{N}, n \geq B \Rightarrow f(n) \leq cg(n)\}$$

Define $f(n) = n^3 - 3n + 1$ and $g(n) = 5n^2 + 7n + 9$. Use the proof structure from this course to PROVE $f \notin \mathcal{O}(g)$. You may NOT use the results and techniques of limits from calculus.

QUESTION 8. [10 MARKS]

Consider the definition of $\mathcal{O}(g)$ below:

$$\mathcal{O}(g) = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0} : \exists c \in \mathbb{R}^+, \exists B \in \mathbb{N}, \forall n \in \mathbb{N}, n \geq B \Rightarrow f(n) \leq cg(n)\}$$

Define $f(n) = 3n$ and $g(n) = \ln(n)$. Use the proof structure from this course to PROVE $f \notin \mathcal{O}(g)$. Unlike the previous question, you MAY use the techniques of limits from calculus, including l'Hôpital's rule. You may find it useful to note that the derivative of $\ln(x)$ is $1/x$.

QUESTION 9. [10 MARKS]

Suppose you have a floating-point system with base $\beta = 3$, exponents from the set $\{-2, \dots, 2\}$, $t = 3$ digits in the significand, a single sign symbol $+$ or $-$, a radix point following the first digit, and the convention that the digit preceding the radix point is non-zero unless we are representing zero itself.

PART (A) [5 MARKS]

Which of the following can be represented exactly in the given system? In each case, explain why or why not.

- (i) $3/4$
- (ii) $4/9$
- (iii) $4/27$
- (iv) -25
- (v) 27

PART (B) [5 MARKS]

Give an example of two numbers that each have representations (possibly inexact) in the given system, but when one is subtracted from the other within the given system, yield a relative error of more than 100%. Explain.

This page left intentionally (nearly) blank, to accommodate work that doesn't fit elsewhere.

This page left intentionally (nearly) blank, to accommodate work that doesn't fit elsewhere.

This page left intentionally (nearly) blank, to accommodate work that doesn't fit elsewhere.

This page left intentionally (nearly) blank, to accommodate work that doesn't fit elsewhere.

Total Marks = 92

Student #: _____

Page 14 of 14

END OF EXAM