

Lecture 4

Analysis of Optimal Trajectories, Transition Dynamics in Growth Model

Macroeconomics EC417

Jane Olmstead-Rumsey

London School of Economics, Fall 2022

Plan of Lecture

1. Linearization around steady state, speed of convergence, slope of saddle path
2. Some transition experiments in the growth model

Linearization around Steady State

- Two questions:
 - can we say more than “there exists a unique steady state and the economy converges to it”? Speed of convergence?
 - How analyze stability if two- or N -dimensional state x (so that cannot draw phase diagram)?
- Can answer these questions by analyzing local dynamics close to steady state

Linearization around Steady State

- Let $y \in \mathbb{R}^n$ and the function $m : \mathbb{R}^n \rightarrow \mathbb{R}^n$ define a dynamical system:

$$\dot{y}(t) = m(y(t)) \text{ for } t \geq 0,$$

- Let y^* be a steady state, i.e. $0 = m(y^*)$

Linearization around Steady State

- Let $y \in \mathbb{R}^n$ and the function $m : \mathbb{R}^n \rightarrow \mathbb{R}^n$ define a dynamical system:

$$\dot{y}(t) = m(y(t)) \text{ for } t \geq 0,$$

- Let y^* be a steady state, i.e. $0 = m(y^*)$
- Consider a first order approximation of m around y^* :

$$\dot{y} \approx m(y^*) + m'(y^*)(y - y^*)$$

where $m'(y^*)$ is the $n \times n$ Jacobian of m evaluated at y^* , i.e. the matrix with entries $\partial m_i(y^*)/\partial y_j$

Linearization around Steady State

- Let $y \in \mathbb{R}^n$ and the function $m : \mathbb{R}^n \rightarrow \mathbb{R}^n$ define a dynamical system:

$$\dot{y}(t) = m(y(t)) \text{ for } t \geq 0,$$

- Let y^* be a steady state, i.e. $0 = m(y^*)$
- Consider a first order approximation of m around y^* :

$$\dot{y} \approx m(y^*) + m'(y^*)(y - y^*)$$

where $m'(y^*)$ is the $n \times n$ Jacobian of m evaluated at y^* , i.e. the matrix with entries $\partial m_i(y^*)/\partial y_j$

- Equivalently

$$\dot{\hat{y}} \approx A\hat{y}, \quad \hat{y} = y - y^*, \quad A = m'(y^*)$$

- Idea is then to analyze this linear differential equation.
- Analysis is valid globally (i.e. for all \mathbb{R}^n) if the system is indeed linear.
- Alternatively it is valid in a neighborhood of the steady state.

Linearized Growth Model

- Recall system of two ODEs

$$\begin{aligned}\dot{c} &= \frac{1}{\sigma}(f'(k) - \rho - \delta)c \\ \dot{k} &= f(k) - \delta k - c\end{aligned}\tag{ODE''}$$

- Let $y = (c, k)$ and do analysis on previous slide

$$\hat{y} \approx A\hat{y}, \quad A = \begin{bmatrix} \partial\dot{c}/\partial c & \partial\dot{c}/\partial k \\ \partial\dot{k}/\partial c & \partial\dot{k}/\partial k \end{bmatrix}, \quad \hat{y} = \begin{bmatrix} c - c^* \\ k - k^* \end{bmatrix}$$

where the partial derivatives are evaluated at (c^*, k^*)

- Have

$$A = \begin{bmatrix} \partial\dot{c}/\partial c & \partial\dot{c}/\partial k \\ \partial\dot{k}/\partial c & \partial\dot{k}/\partial k \end{bmatrix} = \begin{bmatrix} 0 & \frac{1}{\sigma}f''(k^*)c^* \\ -1 & \rho \end{bmatrix}$$

where we used $\partial\dot{k}/\partial k = f'(k^*) - \delta = \rho$

Properties of Linear Systems

- **Theorem** (see e.g. Acemoglu, Theorem 7.18) Consider the following linear differential equation system

$$\dot{\hat{y}}(t) = A\hat{y}(t), \quad \hat{y} = y - y^* \quad (*)$$

with initial value $\hat{y}(0)$, and where A is an $n \times n$ matrix. Suppose that $\ell \leq n$ of the eigenvalues of A have negative real parts. Then, there exists an ℓ -dimensional subspace L of \mathbb{R}^n such that starting from any $\hat{y}(0) \in L$, the differential equation $(*)$ has a unique solution with $\hat{y}(t) \rightarrow 0$.

- **Proof:** next slide
- **Interpretation:** important thing is to compare number of negative eigenvalues ℓ and number of pre-determined state variables m
 - if $\ell = m$ (standard case): “saddle-path stable”, unique optimal trajectory. Neg. eigenvalues govern speed of convergence.
 - if $\ell < m$: unstable, $y(t)$ does not converge to steady state.
 - if $\ell > m$: multiple optimal trajectories (“indeterminacy”)

Proof of Theorem

- First step is to solve (*). Also see http://en.wikipedia.org/wiki/Matrix_differential_equation
- Denote the eigenvalues of A by $\lambda_1, \dots, \lambda_n$ and the corresponding eigenvectors by v_1, \dots, v_n .
- Diagonalizing the matrix A we obtain:

$$A = P\Lambda P^{-1}$$

- Λ is diagonal matrix with eigenvalues of A , possibly complex, on its diagonal.
- Matrix P contains the eigenvectors of A , i.e. $P = (v_1, \dots, v_n)$ (to see this write $AP = P\Lambda$ or $Av_i = \lambda_i v_i$) and is invertible
 - ignores some technicalities discussed in ch. 6 of SLP book
- Write system as

$$\begin{aligned} P^{-1}\dot{\hat{y}}(t) &= \Lambda P^{-1}\hat{y}(t) \\ \Leftrightarrow \dot{z}(t) &= \Lambda z(t), \quad z(t) = P^{-1}\hat{y}(t) \end{aligned}$$

Proof of Theorem

- Since Λ is diagonal $\dot{z}_i = \lambda_i z_i(t)$, $i = 1, \dots, n$, i.e. it can be solved element by element

$$z_i(t) = a_i e^{\lambda_i t}$$

where d_i are constants of integration

- We have that $\hat{y}(t) = Pz(t)$. Using $P = (v_1, \dots, v_n)$ we have

$$\hat{y}(t) = \sum_{i=1}^n d_i e^{\lambda_i t} v_i \quad (**)$$

- For now, assume all λ_i are real
- Let λ_i be such that for $i = 1, 2, \dots, \ell$ we have $\lambda_i < 0$ and for $i = \ell + 1, \ell + 2, \dots, n$ we have $\lambda_i \geq 0$. That is, eigenvalues of A are ordered so that first ℓ are negative.

Proof of Theorem

- Q: when does $\hat{y}(t) \rightarrow 0$? A: initial condition needs to satisfy

$$\hat{y}(0) = \sum_{i=1}^n d_i v_i, \quad d_i = 0, i = \ell + 1, \ell + 2, \dots, n$$

- That is, $\hat{y}(t) \rightarrow 0$ only if $\hat{y}(0)$ lies in particular subspace of \mathbb{R}^n .
Dimension of subspace = # of negative eigenvalues ℓ .
- Exercise: how extend to case where λ_i can be complex?

Linearized Growth Model

- Recall

$$\dot{\hat{y}} \approx Ay, \quad A = \begin{bmatrix} 0 & \frac{1}{\sigma} f''(k^*)c^* \\ -1 & \rho \end{bmatrix}, \quad \hat{y} = \begin{bmatrix} c - c^* \\ k - k^* \end{bmatrix}$$

- Let's look at eigenvalues of A . These satisfy

$$0 = \det(A - \lambda I) = -\lambda(\rho - \lambda) + \frac{1}{\sigma} f''(k^*)c^*$$

$$0 = \lambda^2 - \rho\lambda + \frac{1}{\sigma} f''(k^*)c^*$$

Linearized Growth Model

- Recall

$$\dot{\hat{y}} \approx A\hat{y}, \quad A = \begin{bmatrix} 0 & \frac{1}{\sigma}f''(k^*)c^* \\ -1 & \rho \end{bmatrix}, \quad \hat{y} = \begin{bmatrix} c - c^* \\ k - k^* \end{bmatrix}$$

- Let's look at eigenvalues of A . These satisfy

$$0 = \det(A - \lambda I) = -\lambda(\rho - \lambda) + \frac{1}{\sigma}f''(k^*)c^*$$

$$0 = \lambda^2 - \rho\lambda + \frac{1}{\sigma}f''(k^*)c^*$$

- This is a simple quadratic with two solutions (“roots”)

$$\lambda_{1/2} = \frac{\rho \pm \sqrt{\rho^2 - 4\frac{1}{\sigma}f''(k^*)c^*}}{2}$$

- $f''(k^*) < 0$ so both eigenvalues are real, and $\lambda_1 < 0 < \lambda_2$
- Have one pre-determined state variable.
- Theorem says: $\ell = m = 1 \Rightarrow$ saddle-path stable

Linearized Growth Model

- What does this tell us about the time path of capital $k(t)$?
- From (**), solution to matrix differential equation for growth model is

$$\begin{bmatrix} \hat{y}_1(t) \\ \hat{y}_2(t) \end{bmatrix} \approx d_1 e^{\lambda_1 t} \begin{bmatrix} v_{11} \\ v_{12} \end{bmatrix} + d_2 e^{\lambda_2 t} \begin{bmatrix} v_{21} \\ v_{22} \end{bmatrix}$$

where $\hat{y}_1 = c(t) - c^*$, $\hat{y}_2 = k(t) - k^*$ and v_{i1}, v_{i2} denote elements of v_i

- $\lambda_2 > 0 \Rightarrow$ need $d_2 = 0$

$$c(t) - c^* \approx d_1 e^{\lambda_1 t} v_{11}, \quad k(t) - k^* \approx d_1 e^{\lambda_1 t} v_{12}$$

- Have initial condition for $k(0) = k_0 \Rightarrow d_1 v_{12} = k_0 - k^*$

$$c(t) - c^* \approx \frac{v_{11}}{v_{12}} e^{\lambda_1 t} (k_0 - k^*) \quad (1)$$

$$k(t) - k^* \approx e^{\lambda_1 t} (k_0 - k^*) \quad (2)$$

- From (2), know approximate time path for $k(t)$
- (1) pins down initial consumption (v_{11} and v_{12} are known)

Linearization: Speed of Convergence

- Negative eigenvalue λ_1 governs speed of convergence

$$k(t) - k^* \approx e^{-|\lambda_1|t}(k_0 - k^*)$$

- Half-life for convergence to steady state

$$k(t_{1/2}) - k^* = \frac{1}{2}(k_0 - k^*) \quad \Rightarrow \quad t_{1/2} = \frac{\ln(2)}{|\lambda_1|}$$

Linearization: Speed of Convergence

- Negative eigenvalue λ_1 governs speed of convergence

$$k(t) - k^* \approx e^{-|\lambda_1|t}(k_0 - k^*)$$

- Half-life for convergence to steady state

$$k(t_{1/2}) - k^* = \frac{1}{2}(k_0 - k^*) \Rightarrow t_{1/2} = \frac{\ln(2)}{|\lambda_1|}$$

- Formula from previous slide:

$$\lambda_1 = \frac{\rho - \sqrt{\rho^2 - 4\frac{1}{\sigma}f''(k^*)c^*}}{2}$$

- Convergence fast ($|\lambda_1|$ large) if
 - high f'' (strongly diminishing returns)
 - low σ (high “intertemp. elas. of substit.”)
 - low ρ (more patient)
- Later: for reasonable parameter values, neoclassical growth model features very fast convergence, e.g. about $t_{1/2} = 5$ years.

Linearization: Speed of Convergence

- Insights also go through with general utility function $u(c)$
- can show: with general utility function $u(c)$, formula generalizes to

$$\lambda_1 = \frac{\rho \pm \sqrt{\rho^2 - 4 \frac{f''(k^*)c^*}{\sigma(c^*)}}}{2}$$

where

$$\sigma(c) \equiv -\frac{u''(c)c}{u'(c)}$$

- check: $u(c) = \frac{c^{1-\sigma}}{1-\sigma} \Rightarrow \sigma(c) = \sigma$

Slope of Saddle Path

- Recall conditions for optimum:

$$\begin{aligned}\frac{\dot{c}}{c} &= \frac{1}{\sigma}(f'(k) - \rho - \delta) \\ \dot{k} &= f(k) - \delta k - c\end{aligned}\tag{ODE''}$$

with $k(0) = k_0$ and $\lim_{T \rightarrow \infty} e^{-\rho T} c(T)^{-\sigma} k(T) = 0$.

- **Saddle path** $c(k)$ defines optimal consumption for each k
 - a.k.a. consumption policy function
- For many questions, useful to know **slope of saddle path**

Slope of Saddle Path

- Slope of saddle path satisfies

$$c'(k) = \frac{dc}{dk} = \frac{dc/dt}{dk/dt}$$
$$c'(k) = \frac{\frac{1}{\sigma}(f'(k) - \rho - \delta)c(k)}{f(k) - \delta k - c(k)} \quad (*)$$

- Digression: $(*)$ is a non-linear ODE in $c(k)$ that can be solved numerically
 - once solved, know entire dynamics $\dot{k} = f(k) - \delta k - c(k)$
 - alternative to shooting algorithm, no issues with transversality
 - $(*)$ can also be derived from continuous-time Bellman equation (HJB equation)

Slope of Saddle Path

- Now consider slope of saddle path **at steady state** $c'(k^*)$

$$\begin{aligned}c'(k^*) &= \lim_{k \rightarrow k^*} c'(k) = \lim_{k \rightarrow k^*} \frac{\frac{1}{\sigma}(f'(k) - \rho - \delta)c(k)}{f(k) - \delta k - c(k)} = \\&= \frac{\frac{1}{\sigma}f''(k^*)c^*}{f'(k^*) - \delta - c'(k^*)} = \frac{\frac{1}{\sigma}f''(k^*)c^*}{\rho - c'(k^*)}\end{aligned}$$

where the third equality follows from L'Hopital's rule

(http://en.wikipedia.org/wiki/L'H%C3%B4pital's_rule)

- Rearranging, we see that $\lambda = c'(k^*)$ satisfies

$$-\lambda(\rho - \lambda) + \frac{1}{\sigma}f''(k^*)c^* = 0$$

- Same quadratic as before. Two solutions (“roots”)

$$\lambda_{1/2} = \frac{\rho \pm \sqrt{\rho^2 - 4\frac{1}{\sigma}f''(k^*)c^*}}{2}, \quad \lambda_1 < 0 < \lambda_2$$

Slope of Saddle Path

- Know $c'(k^*) > 0 \Rightarrow$ slope of saddle path = positive root

$$c'(k^*) = \frac{\rho + \sqrt{\rho^2 - 4\frac{1}{\sigma}f''(k^*)c^*}}{2}$$

Slope of Saddle Path

- Know $c'(k^*) > 0 \Rightarrow$ slope of saddle path = positive root

$$c'(k^*) = \frac{\rho + \sqrt{\rho^2 - 4\frac{1}{\sigma}f''(k^*)c^*}}{2}$$

- In growth model
 - negative eigenvalue of linearized system = speed of convergence
 - positive eigenvalue of linearized system = slope of saddle path
 - seems to be a coincidence, same not true in more general models. Instead slope of saddle path related to eigenvectors.

Linearization and Perturbation: Relation

- Popular method in economics: perturbation methods
- Some useful references:
 - Judd (1996) “Approximation, perturbation, and projection methods in economic analysis”, Judd’s (1998) book
 - Schmitt-Grohe and Uribe (2004)
 - Fernandez-Villaverde lecture notes
http://economics.sas.upenn.edu/~jesusfv/Chapter_2_Perturbation.pdf
 - Original references from math literature: Fleming (1971), Fleming and Souganidis (1986)
- **Takeaway:** linearization around steady state is particular application of first-order perturbation method
- Why perturbation? Perturbation methods are more general and there are more powerful theorems

Linearization and Perturbation: Relation

- Recall dynamical system from beginning of lecture

$$\dot{y}(t) = m(y(t)), \quad y(0) = y_0 \quad (*)$$

- In general, to apply perturbation method need:
 - some known solution of equation, call it $y^0(t)$
 - to express equation as a perturbed version of known solution in terms of **scalar** “perturbation parameter”, call it ε
- Application to our system (*):
 - know solution if initial condition is steady state, $y_0 = y^*$
 - view (*) as

$$\dot{y}(t, \varepsilon) = m(y(t, \varepsilon)), \quad y(0, \varepsilon) = y^* + \varepsilon \hat{y}_0 \quad (**)$$

where $\varepsilon \hat{y}_0$ is initial deviation from steady state

- Key idea of perturbation: look for solution of (**) of form

$$y(t, \varepsilon) = y^* + \varepsilon y_1(t) + \varepsilon^2 y_2(t) + \dots$$

- First-order perturbation: $y(t, \varepsilon) \approx y^* + \varepsilon y_1(t)$

Linearization and Perturbation: Relation

- Let's implement first-order perturbation: look for solution of (**) of form

$$y(t, \varepsilon) \approx y^* + \varepsilon y_1(t)$$

- Taylor's theorem: set

$$y_1(t) = \frac{\partial y(t, 0)}{\partial \varepsilon}$$

- Find by differentiating (**) with respect to ε

$$\begin{aligned} \frac{\partial}{\partial \varepsilon} \dot{y}(t, 0) &= m'(y(t, 0)) \frac{\partial y(t, 0)}{\partial \varepsilon} \\ \text{i.e. } \dot{y}_1(t) &= m'(y^*) y_1(t) \end{aligned}$$

- Recall linearized system from beginning of lecture

$$\dot{\hat{y}} = A \hat{y}, \quad A = m'(y^*)$$

- Hence $y_1(t)$ solves same equation as $\hat{y}(t)$. Linearization is 1st-order perturbation with $\varepsilon = 1$ so that $y(t) = y^* + y_1(t)$

Linearization: Discrete Time

- Similar results apply for discrete-time optimal control problems
- **Main difference:** it's about whether eigenvalues are < 1 rather than < 0 .
- See Stokey-Lucas-Prescott chapter 6.
- Let $y \in \mathbb{R}^n$ and the function $m : \mathbb{R}^n \rightarrow \mathbb{R}^n$ define a dynamical system:

$$y_{t+1} = m(y_t)$$

- Let y^* be a steady state, i.e. $y^* = m(y^*)$
- Consider a first order approximation of m around y^* :

$$\begin{aligned} y_{t+1} &\approx m(y^*) + m'(y^*)(y_t - y^*) \\ \hat{y}_{t+1} &\approx A\hat{y}_t, \quad \hat{y}_t = y_t - y^*, \quad A = m'(y^*), \end{aligned}$$

Linearization: Discrete Time

- **Theorem** Consider the following linear difference equation system

$$\hat{y}_{t+1} = A\hat{y}_t, \quad \hat{y}_t = y_t - y^* \quad (*)$$

with initial value $\hat{y}(0)$, and where A is an $n \times n$ matrix. Suppose that $\ell \leq n$ of the eigenvalues of A have real parts that are less than one. Then, there exists an ℓ -dimensional subspace L of \mathbb{R}^n such that starting from any $\hat{y}_0 \in L$, the difference equation $(*)$ has a unique solution with $\hat{y}_t \rightarrow 0$.

- Proof is exact analogue
- **My advice:** if you want to linearize a model/do stability analysis, do it in continuous time
 - always works out more nicely
 - but be my guest and linearize growth model in discrete time

Transition Experiments

- Consider growth model with utility and production functions

$$u(c) = \frac{c^{1-\sigma}}{1-\sigma}, \quad f(k) = \varepsilon k^\alpha$$

- Consider following thought experiment
 - up until $t = 0$, economy in steady state
 - at $t = 0$, ε increases permanently to $\varepsilon' > \varepsilon$
- Question: what can we say about time paths of $k(t)$, $i(t)$ and particularly $c(t)$ as model converges to new steady state?
 - consumption increases in long-run, but what about short-run?

Steady State Effects

- For given ε , steady state capital and consumption are

$$k^* = \left(\frac{\alpha \varepsilon}{\rho + \delta} \right)^{\frac{1}{1-\alpha}}, \quad c^* = \varepsilon (k^*)^\alpha - \delta k^*$$


- So both k^* and c^* increase. Note also that

$$\frac{c^*}{k^*} = \varepsilon (k^*)^{\alpha-1} - \delta = \frac{\rho + \delta}{\alpha} - \delta$$


which is independent of ε

- But what about transition?
 - see phase diagrams on next slides

Case 1: slope of new saddle path $< c^*/k^*$



Case 2: slope of new saddle path $> c^*/k^*$



Intuition: income vs. substitution effect

- Why can consumption decrease on impact?
- Offsetting income and substitution effects
 - income effect: $\varepsilon \uparrow \Rightarrow$ wealthier (PDV of earnings higher)
 \Rightarrow eat more
 - substitution effect: $\varepsilon \uparrow \Rightarrow$ MPK \uparrow = return to saving \uparrow
 \Rightarrow eat less
- In general, overall effect is ambiguous
- “Income vs. substitution effect” is always the right answer!

When does $c(t)$ decrease on impact?

- Have formula for slope of saddle path

$$c'(k^*) = \frac{\rho + \sqrt{\rho^2 - 4\frac{1}{\sigma}f''(k^*)c^*}}{2}$$

\Rightarrow can say more

- Consider special case $\delta = 0$ (makes algebra easier). Have

$$\alpha\varepsilon'(k^*)^{\alpha-1} = \rho, \quad \frac{c^*}{k^*} = \frac{\rho}{\alpha}$$

$$\begin{aligned} c'(k^*) &= \frac{\rho + \sqrt{\rho^2 + 4\frac{1-\alpha}{\sigma}\alpha\varepsilon'(k^*)^{\alpha-2}c^*}}{2} \\ &= \frac{\rho + \sqrt{\rho^2 + 4\frac{1-\alpha}{\sigma}\alpha\varepsilon'(k^*)^{\alpha-1}\frac{c^*}{k^*}}}{2} = \frac{\rho}{2} \left(1 + \sqrt{1 + \frac{4}{\sigma} \frac{1-\alpha}{\alpha}} \right) \end{aligned}$$

When does $c(t)$ decrease on impact?

- So question is when

$$c'(k^*) = \frac{\rho}{2} \left(1 + \sqrt{1 + \frac{4}{\sigma} \frac{1-\alpha}{\alpha}} \right) > \frac{\rho}{\alpha} = \frac{c^*}{k^*}$$
$$\sqrt{\alpha^2 + \frac{4}{\sigma}(1-\alpha)\alpha} > 2 - \alpha$$
$$\alpha > \sigma$$

- Summary:
 - $\sigma > \alpha$: income effect dominates $\Rightarrow c(t)$ increases
 - $\sigma < \alpha$: subst. effect dominates $\Rightarrow c(t)$ decreases
 - $\sigma = \alpha$: income and subst. effects cancel $\Rightarrow c(t)$ constant
- Exercise: what is the cutoff in general case $\delta > 0$

Transition Experiments

- Exercise: analogous experiments for other parameter values
 - increase in ρ
 - increase in δ