

ACTIVE SEGMENTATION HELP

DIMITER PRODANOV^{1,2}

¹*NERF, IMEC, Kapeldreef 75, 3001 Leuven, Belgium;* ²*ITSDP, IICT, Bulgarian Academy of Sciences, Sofia, Bulgaria*

CONTENTS

1. Gaussian Scale Spaces	1
2. α -Scale Spaces	2
3. Weak differentiation	2
4. Differential Invariants	3
5. Orthogonal Laplacian Decomposition (LOD)	3
6. Zero-order Geometric Features	4
6.1. Gauss2D – Gaussian smoothing	4
6.2. FFT Gaussian smoothing	4
7. First-order Geometric Features	4
7.1. Gradient	4
7.2. Gaussian Structure – Structure tensor	4
8. Second-order Geometric Features	5
8.1. LoG – Laplacian of Gaussian	5
8.2. FFT LoG	5
8.3. ALoG – Anisotropic decomposition of LoG	5
8.4. Hessian	5
8.5. Curvature 2D	6
8.6. Gaussian curvature	6
8.7. Curvature 3D	6
9. Higher order Geometric Features	7
9.1. Gaussian Jet	7
9.2. Bi-Laplacian of Gaussian (BoG)	7
References	7

1. GAUSSIAN SCALE SPACES

In the digital domain, smoothing leads to loss of resolution and, therefore, of some information. The axiomatic linear scale space theory was formulated in series of works by Witkin and Koenderink [10, 4]. In its original version, the theory depends on several properties of the Gaussian filters as solutions of the diffusion

equation in the scale-space generated by the image. That is, the generic smoothing kernel G is identified with a radially-symmetric Gaussian kernel of scale $s = \sigma^2 \in \mathbb{R}$

$$G(r) = \frac{e^{-r^2/2s}}{2\pi s} = \frac{e^{-(x^2+y^2)/2s}}{2\pi s}$$

The Gaussian kernels provide several advantages: (i) they are rotationally invariant (ii) they do not produce artificial extrema in the resulting image (iii) successive convolutions with different kernels can be combined. Mathematically, this imposes a very useful semi-group structure, equivalent to the heat/diffusion equation. In this sense, the image structures diffuse or "melt-down", so that the rate of this diffusion indicates the "robustness" of the structure.

2. α -SCALE SPACES

However, the information loss can be limited if one uses multiple smoothing scales Pauwels et al. [6] and later Duits et al. [2] introduced the α -scale spaces for image processing. The basis of the approach is a generalization the heat equation. The resulting convolution kernels can be described best by the tools of fractional calculus. The kernel evolution is governed by two parameters – the scale s and the order of differentiation α . The approach leads to the fractional heat problem:

$$\begin{aligned} u(0, \mathbf{x}) &= I(\mathbf{x}) \\ \partial_s u(s, \mathbf{x}) &= -(-\Delta)^\alpha u(s, \mathbf{x}), \quad 0 \leq \alpha \leq 1 \end{aligned}$$

where the Riesz fractional Laplacian operator is defined in the Fourier domain by:

$$(-\Delta)^\alpha U(\mathbf{k}) := k^{2\alpha} U(\mathbf{k}), \quad k = |\mathbf{k}|,$$

where the k is the modulus of the wave vector \mathbf{k} . Formally, the operator is extended by continuity for $\alpha = 1$ as $(-\Delta)^1 = \Delta \mapsto k^2$, which corresponds to the usual Laplacian; and to identity for $\alpha = 0$, corresponding to invariance in the spatial domain. The Green function of the differential equation is the stretched exponential kernel

$$G(k, s) = e^{-k^{2\alpha} s}$$

in the frequency domain. Direct inversion of the Green function gives the Bessel integral

$$G_\alpha(r, s) = \frac{1}{\pi} \int_0^\infty J_0(kr) e^{-k^{2\alpha} s} k dk$$

involving the Bessel function of first kind $J_0(r)$.

3. WEAK DIFFERENTIATION

Weak differentiation operations in distributional sense can be defined in terms of convolution with the gradient of a smooth kernel function as:

$$\nabla_G F := -F \star \nabla G$$

where the symbol ∇ represents the of the gradient operator for the Euclidean basis (e_1, e_2)

$$\nabla = e_1 \frac{\partial}{\partial x} + e_2 \frac{\partial}{\partial y}$$

Filter	Functionality	Feature order
Gauss2D	Gaussian smoothing	0
Gradient	Gradient amplitude and orientation	1
Gaussian Structure	Structure tensor	1
LoG	Laplacian of Gaussian (LoG)	2
ALoG	Anisotropic decomposition of LoG	2
	Gradient amplitude and orientation	1
Hessian	Eigenvalues of the Hessian	2
	Determinant of the Hessian	2
Curvature 2D	Line curvatures + Hessian determinant	2
Curvature 3D	Mean + Gauss curvature of surfaces	2
BoG	Bi-Laplacian of Gaussian	4
Gaussian Jet	Gaussian Jet of order n	n
LoGN	n-th order PoL	2n
FFT Kernel LoG	Riesz Laplacian	$\alpha \in \mathbb{R}$

TABLE 1. Filters computing geometric features[9]

4. DIFFERENTIAL INVARIANTS

There are several types of geometric features useful for image segmentation. Typical interesting image features are blobs, filaments and corners. The normal Laplacian $\Delta_{\perp G}$ presented below is sensitive to blobs, while its complement – the tangential Laplacian $\Delta_{\parallel G}$ is sensitive to filaments.

Various geometric features computed by the AS/IJ platform are presented in Table 1. Notable differential invariants are the amplitude $|\nabla_G F|$ and the orientation of the gradient, the Hessian determinant, the Hessian eigenvalues, as well as the isophote κ and streamline curvatures μ [3]. Up to a sign convention we have

$$\begin{aligned}
 (1) \quad & \kappa = \nabla_G \cdot \mathbf{n}^0 \\
 (2) \quad & \mu = \nabla_G \cdot \tau, \quad \tau = I_2 \cdot \mathbf{n}^0
 \end{aligned}$$

where I_2 is the pseudoscalar of the Euclidean image plane. From the perspective of scale space theory the study of the differential invariants of the image reduce to the study of the differential invariants of the radially symmetric (generalized) heat kernel.

5. ORTHOGONAL LAPLACIAN DECOMPOSITION (LOD)

The simple representation of the gradient introduced above can be extended to a coordinate independent (!) operator using the tools of Geometric Algebra and Calculus. Readers are directed to [5] for an introductory material on the subject. The Laplace operator can be decomposed in two orthogonal components– on in the direction of the isophote and the other in the direction normal to the isophotes. The Laplacian of scalar function F can be decomposed into a normal component and tangential component, thus breaking the isotropy of the original operator [8]:

$$\nabla^2 F = (\mathbf{n}^0 \cdot \nabla)^2 F + (\mathbf{n}^0 \cdot \nabla F) \nabla \cdot \mathbf{n}^0 = \Delta_{\perp} F + \Delta_{\parallel} F$$

where $\mathbf{n} = \nabla F$ is the gradient, \mathbf{n}^0 is the unit normal vector to the isophote curve $F(x, y) = c$,

$$\mathbf{n}^0 = \mathbf{n}/|\mathbf{n}|$$

and $\mathbf{n}^0 \cdot \nabla$ denotes the directional derivative. Then also using the isophote curvature the components are given by

$$(3) \quad \Delta_{\perp} = (\mathbf{n}^0 \cdot \nabla)^2$$

$$(4) \quad \Delta_{\parallel} = (\nabla \cdot \mathbf{n}^0) (\mathbf{n}^0 \cdot \nabla) = \kappa \mathbf{n}^0 \cdot \nabla$$

This is a coordinate-free definition of Δ_{\perp} and Δ_{\parallel} , which can be specialized to any smooth coordinate system. This comes in contrast to the approach of "gauge coordinates" employed in [3]. Furthermore, if we specialize to weak Gaussian derivatives

$$\Delta_{\parallel G} F = \kappa |A|, \quad |A| = \sqrt{G_x^2 + G_y^2}$$

6. ZERO-ORDER GEOMETRIC FEATURES

6.1. Gauss2D – Gaussian smoothing. The filter convolves an image with the Gaussian kernel in the spatial domain.

$$G(x, y) \star F(x, y)$$

6.2. FFT Gaussian smoothing. The filter convolves an image with the Riesz/Gaussian kernel in the Fourier domain.

$$\mathcal{F}^{-1} e^{-k^{2\alpha}} F(k)$$

7. FIRST-ORDER GEOMETRIC FEATURES

Gradient amplitude	$A = \sqrt{G_x^2 + G_y^2}$
Gradient orientation	$\sin \phi = G_y / \sqrt{G_x^2 + G_y^2}$
	$\cos \phi = G_x / \sqrt{G_x^2 + G_y^2}$

TABLE 2. First-order differential invariants

7.1. Gradient. The filter computes the gradient amplitude and orientation (sine and cosine). The full output also outputs the elements of the gradient G_x and G_y .

7.2. Gaussian Structure – Structure tensor. The structure tensor (ST) is an abstract extension of the gradient. The tensor encodes the predominant directions of the gradient in a specified neighborhood of a point, and the degree to which those directions are coherent as a function of scale. Suppose that we have a scale-space representation of the gradient ∇_G . Then the structure tensor is the smoothed tensor product of the smoothed gradient vector[1]:

$$S_r(F) := G_r \star \{\nabla_G \cdot \nabla_G^T\}$$

From this expression it is apparent that the operator introduces smoothing on two scales. However, because of its quadratic characters the scales do not compose. $S_r(I)$ can be represented by a 2x2 matrix.

Laplacian	$\Delta_G = \text{Tr } \mathbb{H} = G_{xx} + G_{yy}$
determinant of the Hessian	$\det \mathbb{H}_G = G_{xx}G_{yy} - G_{xy}^2$

TABLE 3. Second-order differential invariants

8. SECOND-ORDER GEOMETRIC FEATURES

8.1. LoG – Laplacian of Gaussian. The Laplacian operator can be thought of as the square of the gradient ∇ . The analogy can be made precise using the tools of the Clifford algebra and Geometric Calculus: $\Delta = \nabla^2$. In Cartesian coordinates, the Laplacian has the representation

$$\Delta_G = G_{xx} + G_{yy}$$

The filter convolves an image with the Laplacian of Gaussian.

8.2. FFT LoG. The plugin convolves an image with the Riesz Laplacian kernel of order α .

8.3. ALoG – Anisotropic decomposition of LoG. The theory of the anisotropic decomposition of LoG (ALoG) is given in Sec. 5. In Cartesian coordinates, the LoG is represented by

$$(5) \quad (G_x^2 + G_y^2) \Delta_{\perp G} = (G_x^2) G_{xx} + (2G_x G_y) G_{xy} + (G_y^2) G_{yy}$$

$$(6) \quad (G_x^2 + G_y^2) \Delta_{\parallel G} = (G_x^2) G_{xx} - (2G_x G_y) G_{xy} + (G_y^2) G_{yy}$$

The plugin computes the anisotropic decomposition. The full output option also outputs the components of the gradient and the Hessian.

8.4. Hessian. The weak Hessian tensor with respect to the (Gaussian) kernel G is defined as the tensor product

$$\mathbb{H}_G(F) := \nabla_G \otimes \nabla_G F$$

For smooth signals, the Hessian is symmetric and can be regarded as a metric tensor. In Cartesian coordinates, the Hessian can be specialized to the usual formula

$$\mathbb{H}_G(F) = \begin{pmatrix} G_{xx} & G_{xy} \\ G_{xy} & G_{yy} \end{pmatrix} \star F$$

where subscripts denote differentiation by coordinate variables. The Hessian generates several differential invariants. These are the trace, determinant and the eigenvalues.

For the trace holds

$$\text{Tr } \mathbb{H}_G(F) = \nabla_G^2 F = \Delta_G F$$

where Δ_G denotes the weak Laplacian operator.

The determinant is

$$\det \mathbb{H}_G(F) = G_{xx}G_{yy} - G_{xy}^2$$

The eigenvalues $\lambda_{1,2}$ can be determined locally from the equation

$$\det(\mathbb{H}_G(F) - \lambda \mathbb{I}) = 0$$

This gives the quadratic equation

$$\lambda^2 - \text{Tr } \mathbb{H}_G(F) \lambda + \det \mathbb{H}_G(F) = 0$$

In an explicit form the eigenvalues are

$$\lambda_{1,2} = \frac{1}{2} \left((G_{xx} + G_{yy}) \pm \sqrt{(G_{xx} + G_{yy})^2 - 4(G_{xx}G_{yy} - G_{xy}^2)} \right)$$

The plugin computes the amplitude, sine and cosine of the gradient, the Hessian determinant and its 2 eigenvalues. The full output option also outputs the components of the gradient and the Hessian.

8.5. Curvature 2D. The planar image is represented as a set of isophote contours

$$F(x, y) = c$$

The gradient vector n is orthogonal to the isophote contour.

The plugin computes 2 invariants:

- Isophote curvature

$$\kappa = \frac{G_{xx}G_y^2 - 2G_xG_yG_{xy} + G_x^2G_{yy}}{(G_x^2 + G_y^2)^{3/2}}$$

- Streamline curvature

$$\mu = \frac{G_xG_y(G_{yy} - G_{xx}) + G_{xy}(G_x^2 - G_y^2)}{(G_x^2 + G_y^2)^{3/2}}$$

The full output option outputs the components of the gradient and the Hessian.

8.6. Gaussian curvature. There is a second plugin computing the extrinsic linear curvature

$$\nu = \frac{G_xG_{yy} - G_yG_{xx}}{(G_x^2 + G_y^2)^{3/2}}$$

and the determinant of the Hessian. The full output option outputs the components of the gradient and the Hessian.

8.7. Curvature 3D. The planar image is represented as a surface in the three-dimensional Euclidean space \mathbb{E}^3 , where the elevation z represents the signal intensity.

$$z = F(x, y)$$

The plugin computes

- Mean curvature

$$K_m = \frac{1}{2} \frac{(1 + G_x^2)G_{yy} - 2G_xG_yG_{xy} + (1 + G_y^2)G_{xx}}{(1 + G_x^2 + G_y^2)^{3/2}}$$

- Gaussian curvature

$$K = \frac{G_{xx}G_{yy} - G_{xy}^2}{(1 + G_x^2 + G_y^2)^2}$$

Geometrically, the mean curvature is given by the divergence of the unit normal vector in 3D. The full output option outputs the components of the gradient and the Hessian.

9. HIGHER ORDER GEOMETRIC FEATURES

9.1. Gaussian Jet. In the spatial domain, the Gaussian derivatives for the one dimensional case can be computed in closed form as

$$G_n(x) = \frac{\partial^n}{\partial x^n} G(x) = \frac{(-1)^n}{\sqrt{2\pi s^{n+1}}} He_n(x/\sqrt{s}) e^{-\frac{x^2}{2s}}$$

where $He_n(x)$ is the statistician's Hermite polynomial of order n . The sequence of statistician's Hermite polynomials satisfies the recursion

$$He_{n+1}(x) = xHe_n(x) - nHe_{n-1}(x)$$

starting from $He_0(x) = 1$ and $He_1(x) = x$. This allows for efficient simultaneous computation of all derivatives up to an order n in order to populate the the n -jet space. The filter computes all Gaussian derivatives up to order n .

9.2. Bi-Laplacian of Gaussian (BoG). The Laplacian operator can be composed multiple times to give rise to the Power-of-Laplacian (PoL) operator [7]: $\Delta_G^n I$. The plugin computes $\Delta_G^2 I$. This operator enhances high-frequency features of an images given the scale cut-off. This can be seen easily from the frequency response of the LoG filter.

REFERENCES

- [1] T. Brox, J. Weickert, B. Burgeth, and P. Mrazek. Nonlinear structure tensors. Technical report, Universitat des Saarlandes, 2004.
- [2] R. Duits, M. Felsberg, L. Florack, and B. Platel. Alpha-scale spaces on a bounded domain. scale space methods in computer vision, 494- 510. *Springer*, 2003.
- [3] L. M J Florack, B. M ter Haar Romeny, J. J Koenderink, and M. A Viergever. Scale and the differential structure of images. *Image and Vision Computing*, 10:376–388, 1992.
- [4] J. J. Koenderink. The structure of images. *Biological Cybernetics*, 50(5):363–370, aug 1984.
- [5] A. Macdonald. A survey of geometric algebra and geometric calculus. *Advances in Applied Clifford Algebras*, pages 1–39, 2016.
- [6] E. J Pauwels, L. J. Van Gool, P. Fiddelaers, and T. Moons. An extended class of scale-invariant and recursive scale space filters. *Pattern Analysis and Machine Intelligence, IEEE Transactions on*, 17(7):691–701, 1995.
- [7] D. Prodanov, T. Konopczynski, and M. Trojnar. Selected applications of scale spaces in microscopic image analysis. *Cybernetics and Information Technologies*, 15(7):5–12, dec 2015.
- [8] Dimitar Prodanov and Sumit Kumar Vohra. Active segmentation: Differential geometry meets machine learning. In *International Conference on Computer Systems and Technologies 2022*. ACM, jun 2022.
- [9] S. K. Vohra and D. Prodanov. The active segmentation platform for microscopic image classification and segmentation. *Brain Sciences*, 11(12):1645, dec 2021.
- [10] A.P. Witkin. Scale-space filtering. In *Proc. 8th Int. Joint Conf. Artificial Intelligence (IJCAI 83)*, volume 2, 1019-1022, 1983.