Finite Fields and Polynomial Factorization

N-20221002

§1.1 Constructing Finite Fields

The construction and factorization results follow the standard organization in Lidl and Niederreiter, *Introduction to Finite Fields and their Applications*, Chapters 2–4, with small examples calculated explicitly below.[1] The central question is: how do polynomials factor when the coefficients live in a finite field rather than in $\mathbb{Q}$, $\mathbb{R}$, or $\mathbb{C}$?

The question is not only theoretical: it appears in coding theory, cryptography, computational algebra, and the construction of algebraic extensions. The definition of a field comes first because factorization depends heavily on the coefficient domain.

A field F is a set with addition and multiplication such that:

(i) $(F, +)$ is an abelian group with identity 0;

(ii) $F \setminus \{0\}$ is an abelian group under multiplication with identity 1;

(iii) multiplication distributes over addition.

The exclusion of 0 from the multiplicative group is essential. If 0 had a multiplicative inverse, then from $0 \cdot a = 0$ one would get contradictions such as $1 = 0$. So the phrase "every nonzero element has an inverse" is not a technical nuisance; it is what makes division possible.

Examples are

$$\mathbb{Q}, \qquad \mathbb{R}, \qquad \mathbb{C},$$

and, for a prime p,

$$\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}.$$

The requirement that p be prime is crucial. If p is composite, zero divisors appear. For example in $\mathbb{Z}/6\mathbb{Z}$,

$$2 \cdot 3 \equiv 0 \pmod 6,$$

even though neither factor is zero.

A field extension K/F means F is embedded in a larger field K. The point of passing to K is often to make more polynomial equations solvable. For example,

$$x^2 + 1$$

has no root in $\mathbb{R}$, but it splits in $\mathbb{C}$:

$$x^2 + 1 = (x - i)(x + i).$$

If K is finite-dimensional as a vector space over F, its dimension is the degree of the extension:

$$[K : F] = \dim_F K.$$

[1] R. Lidl and H. Niederreiter, *Introduction to Finite Fields and their Applications*, Cambridge University Press, Chapters 2–4.

Finite extensions are tractable because linear algebra applies directly, whereas infinite extensions require substantially heavier tools.

For a prime p, $\mathbb{F}_p$ is the simplest finite field. To construct fields with p^n elements, use irreducible polynomials. If

$$f(x) \in \mathbb{F}_p[x]$$

is irreducible of degree n, then

$$\mathbb{F}_p[x]/(f(x))$$

is a field with p^n elements.

For example, over $\mathbb{F}_2$, the polynomial

$$f(x) = x^3 + x + 1$$

has no root in $\mathbb{F}_2$, hence is irreducible of degree 3. Therefore

$$\mathbb{F}_2[x]/(x^3 + x + 1)$$

is a field with $2^3 = 8$ elements.

In this quotient field, if α denotes the class of x, then

$$\alpha^3 + \alpha + 1 = 0,$$

so

$$\alpha^3 = \alpha + 1$$

in characteristic 2. Every element can be uniquely written as

$$a + b\alpha + c\alpha^2, \qquad a, b, c \in \mathbb{F}_2.$$

§1.2 The first irreducibility test: degrees one and two

In $\mathbb{F}_q[x]$, every nonzero polynomial factors uniquely into irreducible polynomials, up to order and multiplication by nonzero constants. This is analogous to unique prime factorization of integers.

However, a polynomial need not split into linear factors over its base finite field. For example, a polynomial may be irreducible over $\mathbb{F}_p$ but split over an extension field. This is exactly like $x^2 + 1$ over $\mathbb{R}$, except that the extension fields are finite and highly structured.

Consider

$$f(x) = x^4 + x^2 + 2 \in \mathbb{F}_3[x].$$

First test roots:

$$f(0) = 2 \neq 0,$$

$$f(1) = 1 + 1 + 2 = 4 \equiv 1 \pmod 3,$$

$$f(2) = 2^4 + 2^2 + 2 = 16 + 4 + 2 = 22 \equiv 1 \pmod 3.$$

So f has no linear factor. It might still factor as a product of two quadratics. For a degree-four polynomial, root testing is insufficient: a reducible polynomial without roots must factor into two irreducible quadratics. Over $\mathbb{F}_3$, the monic irreducible quadratics are exactly $x^2 + 1$, $x^2 + x + 2$, and $x^2 + 2x + 2$. Dividing f by these yields respective remainders

$$2, \qquad 2x + 2, \qquad x + 2.$$

None vanishes. Since f also has no linear factor, it is irreducible over $\mathbb{F}_3$. This completes the omitted calculation rather than merely asserting that a check exists.

This illustrates the basic workflow: root testing only detects linear factors. For degree 4, one must also check quadratic factors.

A polynomial can factor differently depending on the coefficient field. For example, $x^2 + 1$ is irreducible over $\mathbb{R}$ only if one insists on real linear factors, but over $\mathbb{C}$ it splits. Over finite fields the same phenomenon becomes arithmetic.

For odd p, a quadratic polynomial

$$ax^2 + bx + c, \qquad a \neq 0,$$

has a root (and hence factors) iff its discriminant

$$\Delta = b^2 - 4ac$$

is a square in $\mathbb{F}_p$, including zero. The restriction $p \neq 2$ matters: the quadratic formula divides by $2a$, which is impossible in characteristic 2. In characteristic 2, the polynomial must instead be analyzed using its actual values or an appropriate Artin–Schreier form. Thus irreducibility is controlled by quadratic residues. This links polynomial factorization directly to modular arithmetic.

§1.3 Frobenius is the field's intrinsic symmetry

The standard facts are:

(i) finite fields have order p^n;

(ii) for every prime power p^n, there is a finite field of that order, unique up to isomorphism;

(iii) the nonzero elements $\mathbb{F}_{p^n}^{\times}$ form a cyclic group of order $p^n - 1$;

(iv) the Frobenius map

$$\mathrm{Frob}(a) = a^p$$

is an automorphism of $\mathbb{F}_{p^n}$.

The cyclicity of the multiplicative group is surprisingly powerful. It means every nonzero element is a power of one primitive element. This is why finite field computations often become exponent arithmetic modulo $p^n - 1$.

In characteristic p, the Frobenius map is

$$\varphi(a) = a^p.$$

It is a field homomorphism because

$$(a + b)^p = a^p + b^p$$

in characteristic p. In a finite field, Frobenius is an automorphism. Its fixed field inside $\mathbb{F}_{p^n}$ is $\mathbb{F}_p$.

The polynomial

$$x^{p^n} - x$$

has all elements of $\mathbb{F}_{p^n}$ as roots, and it factors over $\mathbb{F}_p$ as the product of all monic irreducible polynomials whose degrees divide n. This is a central organizing fact for finite-field factorization.

In characteristic p, the Frobenius map

$$F : x \mapsto x^p$$

is a field homomorphism because

$$(x+y)^p = x^p + y^p.$$

In $\mathbb{F}_{p^n}$, it is an automorphism, and

$$F^n = 1.$$

The Galois group is cyclic:

$$\mathrm{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p) = \langle F \rangle \cong \mathbb{Z}/n\mathbb{Z}.$$

This is the cleanest first example of a Galois group: roots are moved by raising to powers of p, and the fixed field of Frobenius is $\mathbb{F}_p$.

An element $\alpha \in \mathbb{F}_{q^n}$ has conjugates

$$\alpha, \quad \alpha^q, \quad \alpha^{q^2}, \quad \dots$$

under Frobenius. Its minimal polynomial over $\mathbb{F}_q$ is

$$\prod_{i=0}^{d-1} (x - \alpha^{q^i}),$$

where d is the size of the Frobenius orbit. Thus irreducible polynomials over finite fields are Frobenius orbits in disguise.

This explains why the degrees of irreducible factors of a polynomial are controlled by Frobenius cycles.

§1.4 Factor Degrees and Möbius Inversion

Every element of $\mathbb{F}_{q^n}$ is a root of

$$x^{q^n} - x.$$

Over $\mathbb{F}_q$, this polynomial is the product of all monic irreducible polynomials whose degrees divide n. If N_d is the number of monic irreducible polynomials of degree d, then

$$q^n = \sum_{d \mid n} d N_d.$$

By Möbius inversion,

$$N_n = \frac{1}{n} \sum_{d \mid n} \mu(d) q^{n/d}.$$

This formula turns factorization into enumeration. The factor d appears because each degree-d irreducible polynomial contributes d roots.

Finite-field factorization algorithms exploit Frobenius. The polynomial

$$x^{q^m} - x$$

has as roots exactly the elements of $\mathbb{F}_{q^m}$. Therefore

$$\gcd(f(x), x^{q^m} - x)$$

extracts from f the product of irreducible factors whose degrees divide m.

This is the idea behind distinct-degree factorization. First analyze repeated factors by checking $\gcd(f, f')$. If $f' = 0$ over a perfect finite field, this signals that f is a polynomial in x^p; the square-free algorithm must take an appropriate p-th root before continuing, rather than dividing by the zero derivative. Then use Frobenius powers to separate factors by degree. Finally use algorithms such as Cantor–Zassenhaus to split equal-degree parts.

Berlekamp's algorithm turns factorization into a nullspace computation. Over $\mathbb{F}_q$, consider the map on the quotient algebra $A = \mathbb{F}_q[x]/(f)$

$$B : g \mapsto g^q - g.$$

Elements in the kernel satisfy $g^q = g$, and this kernel contains information about decomposing f into factors.

Thus the algorithm uses the finite-dimensional $\mathbb{F}_q$-vector space structure of the quotient ring. This is a concrete instance of a general computational theme: algebraic structure becomes linear algebra after passing to a finite quotient.

§1.5 Evaluation Codes from Polynomials

Finite fields are essential in coding theory because code symbols can be added, multiplied, and evaluated polynomially. Reed–Solomon codes evaluate polynomials over $\mathbb{F}_q$ at distinct field points:

$$f \longmapsto (f(a_1), \ldots, f(a_m)).$$

A nonzero polynomial of degree $< k$ has at most $k - 1$ roots. Thus two distinct polynomials of degree $< k$ differ in at least $m - k + 1$ of the m evaluation coordinates, assuming $m \leq q$ distinct evaluation points. This proves the Reed–Solomon minimum-distance bound $d_{\min} \geq m - k + 1$, with equality for suitable polynomial differences; hence a codeword has enough separation to correct up to $\lfloor (m - k)/2 \rfloor$ symbol errors.

Here the same polynomial-root principle used in factorization becomes a robustness principle for information transmission.

The identity

$$\#\mathbb{A}^1(\mathbb{F}_q) = q$$

and the polynomial $x^q - x$ already hint at a larger story: equations over finite fields define finite sets of points, and Frobenius acts on their solutions. In higher-dimensional algebraic geometry, counting points over $\mathbb{F}_{q^n}$ reveals cohomological information.

Thus elementary finite-field polynomial factorization is a first step toward modern arithmetic geometry: finite fields provide a world where algebraic equations, symmetries, and point counts interact exactly.