

Set Problems, Extremal Counting, and Venn-Diagram Reasoning

N-20221207

§1.1 A set construction problem

The first problem defines a finite set

$$A = \{a_1 < a_2 < \cdots < a_k\}$$

of positive integers and builds two derived sets, one from sums and one from differences. The handwritten solution records the important idea: use the union $S \cup T$, and do not try to identify every element exactly. Bound its size from below and above.

The sum set contains at least the ordered chain

$$2a_1, a_1 + a_2, 2a_2, \dots, a_{k-1} + a_k, 2a_k,$$

so

$$|S| \geq 2k - 1.$$

The difference set contains

$$0, a_2 - a_1, a_3 - a_1, \dots, a_k - a_1,$$

so

$$|T| \geq k.$$

Under the condition of the problem, $S \cap T = \emptyset$, hence

$$|S \cup T| = |S| + |T| \geq 3k - 1.$$

On the other hand, all these numbers lie between 0 and $2a_k$, so

$$|S \cup T| \leq 2a_k + 1.$$

Thus

$$3k - 1 \leq 2a_k + 1.$$

If $a_k \leq 2021$, then

$$k \leq \frac{2 \cdot 2021 + 2}{3} = 1348.$$

The construction

$$A = \{674, 675, \dots, 2021\}$$

attains this bound, so the maximum is

$$|A| = 1348.$$

§1.2 What this problem teaches

The note's blue comment says that using $S \cup T$ is a good way to avoid proving a complicated extremal statement directly. This is the key lesson. When a problem asks for the largest possible size of a set, it often helps to build auxiliary sets whose sizes are forced by the structure.

A lower bound comes from a construction. An upper bound comes from counting constraints. A complete solution needs both.

§1.3 A Venn-diagram necessary-and-sufficient problem

Another problem defines set difference

$$A - B = \{x : x \in A, x \notin B\}$$

and asks for the relation among A, B, C when

$$(A - B) \cup (B - A) \subseteq C.$$

The question asks whether this is equivalent to a condition involving

$$A \subseteq (C - B) \cup (B - C)$$

or


$$A \cap B \cap C = \emptyset.$$

The handwritten solution first draws a Venn diagram, then rewrites the pieces algebraically. The symmetric difference

$$(A - B) \cup (B - A)$$

is the part where membership in A and B differs. If this is contained in C , then outside C the sets A and B must agree.

The correct way to solve such problems is to analyze each of the eight Venn regions. The note records a false start and then corrects it, which is valuable: set identities are easy to get wrong if one reasons only verbally.



A Venn diagram is a region bookkeeping device.

Figure 1.1: For three sets one should reason region by region, not by visual guesswork alone.

§1.4 Difference identities

One useful identity appearing in the handwritten reasoning is

$$B - C = B \cap C^c.$$

Therefore

$$B - C - (A \cap B) = B \cap C^c \cap (A \cap B)^c.$$

Using De Morgan,

$$(A \cap B)^c = A^c \cup B^c,$$

so

$$B \cap C^c \cap (A^c \cup B^c) = (B \cap C^c \cap A^c) \cup (B \cap C^c \cap B^c).$$

The second term is empty, hence

$$B - C - (A \cap B) = B \cap A^c \cap C^c.$$

This is exactly the kind of algebraic cleanup that a Venn diagram suggests but does not replace.

§1.5 Coordinate and algebraic exercises

The later pages contain a mixture of algebraic and coordinate exercises. They include problems about quadratic expressions, tangent or chord diagrams, and conditions that must be translated into equations. The handwritten computations repeatedly use the same pattern:

- (i) introduce the unknowns explicitly;
- (ii) translate each condition into an equation or inequality;
- (iii) reduce by substitution or by completing the square;
- (iv) check equality or boundary cases.

Several sketches show lines touching curves or intersecting regions. Rather than reproducing them as raw images, the important content is the method: a picture tells us which algebraic condition to impose, such as tangency, parallelism, or containment.

§1.6 A note on proof writing

The red handwritten reflection says that a purely diagrammatic method is intuitive but can be unreliable. The mature solution uses the diagram to discover the statement, then uses set algebra to prove it. This is an excellent general lesson. Geometry and diagrams are for finding the path; formal manipulations are for making the path safe.

§1.7 Extremal set problems: why constructions matter

In an extremal counting problem, an upper bound is only half the solution. One must also give a construction showing the bound is sharp. The note's sum-set and difference-set arguments follow this pattern: first prove that the derived sets must contain enough elements; then exhibit a set A for which the lower bound is actually attained.

This is a common combinatorial habit. Inequalities show impossibility beyond a threshold. Constructions show that the threshold is real.

§1.8 Venn diagrams versus proofs

Venn diagrams are excellent for discovering set identities, but the proof should eventually be written elementwise. For example,

$$A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$$

is proved by taking an arbitrary x and translating membership:

$$x \in A \setminus (B \cup C) \iff x \in A, x \notin B, x \notin C.$$

This is exactly the condition for membership in the right-hand side. The diagram gives intuition; the element chase gives proof.

§1.9 A wider frame

This note is about extremal and logical reasoning. The extremal problem uses auxiliary sets and cardinality bounds. The Venn problem uses Boolean algebra and region decomposition. Both are early forms of a common method: replace a vague configuration by a finite list of regions or quantities that can be counted exactly. This is the same spirit behind combinatorial proofs, measure decompositions, and even sigma-algebra calculations later in analysis.

§1.10 Extremal problems: upper bound plus construction

The sum-set/difference-set problem has the standard shape of extremal combinatorics. One proves an upper bound by showing that certain auxiliary objects must fit inside a finite container. Then one gives a construction meeting the bound.

This two-part structure is essential:

$$\text{upper bound} + \text{matching construction} = \text{exact extremal value}.$$

A construction alone proves possibility; an upper bound alone proves limitation. A complete extremal solution needs both.

§1.11 Additive combinatorics viewpoint

For a finite set $A \subseteq \mathbb{Z}$, the sum set and difference set are

$$A + A = \{a + a' : a, a' \in A\}, \quad A - A = \{a - a' : a, a' \in A\}.$$

The size of these sets measures additive structure. Arithmetic progressions have unusually small sumsets. Random sets usually have much larger sumsets.

A central theorem in additive combinatorics is Freiman's philosophy: if $|A + A|$ is small compared with $|A|$, then A must resemble a generalized arithmetic progression. The elementary problem is a small concrete case of studying what set structure is forced by sum and difference constraints.

§1.12 Cauchy–Davenport shadow

In a prime cyclic group $\mathbb{Z}/p\mathbb{Z}$, the Cauchy–Davenport theorem says

$$|A + B| \geq \min(p, |A| + |B| - 1).$$

This is the modular analogue of the simple lower bound that a sumset of ordered integer sets contains a chain of at least $|A| + |B| - 1$ elements.

The theorem shows that the lower-bound logic in the note is not accidental. Sumsets are forced to grow unless the ambient group wraps around or the set has special structure.

§1.13 Boolean atoms and Venn regions

For three sets A, B, C , the universe is partitioned into at most eight atoms:

$$A^{\epsilon_1} \cap B^{\epsilon_2} \cap C^{\epsilon_3}, \quad \epsilon_i \in \{0, 1\},$$

where $A^1 = A$ and $A^0 = A^c$. Every Boolean expression in A, B, C is a union of some of these atoms.

This gives a systematic proof method: reduce both sides of a set identity to the same atoms. Venn diagrams are pictures of this atomic decomposition; elementwise proofs are the formal version.

§1.14 Symmetric difference as addition mod two

The symmetric difference

$$A \triangle B = (A \setminus B) \cup (B \setminus A)$$

behaves like addition modulo 2 on characteristic functions:

$$\mathbf{1}_{A \triangle B} = \mathbf{1}_A + \mathbf{1}_B \pmod{2}.$$

Thus the power set $\mathcal{P}(U)$ becomes a vector space over $\mathbb{F}_2$ under symmetric difference. Intersection acts like multiplication.

This explains why expressions such as $(A - B) \cup (B - A)$ have algebraic structure: they are not just set differences, but mod-two sums of predicates.

§1.15 Sperner and intersecting-family context

Many extremal set problems ask for the largest family avoiding a containment or disjointness pattern. Sperner’s theorem says that an antichain $\mathcal{F} \subseteq \mathcal{P}([n])$ satisfies

$$|\mathcal{F}| \leq \binom{n}{\lfloor n/2 \rfloor}.$$

The largest example is the middle layer.

For intersecting families, the Erdős–Ko–Rado theorem says that if $\mathcal{F} \subseteq \binom{[n]}{k}$ is pairwise intersecting and $n \geq 2k$, then

$$|\mathcal{F}| \leq \binom{n-1}{k-1}.$$

The extremal example is a star: all k -sets containing a fixed element.

These theorems are advanced versions of the note’s method: identify the forbidden configuration, prove an upper bound, and classify sharp examples.

§1.16 Shifting and stability

Shifting transforms a set family into a more ordered one without changing its size or destroying the relevant forbidden configuration. Repeated shifting often reduces an extremal problem to a canonical family.

A stability theorem asks more than the maximum: if a family is close to maximum size, must it be structurally close to an extremal example? This is the research-level continuation of checking equality cases in an elementary inequality.

§1.17 Extremal sets, Boolean algebra, and stability

The note combines two important habits. Extremal counting uses auxiliary sets, upper bounds, and sharp constructions. Venn reasoning uses Boolean atoms and set algebra. At a higher level these become additive combinatorics, extremal set theory, vector-space structure over $\mathbb{F}_2$, and stability methods.