

Congruence Classes, Complete Residue Systems, and Euler's Function

N-20221126

§1.1 Congruence

Let m be a positive integer and $a, b \in \mathbb{Z}$. We say that a and b are congruent modulo m , written

$$a \equiv b \pmod{m},$$

if

$$m \mid (a - b).$$

This definition is just divisibility language with a new notation. It allows us to work with remainders rather than whole integers.

§1.2 Basic properties

The note lists the standard properties:

- (i) Reflexivity: $a \equiv a \pmod{m}$.
- (ii) Symmetry: if $a \equiv b \pmod{m}$, then $b \equiv a \pmod{m}$.
- (iii) Transitivity: if $a \equiv b \pmod{m}$ and $b \equiv c \pmod{m}$, then $a \equiv c \pmod{m}$.
- (iv) Compatibility with addition and multiplication:

$$a \equiv b, c \equiv d \pmod{m} \Rightarrow a + c \equiv b + d, \quad ac \equiv bd \pmod{m}.$$

- (v) If $a \equiv b \pmod{m}$, then

$$a^n \equiv b^n \pmod{m}$$

for every positive integer n .

The last two rules are what make congruence useful: one can replace a large number by a small remainder and keep computing.

§1.3 Residue classes

A residue class modulo n is a set

$$M_i = \{x \in \mathbb{Z} : x \equiv i \pmod{n}\}.$$

The integers split into n disjoint classes:

$$M_0, M_1, \dots, M_{n-1}.$$

This is a partition of $\mathbb{Z}$. The note records two important consequences:

- (i) among any $n + 1$ integers, two are congruent modulo n ;
- (ii) if $(i, n) = 1$, then every integer in M_i is coprime to n .

The first is the pigeonhole principle; the second follows because congruent numbers have the same gcd with n .

§1.4 Complete and reduced residue systems

A complete residue system modulo n is any set of n representatives, one from each residue class. The standard example is

$$0, 1, 2, \dots, n - 1.$$

A reduced residue system modulo n consists of one representative from each residue class coprime to n . The number of such classes is Euler's function

$$\varphi(n).$$

If

$$n = p_1^{\alpha_1} \cdots p_k^{\alpha_k},$$

then

$$\varphi(n) = n \left(1 - \frac{1}{p_1}\right) \cdots \left(1 - \frac{1}{p_k}\right).$$

In particular,

$$\varphi(p) = p - 1, \quad \varphi(p^a) = p^a - p^{a-1}.$$

§1.5 Example: last digit

To compute the last digit of 15^{200} , work modulo 10. Since

$$15 \equiv 5 \pmod{10},$$

we get

$$15^{200} \equiv 5^{200} \equiv 5 \pmod{10}.$$

Thus the last digit is 5.

The note's method is to reduce early. Do not expand powers; replace the base by its residue.

§1.6 Example: modulo 17

To compute

$$15^{100} \pmod{17},$$

notice

$$15 \equiv -2 \pmod{17}.$$

Then

$$15^{100} \equiv (-2)^{100} = 2^{100} \pmod{17}.$$

Since

$$2^4 = 16 \equiv -1 \pmod{17},$$

we have

$$2^8 \equiv 1 \pmod{17}.$$

Therefore

$$2^{100} = 2^{96} 2^4 \equiv 1^{12} (-1) \equiv 16 \pmod{17}.$$

§1.7 Example: reducing a long expression

For an expression such as

$$2001 \cdot 2002 + 2003 \cdot 2004 + 2005 \cdot 2006$$

modulo 45, reduce every factor modulo 45 first. The arithmetic becomes small. The main habit is:

$$\text{large expression} \longrightarrow \text{small residues.}$$

This is the same principle behind all modular computation.

§1.8 A polynomial congruence by induction

The note also includes a divisibility proof for an expression depending on k , of the form

$$12^{2k+1} + 36^k + 5^k + 3.$$

The standard method is induction or modular periodicity. One chooses the modulus, checks the base case, and then shows the expression changes by a multiple of the modulus when k increases. Equivalently, reduce each base modulo the modulus and use periodicity of powers.

§1.9 Quadratic residues modulo 8

A consolidation exercise proves that a perfect square modulo 8 is congruent to

$$0, 1, \text{ or } 4.$$

Indeed, every integer is congruent modulo 8 to one of

$$0, 1, 2, 3, 4, 5, 6, 7.$$

Squaring gives residues

$$0, 1, 4, 1, 0, 1, 4, 1.$$

Thus only 0, 1, 4 occur. This is a powerful small filter in number theory problems.

§1.10 Where the idea leads

Congruence classes are quotient thinking. Instead of remembering the whole integer, we remember only its class in

$$\mathbb{Z}/n\mathbb{Z}.$$

Complete residue systems are representatives of all classes; reduced residue systems are the invertible classes. Euler's function counts the units. This note is therefore the elementary entrance to the ring $\mathbb{Z}/n\mathbb{Z}$ and its unit group.

§1.11 Residue systems as representatives

A complete residue system modulo n is a choice of one representative from each class in $\mathbb{Z}/n\mathbb{Z}$. A reduced residue system is a choice of representatives from the unit group

$$(\mathbb{Z}/n\mathbb{Z})^\times.$$

Thus the difference between complete and reduced systems is the difference between the whole quotient ring and its invertible elements.

§1.12 Euler function through prime powers

For a prime power,

$$\varphi(p^k) = p^k - p^{k-1} = p^k \left(1 - \frac{1}{p}\right).$$

Using the Chinese remainder theorem gives

$$\varphi(n) = n \prod_{p|n} \left(1 - \frac{1}{p}\right).$$

§1.13 Orders and exponent reduction

If a is a unit modulo n , its order is the smallest d with $a^d \equiv 1 \pmod{n}$. Exponent reduction is really reduction modulo the order of a , not blindly modulo $\varphi(n)$.

§1.14 Residues, units, and exponent cycles

Residue classes form quotient rings, reduced residue systems form unit groups, Euler's function counts units, and exponent tricks are statements about orders in finite groups.

§1.15 Residue systems as groups of units

A reduced residue system modulo n is the unit group

$$(\mathbb{Z}/n\mathbb{Z})^\times.$$

Its order is Euler's function $\varphi(n)$. Euler's theorem says that if $\gcd(a, n) = 1$, then

$$a^{\varphi(n)} \equiv 1 \pmod{n}.$$

This is simply Lagrange's theorem applied to the finite group of units.

Thus the elementary reduced-residue computations are group theory in disguise.

§1.16 CRT decomposition of unit groups

If m, n are coprime, then

$$\mathbb{Z}/mn\mathbb{Z} \cong \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}.$$

Taking units gives

$$(\mathbb{Z}/mn\mathbb{Z})^\times \cong (\mathbb{Z}/m\mathbb{Z})^\times \times (\mathbb{Z}/n\mathbb{Z})^\times.$$

This explains the multiplicativity of φ .

For prime powers, the structure of the unit group is more subtle. For odd primes p , the group $(\mathbb{Z}/p^k\mathbb{Z})^\times$ is cyclic of order $p^{k-1}(p-1)$. Powers of 2 require separate treatment.

§1.17 Primitive roots and characters

A primitive root modulo n is a generator of the unit group. When the unit group is cyclic, every unit is a power of one element. This converts multiplicative congruence problems into exponent arithmetic.

Dirichlet characters are homomorphisms

$$\chi : (\mathbb{Z}/n\mathbb{Z})^\times \rightarrow \mathbb{C}^\times$$

extended to all integers by setting $\chi(a) = 0$ when $\gcd(a, n) > 1$. They are Fourier characters on the finite abelian group of units and are central in Dirichlet's theorem on primes in arithmetic progressions.

§1.18 Residue systems are calculations inside finite groups

The note's residue tables, reduced systems, and Euler-function formulas are not just modular arithmetic. They are computations inside finite groups. The next levels are group decomposition, primitive roots, and characters.

§1.19 Primitive roots as logarithms in finite groups

When $(\mathbb{Z}/n\mathbb{Z})^\times$ is cyclic with generator g , every unit has the form g^k . This turns multiplication into addition of exponents modulo the group order. In other words, a primitive root supplies a discrete logarithm coordinate.

This is powerful but also limited: not every modulus has primitive roots. The classification says primitive roots exist for

$$n = 1, 2, 4, p^k, 2p^k$$

where p is an odd prime. This explains why some modular arithmetic problems become simple after choosing a primitive root, while others require decomposing the unit group into several cyclic factors.

§1.20 Characters as Fourier analysis on unit groups

Dirichlet characters are not arbitrary functions; they are group homomorphisms from the unit group to complex roots of unity. Orthogonality of characters gives filters for congruence classes. This is the multiplicative analogue of using roots of unity to filter residues modulo n .

The elementary reduced-residue system is therefore the domain on which finite harmonic analysis is performed.