

A
Knockout inhibition of the sprouting (ISV) models across the ensemble
FASTCORE loose
FASTCORE default
FASTCORE strict
GIMME
acaa1 ada acox1 uqcrb lcat nmnat1 slc29a1b cpt2 slc19a3a sdhc sdhb sdhdb sdha suclg1 dldh fh dlst mdh2 slc25a20 slc19a1 slc25a19 acat2 fdps slc25a36a idh2 cmpk ptdss1a
inhibition of sprouting growth
1.0 0.8 0.6 0.4 0.2 0.0
called, two-lineage rule
called, all-lineage rule
conventional-screen candidates (red names)
B
Calls per member
FASTCORE loose 13 0
FASTCORE default 11 0
FASTCORE strict 5 2
GIMME 5 1
0 10
genes called
two-lineage rule
all-lineage rule
C
Media: FASTCORE default
Media: GIMME
slc25a20
slc25a36a
idh2
lcat 1.00 1.00 1.00 1.00 1.00 1.00
slc25a19
sdha 0.14 0.02 0.13 0.14
mdh2 0.09 0.09
cpt2
uqcrb 0.74 0.08 0.46 0.26 0.74
0.05 0.10 0.02 0.04 0.05 0.04
0.50 0.50 0.50 0.50 0.50 0.50
0.77 0.01 0.07 0.36 0.77 0.23
1.00 1.00 1.00 1.00 1.00 1.00
0.92 0.63 0.80 0.88 0.92 0.70
0.77 0.01 0.10 0.36 0.77 0.23
1.00 1.00 1.00 1.00 1.00 1.00
1.00 1.00 1.00 1.00 1.00 1.00
baseline hypoxia ischemia hyperglycemia +carnitine low amino acids
D
slc25a36a call rests on guk1a
threshold
HE_EMP
aTrunk_mixed
vPCV
vPCV_cyc
aCran
ISV_tip
angioblast_cyc
vCV
aDA
cranV
angioblast
cranV_cyc
endocard
ISV_cyc
0.0 0.1 0.2 0.3 0.4
guk1a expression (log1p CP10k)