

# Magnetic and Geometric Properties of the Ising Model on Lattice Random Walks

Ilya Pchelintsev

Scientific Adviser: Burovsky Evgeny Andreevich

The Department of Applied Mathematics

National Research University "Higher School of Economics"

The Moscow Institute of Electronics and Mathematics

Moscow, Russia

iipchelintsev@edu.hse.ru

22nd of March, 2023

# Table of contents


1 Introduction

2 Results

3 Future work

# Ising-ISAW model


- Linear self-avoiding conformation (**SAW**-models)
- Spin subsystem inside of the monomers (regular **Ising model**)
- Close-range interaction (1)
- Tricriticality of the phase transition point [1]




$$H_{u,N,\{\sigma\}} = - \sum_{\langle i,j \rangle} J \sigma_i \sigma_j, \quad i, j \in u, \quad |u| = N \quad (1)$$

# Universality of critical properties of the regular Ising model

$$H_{L,r,\{\sigma\}} = - \sum_{\langle i,j \rangle} J \sigma_i \sigma_j \quad (2)$$



(a) Example of the regular Ising model and its gyration ellipse






(b) Critical cumulant as a function of the lattice ratio [2]

## Question №1

Equal critical geometrical properties  $\Rightarrow$  equal critical magnetic ones?

# Lattice nearest-neighbors modifications



## Question №2

Universality of lattices with equal numbers of dimensions OR equal coordination numbers?

# Observables

Critical cumulant:

$$U_4 = 1 - \frac{\langle m^4 \rangle}{3\langle m^2 \rangle^2} \quad (3)$$

Gyration tensor of SAW-system [3]:

$$Q_{N,\alpha\beta} = \frac{1}{N} \sum_{i=1}^N (w_{i,\alpha} - w_{c,\alpha})(w_{i,\beta} - w_{c,\beta}) \quad (4)$$


Aspect ratio of the SAW-system:

$$r = \sqrt{\frac{\langle q_1 \rangle_N}{\langle q_2 \rangle_N}} \quad (5)$$


Asphericity:

$$\mathcal{A} = \left\langle \frac{(q_1 - q_2)^2}{(q_1 + q_2)^2} \right\rangle_N \quad (6)$$

# Critical Asphericity



(a) Asphericity of the regular Ising as a function of the lattice ratio



(b) Asphericity of the Ising-ISA as a function of  $J$  in the crit. region

Structure	lattice	$J_c$
SAW	Square	0.8340(5) [4]
lattice	Rectangular	$\ln(1 + \sqrt{2})/2$ [5]

Ising-ISA			
J	$\mathcal{A}$	r	$U_4$ Rectangular
0.831	0.415	0.465	$0.338 \pm 0.006$
0.832	0.4072	0.47	$0.343 \pm 0.006$
0.836	0.373	0.492	$0.349 \pm 0.006$

## Result



As for Ising-ISA  $U_4 = 0.308(8)$ , critical cumulant showed **complete mismatch**.





# Bulks of the Square lattice

Monomers according to the number of interactions:

- 2 neighbors - 1D-chains
- 3 neighbors - boundaries of the cluster
- 4 neighbors - core of the cluster




# Bulk results



- First order transition was determined in cubic Ising-ISAW
- Continuous transition in triangular Ising-ISAW
- Clear depiction of the chain consolidation






# Anticipated results



## $J=0$ case research:

- fractions scaling nature as functions of the chain length?
- comparability of the lattice modifications

# References

-  P.-G. de Gennes, *Scaling concepts in polymer physics*. Cornell University Press, 1979.
-  W. Selke, “Critical Binder cumulant of two-dimensional Ising models,” *Eur. Phys. J. B*, vol. 51, no. 2, pp. 223–228, 2006.
-  S. Caracciolo, M. Gherardi, M. Papinutto, and A. Pelissetto, “Geometrical properties of two-dimensional interacting self-avoiding walks at the  $\theta$ -point,” *J. Phys. A: Math. Theor.*, vol. 44, no. 11, pp. 1–24, 2011.
-  K. Faizullina, I. Pchelintsev, and E. Burovski, “Critical and geometric properties of magnetic polymers across the globule-coil transition,” 2021.
-  L. Onsager, “Crystal statistics. i. a two-dimensional model with an order-disorder transition,” *Phys. Rev.*, vol. 65, pp. 117–149, Feb 1944. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRev.65.117>

*Thanks for watching!*