

The Congruence Lattice Problem: A Historical Survey

Štěpán Chrast

February 24, 2026

Abstract

The Congruence Lattice Problem (CLP) asks whether every distributive algebraic lattice is isomorphic to the congruence lattice of a lattice. After early positive results in the finite case and several infinite cardinalities, the problem was solved negatively by Wehrung, who constructed a distributive algebraic lattice that is not representable as a congruence lattice. This article provides a historical and structural overview of the problem, the main positive results, and the ideas behind the counterexample and the optimal cardinal bound.

Contents

1	Introduction	2
2	Preliminaries	2
3	The Finite Case	3
4	Partial Results for Infinite Lattices	3
5	Obstructions in the Infinite Case	3
6	Wehrung's Counterexample and the Negative Solution	4
7	Conclusion	5

1 Introduction

The Congruence Lattice Problem (CLP) is a classical question in lattice theory concerning the possible structures of congruence lattices of lattices. Attributed to Dilworth in the 1940s and open for more than sixty years, it became a central problem in the interaction between lattice theory and universal algebra.

The problem originates from a fundamental observation about congruences: In 1942, Funayama and Nakayama proved that for any lattice L , the congruence lattice of L is distributive [2]. This result led to the inverse question of which distributive lattices can arise as congruence lattices of some lattice.

Later in 1948, Birkhoff and Frink proved that for any algebra A , the congruence lattice $\text{Con}(A)$ is an *algebraic* lattice. Consequently, distributivity alone cannot characterize congruence lattices of lattices.

Together, these observations form the following fact about lattices:

For any lattice L , the congruence lattice of L is a distributive algebraic lattice.

From this observation, one might ask whether the inverse is true - whether those properties are enough for a lattice to be the congruence lattice of some lattice L :

Is every distributive algebraic lattice isomorphic to the congruence lattice of some lattice L ?

This question became known as the Congruence Lattice Problem.

A crucial structural feature of algebraic lattices is that they are determined by their compact elements. In the case of congruence lattices, their compact elements are the finitely generated congruences, which form a distributive $(\vee, 0)$ -semilattice $\text{Con}_c(L)$. Accordingly, many authors reformulate the CLP in the language of distributive $(\vee, 0)$ -semilattices in the following equivalent formulation [3]:

Is every distributive $(\vee, 0)$ -semilattice S isomorphic to $\text{Con}_c(L)$ of some lattice L ?

The aim of this paper is to present a brief historical and conceptual survey of the CLP. After recalling the necessary background in lattice theory, we outline the finite solution and the subsequent partial results for infinite lattices. We then describe the structural obstructions that emerge in higher cardinalities and show Wehrung's counterexample, the negative resolution of the Congruence Lattice Problem and further results.

2 Preliminaries

To formulate the Congruence Lattice Problem, we first recall several basic notions from lattice theory:

Distributive lattice. A *lattice* is an algebra $L = (L, \vee, \wedge)$ consisting of a nonempty set equipped with two binary operations, join ($\vee$) and meet ($\wedge$), that are idempotent, commutative, associative, and satisfy the absorption identities. A lattice is called *distributive* if it satisfies the distributive law

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$

for all $x, y, z \in L$.

Congruence. A *congruence relation* on a lattice L is an equivalence relation θ such that: if $x \equiv_\theta y$ and $u \equiv_\theta v$, then

$$x \vee u \equiv_\theta y \vee v, \quad x \wedge u \equiv_\theta y \wedge v.$$

The set of all congruences of some algebra A , ordered by inclusion, forms a lattice denoted $\text{Con}(A)$.

Complete lattice. A lattice is *complete* if every subset has a supremum and an infimum.

Compact element. Let L be a complete lattice. An element $x \in L$ is *compact* if for every $A \subseteq L$,

$$x \leq \bigvee A \Rightarrow \exists F_{\text{finite}} \subseteq A \text{ such that } x \leq \bigvee F.$$

Algebraic lattice. A lattice L is *algebraic* if it is complete and every element is the supremum of the compact elements below it.

$\text{Con}_c(A)$. Let A be an algebra. The set of compact elements of the lattice $\text{Con}(A)$ forms a distributive $(\vee, 0)$ -semilattice, denoted by $\text{Con}_c(A)$.

Representability. A distributive $(\vee, 0)$ -semilattice S is *representable* if S is isomorphic to the $\text{Con}_c(L)$ for some lattice L , as $(\vee, 0)$ -semilattices.

Weakly distributive homomorphisms [3]. A homomorphism of join-semilattices $\mu : S \rightarrow T$ is *weakly distributive at* $x \in S$ if whenever $\mu(x) \leq y_0 \vee y_1$ in T , there exist $x_0, x_1 \in S$ such that $x \leq x_0 \vee x_1$ and $\mu(x_i) \leq y_i$ for $i < 2$. We say that μ is *weakly distributive* if it is weakly distributive at every $x \in S$.

For more details see [1].

3 The Finite Case

For the finite cases of lattices, the Congruence Lattice Problem was actually solved before the general CLP was proposed and the notion of algebraicity was introduced in this context. In the 1940s, Dilworth discovered a theorem concerning finite distributive lattices, with a complete proof being later published by Grätzer and Schmidt in 1963. The theorem states that:

Theorem 3.1. *Every finite distributive lattice is isomorphic to the congruence lattice of some finite lattice.*

For proof see [4, Theorem 8.1, p. 101].

Since every finite lattice is complete and every element of a finite lattice is compact, every finite lattice is algebraic, this theorem provides a positive solution for finite lattices of the congruence lattice problem:

Every finite (and therefore algebraic) distributive lattice is isomorphic to the congruence lattice of some (finite) lattice.

4 Partial Results for Infinite Lattices

The finite solution of the Congruence Lattice Problem does not extend directly to infinite lattices, but some strong partial results were obtained for infinite distributive lattices.

In case of infinite lattices, algebraicity is not automatic: not every distributive lattice is algebraic in the finite case. Distributive lattices that are not algebraic are not isomorphic to the congruence lattice of some lattice, hence in the infinite setting we need to focus on algebraic distributive lattices.

A sequence of positive results was obtained during the 1970s–1990s, establishing representability for broad classes of infinite algebraic distributive lattices. Building on structural conditions introduced by E. T. Schmidt, several authors proved representability under certain cardinalities and other properties:

Theorem 4.1 (Wehrung). *Every countable union of distributive lattices with zero is representable.*

Theorem 4.2 (Dobbertin). *Every distributive join-semilattice with zero in which every principal ideal is countable is representable.*

Theorem 4.3 (Huhn). *Every distributive join-semilattice with zero of cardinality at most $\aleph_1$ is representable.*

for details, see [6, p. 705].

Together, these and related results established representability for several broad classes of infinite distributive $(\vee, 0)$ -semilattices, covering cardinalities (on the semilattice side) up to $\aleph_1$. For a long time, no counterexample to the Congruence Lattice Problem was known, while partial positive solutions have been accumulating. Thus leading to further strengthening the confidence that the CLP could be true.

5 Obstructions in the Infinite Case

All those positive results show that many distributive $(\vee, 0)$ -semilattices are representable at small cardinalities, but that does not automatically extend to the general infinite case. In higher cardinalities, some structural obstructions emerge.

A key fact leading to Wehrung's counterexample is the following observation:

Observation 5.1. *Let S be a distributive $(\vee, 0, 1)$ -semilattice. If $S \cong \text{Con}_c(L)$ for some lattice L , then there exists a weakly distributive $(\vee, 0)$ -homomorphism $\mu : \text{Con}_c(L) \rightarrow S$, such that $1 \in \text{rng}(\mu)$. [3]*

With this observation, finding a distributive $(\vee, 0)$ -semilattice such that there is not such weakly distributive homomorphism μ , one would prove CLP to be false, which is exactly what Wehrung did with his counterexample.

6 Wehrung's Counterexample and the Negative Solution

We now sketch the mechanism behind Wehrung's negative solution. The proof is formulated on the level of compact congruences $\text{Con}_c(L)$ and a target distributive $(\vee, 0, 1)$ -semilattice $\mathcal{G}(\Omega)$.

Theorem 6.1 (Wehrung, Theorem 6.1). *Let Ω be a set of cardinality at least $\aleph_{\omega+1}$, and let A be an algebra. If A has a congruence-compatible structure of a $(\vee, 1)$ -semilattice, then there is no weakly distributive $(\vee, 0)$ -homomorphism $\mu : \text{Con}_c(A) \rightarrow \mathcal{G}(\Omega)$ with 1 in its range.*

The semilattice $\mathcal{G}(\Omega)$ construction (sketch)

Fix a set Ω . Start with the $(\vee, 0, 1)$ -semilattice $A(\Omega)$ generated by 1 and elements

$$a_\xi^0, a_\xi^1 \quad (\xi \in \Omega),$$

subject to the relations $a_\xi^0 \vee a_\xi^1 = 1$ for all $\xi \in \Omega$. Wehrung then applies a distributive extension construction D and sets

$$\mathcal{G}(\Omega) = D(A(\Omega)).$$

A key feature is the *finite-support* property: every $x \in \mathcal{G}(\Omega)$ has a least finite support $\text{supp}(x) \subseteq \Omega$ such that $x \in \mathcal{G}(\text{supp}(x))$.

Idea of the proof of Theorem 6.1

Assume toward a contradiction that $\mu : \text{Con}_c(A) \rightarrow \mathcal{G}(\Omega)$ is weakly distributive and $1 \in \text{rng}(\mu)$. Then:

- Using that $1 \in \text{rng}(\mu)$, pick $\alpha \in \text{Con}_c(A)$ with $\mu(\alpha) = 1$ and write α as a finite join of principal congruences, yielding a finite decomposition $1 = \bigvee_{r < m} \mu\Theta_A(t_r, 1)$.
- For each $\xi \in \Omega$, apply weak distributivity to the decomposition $1 = a_\xi^0 \vee a_\xi^1$ in $\mathcal{G}(\Omega)$, producing finite *alternating* congruence chains in A whose successive links map alternately below a_ξ^0 and a_ξ^1 .
- Using finite supports in $\mathcal{G}(\Omega)$ and a Kuratowski free-set argument (available at $|\Omega| \geq \aleph_{\omega+1}$), Wehrung applies his technical lemmas (Erosion + Evaporation) to force a collapse of these chains, contradicting the initial finite decomposition.

□

For details of the construction and proof of Wehrung's theorem, see [3, Theorem 6.1, p. 619].

From Theorem 6.1 to the negative solution of CLP

Let Ω be as in Theorem 6.1 and put $S := \mathcal{G}(\Omega)$. If S were representable, i.e. if $S \cong \text{Con}_c(K)$ for some lattice K , then the isomorphism $\text{Con}_c(K) \rightarrow S$ would be weakly distributive and would map some element to 1, contradicting Theorem 6.1. Hence $\mathcal{G}(\Omega)$ is not representable. Passing from semilattices to algebraic lattices via the ideal lattice $\text{Id}(\mathcal{G}(\Omega))$, one obtains an algebraic distributive lattice that is not isomorphic to $\text{Con}(K)$ for any lattice K , so the Congruence Lattice Problem has a negative answer.

For details and other results of Wehrung's counterexample, see [3].

Růžička's optimal bound

Růžička later improved the cardinal bound to the optimal value $\aleph_2$: there exists a distributive $(\vee, 0, 1)$ -semilattice G of size $\aleph_2$ such that no weakly distributive $(\vee, 0)$ -homomorphism $\text{Con}_c(A) \rightarrow G$ has 1 in its range. In particular, a counterexample to representability already exists at $\aleph_2$. This bound is optimal because every distributive $(\vee, 0)$ -semilattice of cardinality at most $\aleph_1$ is representable (Huhn), so $\aleph_2$ is the smallest cardinal at which non-representability can occur.

More about Růžička's results in [5]

7 Conclusion

The Congruence Lattice Problem has a negative solution: not every distributive algebraic lattice is isomorphic to the congruence lattice of a lattice. The obstruction appears only at the level of higher cardinalities, while representability holds in all finite cases and for distributive $(\vee, 0)$ -semilattices of cardinality at most $\aleph_1$. Wehrung’s construction and Růžička’s optimal bound precisely determine the cardinal threshold, at which the nonrepresentability occurs.

References

- [1] G. Grätzer, *General Lattice Theory*, 2nd ed., new appendices by the author with B. A. Davey, R. Freese, B. Ganter, M. Greferath, P. Jipsen, H. A. Priestley, H. Rose, E. T. Schmidt, S. E. Schmidt, F. Wehrung, and R. Wille, Birkhäuser Verlag, Basel, 1998, xx+663 pp.
- [2] N. Funayama and T. Nakayama, *On the distributivity of a lattice of lattice-congruences*, Proceedings of the Imperial Academy (Tokyo) **18** (1942), 553–554.
- [3] F. Wehrung, *A solution to Dilworth’s congruence lattice problem*, Advances in Mathematics **216** (2007), no. 2, 610–625. doi: 10.1016/j.aim.2007.05.016.
- [4] G. Grätzer, *The Congruences of a Finite Lattice: A “Proof-by-Picture” Approach*, 3rd ed., Birkhäuser / Springer Nature Switzerland, 2023. ISBN 978-3-031-29062-6. doi: 10.1007/978-3-031-29063-3.
- [5] P. Růžička, *Free trees and the optimal bound in Wehrung’s theorem*, preprint (dated 24/9/2007).
- [6] G. Grätzer, *Two Problems That Shaped a Century of Lattice Theory*, Notices of the American Mathematical Society **54** (2007), no. 6, 696–707.