

Exercise 10 (online: 03.07.2023. Return by: **Mo 10.07.2023 10:00**) **14P****1. Magnetic nanoparticles** **10P**

The Stoner-Wohlfarth model describes the angular dependence of the energy E of a spherical, single-domain magnetic nanoparticle with uniaxial crystalline anisotropy by

$$E = K_1 V \sin^2(\theta) - \mu_0 H M_s V \cos(\theta - \phi).$$

Here, θ is the angle between the magnetization \mathbf{M} and the easy axis, and ϕ is the angle between the external magnetic field \mathbf{H} (with $|\mathbf{H}| = H$) and the easy axis. Within the entire volume of the nanoparticle one has $|\mathbf{M}| = M_s$ (saturation magnetization). K_1 is the anisotropy constant, and $\mu_0 \approx 1.26 \cdot 10^{-6} \text{ Vs/(Am)}$ is the permeability of free space. Hint: To shorten your calculations, normalize the energy as $\epsilon = E/(K_1 V)$ and magnetic field as $h = H/H_a$, where $H_a = 2K_1 V/(\mu_0 M_s V)$ is the anisotropy field.

- (a) **Stability criterion and magnetization reversal.** For the case $\phi = \pi$ and $h \geq 0$ find the stable configurations θ of a magnetic nanoparticle. Show that by increasing h the magnetization reverses at $h = 1$ if the particle was in the state $\theta = 0$ at $h < 1$. **(3P)**
- (b) **Angle dependence and hysteresis.** Plot the normalized particle energy $\epsilon(\theta)$ for an external magnetic field with an angle to the easy axis $\phi = \pi, (5/6)\pi, (2/3)\pi$ and $\pi/2$ (4 plots in total). In each case use for plotting at least five different magnetic field amplitudes $h = 0, 0.25, 0.5, 0.75$ and 1 (or more). Describe qualitatively the behaviour of the particle magnetization with increasing external field h for the different θ and discuss the connection to the hysteresis loops presented in the manuscript in Fig. 5.18. **(4P)**
- (c) **Thermally activated magnetization reversal.** Consider a spherical magnetic nanoparticle with $K_1 = 0.2 \text{ J/cm}^3$ and a radius $R = 4 \text{ nm}$ without externally applied magnetic field. How long (on average) should one measure the magnetization of the particle to observe thermally activated magnetization reversal at (i) $T_1 = 300 \text{ K}$, (ii) $T_2 = 77 \text{ K}$? Assume a characteristic time $\tau_0 = 1 \text{ ns}$. The Boltzmann constant is $k_B = 1.38 \cdot 10^{-23} \text{ J/K}$. **(2P)**
- (d) **Memory medium.** Assume that a memory medium based on spherical magnetic nanoparticles with $K_1 = 0.2 \text{ J/cm}^3$ and $\tau_0 = 1 \text{ ns}$ should be developed, which guarantees preserving the information for 100 years at room temperature. How large (at least) should be the diameter of the nanoparticles to implement such a memory? Hint: ignore possible effects due to magnetic domains. **(1P)**

2. **Stoner-Wohlfarth nanoparticle with negative uniaxial anisotropy** 4P

Consider a uniaxial nanomagnetic particle, the energy of which is described by a Stoner-Wohlfarth model

$$E = K_1 V \sin^2 \theta - \mu_0 H M_s V \cos(\theta - \phi) \quad (1)$$

where $H = |\mathbf{H}|$ is the magnitude of an external magnetic field, $M_s = |\mathbf{M}|$ is the magnitude of the particle magnetization (constant and homogeneous at all times), and V the particle volume. In contrast to the usual case, the anisotropy constant shall be negative $K_1 < 0$.

Hint: To shorten your calculations, normalize the energy as $\epsilon = E/(|K_1|V)$ and magnetic field as $h = H/H_a$, where $H_a = 2|K_1|V/(\mu_0 M_s V)$ is the so-called anisotropy field.

- (a) **Sketch the system.** Make a simple drawing showing the directions of \mathbf{M} and \mathbf{H} and explain the two angles θ and ϕ . Consider the case $H = 0$ and plot the normalized energy $\epsilon(\theta)$ for $\theta = -\pi \dots + \pi$ in a separate graph. What is the meaning of $\theta = 0$, which axis is described by this? (2P)
- (b) **Particle groundstate.** We introduce a Cartesian coordinate system with z being the coordinate along $\theta = 0$. In which direction (x, y, z) is the magnetization of the particle pointing for $H = 0$? Could this particle be used for magnetic storage of one bit of information? (2P)