

Index

- absorption
 - large polaron, 75–77
 - large-polaron low-frequency Drude, 78–79
 - small polaron, 77–78, 83
 - small-(bi)polaron low-frequency polarization
 - conductivity, 79–80
 - softening bipolaron, 63, 78
- adiabatic
 - formalism, 21–23
 - limit, definition, 20, 29
 - small-polaron hopping, 101
- bandwidth, polaron
 - strong-coupling, 34
 - strong-coupling, large polaron, 34, 42
 - strong-coupling, small polaron, 35–36, 42
 - weak-coupling, 18
- barrier to self-trapping
 - free-carrier small-polaron dichotomy, 30
 - large-polaron small-polaron dichotomy, 32
 - persistent photo-conductivity, 31
 - small-polaron formation, 30
 - time delay for self-trapping, 30
- bipolaron
 - attraction between large bipolarons, 164–165
 - definition, 3, 54
 - electron correlation, 57–58
 - energy barrier to formation, 57
 - ferroelectric materials, 56
 - inert-pair effect, 59
 - Jahn–Teller, 55, 63
 - large, 54, 55–58
 - large four-lobed, 58
 - negative U , 59
 - pair-breaking in semiclassical hopping, 121–123
 - small, 54, 58–59, 96
 - softening, 55, 59–64, 78
 - softening, boron carbides, 59–60, 63–64
- bipolarons
 - mutual interactions, 164–165
- Born–Oppenheimer method. *See* adiabatic formalism
- boron carbides
 - neutron detector, 133
 - properties, 59–60
 - Seebeck coefficient, 132
 - softening bipolaron, 63–64
- carrier mobilities
 - conductivity mobility, 135
 - drift mobility, 135
 - Hall mobility, 135
 - standard, 135
 - time-of-flight mobility, 135
- carrier-induced atomic changes
 - altered atomic equilibrium positions, 1
 - altered atomic vibration frequencies, 1
- carrier-induced softening
 - atomic desorption, 186–190
 - causes, 129–130
 - definition, 25
 - electronic polarizability, 26, 92
 - enhanced Seebeck coefficient, 130–133
 - Holstein’s molecular-crystal model, 27, 52
 - increased thermal resistivity, 130
 - large-polaron scattering, 92
 - molecular polaron, 51
 - phonon localization, 130
 - Seebeck coefficient enhancement, 64
 - softening bipolaron, 59–64
 - strong-coupling polaron, 28–29, 33–34
 - vibrations’ free-energy, 28–29, 33–34
 - weak-coupling polaron, 13–20
- chalcogenide glasses
 - Hall effect sign anomalies, 158
 - optical properties, 84–85
 - small polarons, 185
 - threshold switching, 184
- coherent transport
 - definition, 86
 - Hall mobility, 136–138
 - large polaron, 88
 - minimum mobility, 87
- desorption
 - carrier-induced softening, 187–189
 - delayed desorption, 187, 190

- desorption (cont.)
 - immediate desorption, 187, 190
 - surface electronic localization, 186–187
- dielectric constants
 - covalent materials, 10
 - ferroelectric materials, 10
 - ionic materials, 10
 - large-bipolaron formation, 56
 - optical, 10
 - static, 10
- effective mass, polaron
 - at high frequencies, 89
 - large-polaron, atomic tunneling regime, 89–91
 - large-polaron, classical treatment, 88–89
 - strong coupling, 3
 - strong coupling, large polaron, 34, 88–91
 - weak coupling, 14
- electronic polarizability, 26
- electronic transfer energy, 100–101
- electronic velocity operator, 103
- electron–phonon interactions, 9–12
 - deformation-potential short-range interaction, 2, 11, 28, 30, 31
 - Fröhlich coupling constant, 14
 - Fröhlich long-range interaction, 2, 9, 28, 30
- exciton
 - Frenkel, 82
 - self-trapped, 82
 - Wannier, 82
- Frenkel–Poole effect, 118
 - for polar molecular solids, 51
- geminate recombination
 - definition, 81
 - Onsager, 81
 - small polaron, 81–82
- Hall effect
 - adiabatic Hamiltonian, 23
 - adiabatic small-polaron hopping, 147–155
 - Hall coefficient, definition, 135
 - Hall mobility, Boltzmann solution, 136–138
 - Hall mobility, definition, 135
 - LaMnO₃, 47–48
 - magnetic phase factors, 139
 - MnO, 158
 - non-adiabatic small-polaron hopping, 141–147
 - sign anomalies, 3, 47–48, 137, 155–159
 - sign anomalies in amorphous covalent semiconductors, 157–158
 - sign anomalies in chalcogenides glasses, 158
 - small-polaron hopping, square lattice, 143–146
 - small-polaron hopping, triangular lattice, 141–143, 147–155
- Holstein’s molecular-crystal model, 32, 51–52
 - for molecular polarons, 51–52
 - for transition-metal oxides, 27, 52
 - hopping activation energy, 106
 - pseudo-molecules, 27, 52
- hopping. *See* phonon-assisted hopping
- incoherent transport
 - definition, 86
 - small bipolaron, 96
 - small polaron, 95–96, 138
- large bipolaron
 - Drude absorption, 169
 - ghost modes, 169
 - ordering, 170
 - photo-absorption, 168
 - photo-emission, 168
 - positron annihilation, 169
 - superconductivity. *See* superconductivity
 - suppression of Seebeck’s spin entropy in cuprates, 171
- large polaron
 - absorption, 75–77
 - definition, 3
 - Drude absorption, 78–79
 - Hall mobility, 138
- large-polaron formation
 - adiabatic limit, scaling analysis, 30
 - quasi-one-dimensional, 32
- large-polaron motion
 - coherent, 3, 88
 - Drude tail, 94
 - effective mass, 34, 88–91
 - Einstein-phonon scattering limit, 94
 - Hall mobility, 138
 - mean-free-path, 93
 - mobility, 93, 94
 - phonon scattering, 3, 91–94
 - phonons’ thermal diffusion, 94
- light-atom diffusion
 - Condon approximation breakdown, 192–193
 - high-temperature hopping, 191–193
 - hydrogen diffusion in bcc metals, 193
 - low-temperature isotope dependence, 196–197
 - semiclassical isotope dependence, 193–196
 - versus classical diffusion, 195
 - versus small-polaron hopping, 195
- luminescence
 - exciton, 82
 - optical cooling, 83–84
 - self-trapped exciton, 82
 - small polaron, 82–83
 - Stokes shift heating, 83
- magnetic field
 - affect on atoms’ motion, 23
- magnetic polaron
 - bound magnetic polaron BMP, 68, 69–72
 - colossal magnetoresistance CMR, 65, 71
 - definition, 65
 - EuO, 65, 71–72
 - EuS, 71
 - EuSe, 71
 - EuTe, 65, 68, 71

- ferromagnetism, 65
- magnetic susceptibility, 65, 68
- small-polaron collapse, 68
- small-polaron collapse, thermally induced, 68–72
- stability, 67–68
- magnetic semiconductors
 - small-polaron hopping, 123–124
- Meyer–Neldel compensation effect, 116
 - molecular polaron, 51
- molecular polaron
 - definition, 49
 - Frenkel–Poole behavior, 51
 - Meyer–Neldel compensation effect, 51
 - scaling analysis, 49–51
 - Seebeck coefficient enhancement, 51
- Peltier heat
 - definition. *See* Seebeck coefficient
- persistent photo-conductivity. *See* barrier to self-trapping
- phonon-assisted hopping
 - activation energy, 106, 115
 - adiabatic semiclassical small-polaron, 110–117
 - adiabatic small-polaron, 98
 - adiabatic versus non-adiabatic regimes, 116–117
 - bipolaron pair breaking, 121–123
 - detailed balance, 108
 - Frenkel–Poole effect, 117–118
 - Hall effect sign anomalies, 155–159
 - inverted regime, 108
 - jump distance dependence, 117
 - magnetic semiconductors, 123–124
 - negative differential conductivity, 108
 - non-adiabatic impurity conduction, 98
 - non-adiabatic semiclassical small polaron, 108–110
 - non-adiabatic small polaron, 102–108
 - non-Ohmic at high fields, 180–184
 - non-Ohmic at low temperatures, 175–180
 - non-Ohmic hopping formalism, 173–175
 - shallow impurity, 97
 - small polaron, 97–102
 - threshold switching, 180–185
- polaron
 - absorption, 75–78
 - solvated electron, 3
 - strong-coupling, definition, 1
 - weak-coupling, definition, 1
- polaron radius
 - quasi-one-dimensional polaron, 32
 - strong-coupling, large polaron, 30, 34, 42
 - strong-coupling, small polaron, 31, 42
 - weak-coupling, 13
- rare-earth chalcogens
 - EuO, 65, 71–72, 133
 - EuS, 71
 - EuSe, 71
 - EuTe, 65, 68, 71
- recombination
 - geminate, 81–82
 - luminescence, 82–84
 - non-radiative, 83–84
- scaling analysis. *See* self-trapping: adiabatic limit
- Seebeck coefficient
 - carrier-induced softening's enhancement, 53, 130–133
 - definition, 125
 - hopping carrier, 127–129
 - LaMnO₃, 47
 - low-temperature limit, 127
 - magnetic semiconductors, 133–134
 - magnetic-field dependence, 126
 - narrow-band limit, 125–126
 - polaron hopping, 128
 - singlet bipolaron, 126
 - spin degeneracy, 126
 - trap-modulated conduction, 129
 - with temperature-dependent energy levels, 126–127
- self-trapping
 - adiabatic approach, 1, 23–29
 - adiabatic limit, scaling analysis, 29–32, 189–190
 - barrier to self-trapping, 30, 190
 - beyond adiabatic limit, 33–39
 - definition, 1, 42
 - desorption, 186–187
 - electronic dimensionality effect, 29, 31–32, 189–190
 - electronically constrained, 2
 - nonlinear eigenvalue equation, 27
 - nonlinearity, 2
- small polaron
 - absorption, 77–78
 - definition, 3
- small polaron, defect-related
 - from breaking a trapped electron pair, 45
 - from trapping a large polaron, 44
- impurity conduction. *See* small-polaron motion
- trapped small polaron, 44
- small-polaron formation
 - adiabatic limit, scaling analysis, 30
 - band narrowing, 36
 - bandwidth, 35
 - barrier to self-trapping, 30, 32
 - beyond adiabatic limit, 33–39
- small-polaron formation, coexistence. *See* barrier to self-trapping
 - with free carriers, 30, 31, 36–39
 - with large polarons, 32
- small-polaron formation, induced by, 42
 - disorder, 41
 - electronic localization, 39–40
 - long-range electron–phonon interaction, 32
 - magnetic disorder, 5, 68–72
- small-polaron motion
 - adiabatic hopping Hall Effect, 147–155
 - adiabatic semiclassical hopping, 110–117
 - adiabatic versus non-adiabatic hopping, 116–117
 - bipolaron pair breaking, 121–123
 - coherent with atomic tunneling, 35–36
 - coherent-motion's suppression, 36, 96, 138
 - Frenkel–Poole effect, 117–118
 - generalized for light-atom's hopping, 191–193
 - Hall-effect sign anomalies, 3, 45–48, 155–159
 - hopping, 3, 36, 98–102

- small-polaron motion (cont.)
 - hopping activation energy, 106, 115
 - hopping Hall effect. *See* Hall effect
 - impurity conduction, 45–48
 - incoherent, 3, 95–96
 - incoherent for classical atoms, 35, 95–96
 - jump-distance dependence, 117
 - low-frequency polarization conduction, 79–80
 - magnetic semiconductors, 123–124
 - non-adiabatic Hall effect, 141–143
 - non-adiabatic hopping, 102–108
 - non-adiabatic semiclassical hopping, 108–110
 - non-Ohmic hopping, 180–185
 - photo-conductivity, 81
 - pinned Fermi level, 45
 - Seebeck coefficient, 127–129
 - Seebeck coefficient, magnetic semiconductors, 133–134
 - temporally correlated hopping, 118–119
 - thermally induced conductivity transition, 39, 42
 - threshold switching model, 180–185
 - time-of-flight mobility, 119–120
- strong-coupling polaron
 - effective mass, 3, 34
 - Fröhlich polaron energy, 14
 - large. *See* large polaron
 - optical absorption, 2
 - small. *See* small polaron
- superconductivity
 - BCS condensate, 163, 165
 - compared with superfluidity, 163
 - large bipolarons, 164
 - large-bipolaron liquid, 165–166
 - large-bipolaron properties, 168–172
 - large-bipolarons' excitation spectrum, 167–168
 - large-bipolarons' proximity effect, 172
 - mutual attraction between large bipolarons, 164–165
 - small- versus large-bipolaronic superconductivity, 163
 - superfluid helium's condensate, 163
 - suppression by large-bipolaron ordering, 171
- thermoelectric figure-of-merit, 132
- threshold switching, 185
 - chalcogenide glasses, 185
 - small-polaron model, 180–185
- time delay for self-trapping. *See* barrier to self-trapping
- time-of-flight mobility, 119–120
- transition-metal oxides
 - La_2CuO_4 , 170, 171
 - LaMnO_3 , 45–48
 - LiNbO_3 , 85
 - MnO , 43, 52, 133, 158
 - NiO , 44
 - SrTiO_3 , 44, 170
- vibrational dispersion
 - adiabatic small-polaron Hall effect, 150
 - adiabatic small-polaron hopping, 112–116
 - correlated hopping, 118–120
 - large-polaron scattering, 93–94
 - phonon-assisted hopping, 98
 - semiclassical small-polaron hopping, 109
 - small-polaron hopping, 105
- Wannier–Stark localization, 86–87, 103
- weak-coupling polaron
 - carrier-induced softening, 15–20
 - deformation-potential short-range electron–phonon interaction, 14
 - effective mass, 14
 - Fröhlich long-range electron–phonon interaction, 13
 - Fröhlich polaron energy, 14