

Hamiltonian

I am considering the interaction term H_I of a Hamiltonian of a system of N particles with positions $\{\bar{\mathbf{R}}_l\}_{l \in \{1, \dots, N\}}$. It shall consist of two-body interactions, which depend on the position of the involved particles. Thus H_I can be written in the form

$$H_I(\{\bar{\mathbf{R}}_l\}) = \sum_{i,j; i \neq j} V_{i,j}(\bar{\mathbf{R}}_i, \bar{\mathbf{R}}_j)$$

In order to being able to derive the second order expansion of H_I , that will be needed later, in a more formal way, I write the two-body interactions as dependend on all positions.

$$H_I(\{\bar{\mathbf{R}}_l\}) = \sum_{i,j; i \neq j} V_{i,j}(\{\bar{\mathbf{R}}_l\})$$

The $V_{i,j}$ are formally functions of all positions $\{\bar{\mathbf{R}}_l\}$, i.e $V_{i,j}(\{\bar{\mathbf{R}}_l\})$. But of course they only depend on $\bar{\mathbf{R}}_i$ and $\bar{\mathbf{R}}_j$, i.e.

$$\frac{\partial V_{i,j}}{\partial \bar{\mathbf{R}}_l} = 0 \quad \text{if } i \neq l \neq j$$

Now I expand the interaction Hamiltonian in small displacements $\{\delta \mathbf{R}_l\}$ from the tweezer centers $\{\mathbf{R}_l\}$.

$$\begin{aligned} H_I(\{\mathbf{R}_l + \delta \mathbf{R}_l\}) &= H_I(\{\mathbf{R}_l\}) + \sum_n \frac{\partial H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n} \delta \mathbf{R}_n + \frac{1}{2} \sum_{n,m} \delta \mathbf{R}_n \frac{\partial^2 H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n \partial \mathbf{R}_m} \delta \mathbf{R}_m \\ &= H_I(\{\mathbf{R}_l\}) + \sum_n \frac{\partial H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n} \delta \mathbf{R}_n + \frac{1}{2} \sum_{n \neq m} \delta \mathbf{R}_n \frac{\partial^2 H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n \partial \mathbf{R}_m} \delta \mathbf{R}_m + \frac{1}{2} \sum_{n=m} \delta \mathbf{R}_n \frac{\partial^2 H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n \partial \mathbf{R}_m} \delta \mathbf{R}_m \\ &= H_I(\{\mathbf{R}_l\}) + \sum_n \frac{\partial H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n} \delta \mathbf{R}_n + \frac{1}{2} \sum_{n \neq m} \sum_{i,j; i \neq j} \delta \mathbf{R}_n \frac{\partial^2 V_{i,j}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n \partial \mathbf{R}_m} \delta \mathbf{R}_m \\ &\quad + \frac{1}{2} \sum_n \sum_{i,j; i \neq j} \delta \mathbf{R}_n \frac{\partial^2 V_{i,j}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n^2} \delta \mathbf{R}_n \\ &= H_I(\{\mathbf{R}_l\}) + \sum_n \frac{\partial H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n} \delta \mathbf{R}_n + \frac{1}{2} \sum_{n \neq m} \sum_{i,j; i \neq j} \delta \mathbf{R}_n \frac{\partial^2 V_{i,j}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n \partial \mathbf{R}_m} \delta \mathbf{R}_m (\delta_{n=i, m=j} + \delta_{n=j, m=i}) \\ &\quad + \frac{1}{2} \sum_n \sum_{i,j; i \neq j} \delta \mathbf{R}_n \frac{\partial^2 V_{i,j}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n^2} \delta \mathbf{R}_n (\delta_{n=i} + \delta_{n=j}) \end{aligned}$$

Using the fact, that

$$\delta \mathbf{R}_i \frac{\partial^2 V_{i,j}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_i \partial \mathbf{R}_j} \delta \mathbf{R}_j = \delta \mathbf{R}_j \frac{\partial^2 V_{i,j}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_j \partial \mathbf{R}_i} \delta \mathbf{R}_i$$

the expansion can be written as

$$\begin{aligned} H_I(\{\mathbf{R}_l + \delta \mathbf{R}_l\}) &= H_I(\{\mathbf{R}_l\}) + \sum_i \frac{\partial H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_i} \delta \mathbf{R}_i + \sum_{i,j; i \neq j} \delta \mathbf{R}_i \frac{\partial^2 V_{i,j}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_i \partial \mathbf{R}_j} \delta \mathbf{R}_j \\ &\quad + \sum_{i,j; i \neq j} \delta \mathbf{R}_i \frac{\partial^2 V_{i,j}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_i^2} \delta \mathbf{R}_i \end{aligned}$$

If I introduce a matrix U the expanded Hamiltonian can be brought into the same form as in the paper.

$$H_I(\{\mathbf{R}_l + \delta \mathbf{R}_l\}) = H_I(\{\mathbf{R}_l\}) + \sum_n \frac{\partial H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n} \delta \mathbf{R}_n + \sum_{n,m} \delta \mathbf{R}_n U_{n,m} \delta \mathbf{R}_m$$

with

$$U_{n,m} = \frac{\partial^2 V_{n,m}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n \partial \mathbf{R}_m} \quad \text{if } n \neq m$$

$$U_{n,n} = \sum_{j; j \neq n} \frac{\partial^2 V_{n,j}(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n^2}$$

Swithing again to the reduced writting, where V_{ij} is only written as a function of the variables of which it actually depends, reads

$$U_{n,m} = \frac{\partial^2 V_{n,m}(\mathbf{R}_n, \mathbf{R}_m)}{\partial \mathbf{R}_n \partial \mathbf{R}_m} \quad \text{if } n \neq m$$

$$U_{n,n} = \sum_{j; j \neq n} \frac{\partial^2 V_{n,j}(\mathbf{R}_n, \mathbf{R}_j)}{\partial \mathbf{R}_n^2}$$

This result is a good point to further specify the potential that is used in the next sections. The potential will only depend on the relative vector between the two positions, i.e. $\mathbf{r}_{ij} = \mathbf{R}_i - \mathbf{R}_j$. Using the derivative

$$\frac{\partial \mathbf{r}_{ij}}{\partial \mathbf{R}_i} = \mathbb{1} = -\frac{\partial \mathbf{r}_{ij}}{\partial \mathbf{R}_j}$$

the derivatives of the U matrix can be written as

$$\begin{aligned} \frac{\partial^2 V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{R}_n \partial \mathbf{R}_m} &= \frac{\partial}{\partial \mathbf{R}_n} \left(\frac{\partial V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{r}_{nm}} \frac{\partial \mathbf{r}_{nm}}{\partial \mathbf{R}_m} \right) \\ &= \frac{\partial^2 V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{r}_{nm}^2} \frac{\partial \mathbf{r}_{nm}}{\partial \mathbf{R}_m} \frac{\partial \mathbf{r}_{nm}}{\partial \mathbf{R}_n} + \frac{\partial V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{r}_{nm}} \frac{\partial^2 \mathbf{r}_{nm}}{\partial \mathbf{R}_n \partial \mathbf{R}_m} \\ &= -\frac{\partial^2 V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{r}_{nm}^2} \end{aligned}$$

and

$$\begin{aligned} \frac{\partial^2 V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{R}_m^2} &= \frac{\partial^2 V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{r}_{nm}^2} \frac{\partial \mathbf{r}_{nm}}{\partial \mathbf{R}_m} \frac{\partial \mathbf{r}_{nm}}{\partial \mathbf{R}_m} \\ &= \frac{\partial^2 V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{r}_{nm}^2} \end{aligned}$$

Thus

$$U_{n,m} = -\frac{\partial^2 V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{r}_{nm}^2} \quad \text{if } n \neq m$$

$$U_{n,n} = \sum_{j; j \neq n} \frac{\partial^2 V_{n,j}(\mathbf{r}_{nj})}{\partial \mathbf{r}_{nj}^2}$$

At this point I can eliminate the linear term of the Hamiltonian by introducing the trap potential

$$H_0(\{\delta \mathbf{R}_l\}) = \sum_n \frac{1}{2} m \omega^2 \delta \mathbf{R}_n^2$$

and redefining the trap centers by a equal shift of all positions $\{\delta \mathbf{R}_l\} = \{\delta \tilde{\mathbf{R}}_l + \mathbf{\Delta}\}$. Neglection all constant terms the Hamiltonian up to second order reads

$$\begin{aligned} H(\{\delta \mathbf{R}_l\}) &= \sum_n \frac{1}{2} m \omega^2 (\delta \tilde{\mathbf{R}}_n + \mathbf{\Delta}_n)^2 + \sum_n \frac{\partial H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n} (\delta \tilde{\mathbf{R}}_n + \mathbf{\Delta}_n) + \sum_{n,m} (\delta \tilde{\mathbf{R}}_n + \mathbf{\Delta}_n) U_{n,m} (\delta \tilde{\mathbf{R}}_m + \mathbf{\Delta}_m) \\ &= \sum_n \frac{1}{2} m \omega^2 \delta \tilde{\mathbf{R}}_n^2 + \sum_{n,m} \delta \tilde{\mathbf{R}}_n U_{n,m} \delta \tilde{\mathbf{R}}_m + \sum_n \left(\frac{\partial H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n} + \left(2 \sum_m \delta \tilde{\mathbf{R}}_m U_{m,n} + m \omega^2 \right) \mathbf{\Delta}_n \right) \delta \tilde{\mathbf{R}}_n, \end{aligned}$$

where it was used that U is symmetric. The linear term vanishes, if Δ_n is chosen such that

$$\Delta_n = - \left(2 \sum_m \delta \tilde{\mathbf{R}}_m U_{m,n} + m\omega^2 \right)^{-1} \frac{\partial H_I(\{\mathbf{R}_l\})}{\partial \mathbf{R}_n}$$

Thus

$$H(\{\delta \tilde{\mathbf{R}}_l\}) = \sum_n \frac{1}{2} m\omega^2 \delta \tilde{\mathbf{R}}_n^2 + \sum_{n,m} \delta \tilde{\mathbf{R}}_n U_{n,m} \delta \tilde{\mathbf{R}}_m$$

To describe the deviation from the center in a quantum mechanical way, the displacements are now quantized as phonons using the quadrature operators

$$\delta \tilde{\mathbf{R}}_n = \frac{1}{\sqrt{m\omega}} (a_n + a_n^\dagger)$$

If we exploit that the interactions realized in U are much smaller compared to the trap frequency, i.e. $U_{n,m} \ll m\omega^2$, the creation and annihilation of phonons is suppressed and the dynamics only involves phonon hopping, thus combinations aa or $a^\dagger a^\dagger$ are discarded.

$$\begin{aligned} H &= \sum_n \omega a_n^\dagger a_n + 2 \sum_{n,m} \frac{U_{n,m}}{m\omega} a_n^\dagger a_m \\ &=: \sum_n \omega a_n^\dagger a_n + \sum_{n,m} \tilde{U}_{n,m} a_n^\dagger a_m \end{aligned}$$

Aligned Dipole-Dipole Interactions

In this section the derivative of the dipole interaction will be performed in the case where all dipoles point into the same direction. The subscripts labeling the different positions are dropped in order to make the notation plainer.

$$V_{dd}(\mathbf{r}) = \frac{1}{r^3} - 3 \frac{(\mathbf{r} \cdot \hat{\mathbf{d}})^2}{r^5},$$

with $\mathbf{r} = \mathbf{R}_i - \mathbf{R}_j$ being the distance between any two sites i and j ($i \neq j$). The first derivative with respect to component α of $\mathbf{r}$, $\alpha \in \{x, y, z\}$ reads

$$\frac{\partial V_{dd}}{\partial r_\alpha} = -\frac{3r_\alpha}{r^5} + \frac{15r_\alpha(\mathbf{r} \cdot \hat{\mathbf{d}})^2}{r^7} - \frac{6(\mathbf{r} \cdot \hat{\mathbf{d}})d_\alpha}{r^5},$$

the second derivative reads consequently

$$\begin{aligned} \frac{\partial^2 V_{dd}}{\partial r_\alpha \partial r_\beta} &= -3 \frac{\delta_{\alpha\beta}}{r^5} + 15 \frac{r_\alpha r_\beta}{r^7} + 15 \frac{\delta_{\alpha\beta}(\mathbf{r} \cdot \hat{\mathbf{d}})^2}{r^7} - 15 \cdot 7 \frac{r_\alpha r_\beta (\mathbf{r} \cdot \hat{\mathbf{d}})^2}{r^9} \\ &\quad + 30 \frac{d_\alpha r_\beta + d_\beta r_\alpha}{r^7} (\mathbf{r} \cdot \hat{\mathbf{d}}) - 6 \frac{d_\alpha d_\beta}{r^5} \\ &= \delta_{\alpha\beta} \left(-3 \frac{1}{r^5} + 15 \frac{(\mathbf{r} \cdot \hat{\mathbf{d}})^2}{r^7} \right) - 6 \frac{d_\alpha d_\beta}{r^5} + 30 \frac{d_\alpha r_\beta + d_\beta r_\alpha}{r^7} (\mathbf{r} \cdot \hat{\mathbf{d}}) \\ &\quad + 15 r_\alpha r_\beta \left(\frac{1}{r^7} - 7 \frac{(\mathbf{r} \cdot \hat{\mathbf{d}})^2}{r^9} \right). \end{aligned}$$

General Dipole-Dipole Interactions

In the case where not all dipole moments are aligned, the properties of the interaction get extended to

$$V_{dd}(\mathbf{r}) = \frac{\hat{\mathbf{d}}_1 \cdot \hat{\mathbf{d}}_2}{r^3} - 3 \frac{(\mathbf{r} \cdot \hat{\mathbf{d}}_1)(\mathbf{r} \cdot \hat{\mathbf{d}}_2)}{r^5}.$$

The first derivative reads

$$\frac{\partial V_{dd}}{\partial r_\alpha} = -\frac{3(\hat{\mathbf{d}}_1 \cdot \hat{\mathbf{d}}_2) r_\alpha}{r^5} + \frac{15 r_\alpha (\mathbf{r} \cdot \hat{\mathbf{d}}_1)(\mathbf{r} \cdot \hat{\mathbf{d}}_2)}{r^7} - \frac{3 \left((\mathbf{r} \cdot \hat{\mathbf{d}}_1) d_{2,\alpha} + (\mathbf{r} \cdot \hat{\mathbf{d}}_2) d_{1,\alpha} \right)}{r^5},$$

the second derivative reads consequently

$$\begin{aligned} \frac{\partial^2 V_{dd}}{\partial r_\alpha \partial r_\beta} &= -3 \frac{\delta_{\alpha\beta} (\hat{\mathbf{d}}_1 \cdot \hat{\mathbf{d}}_2)}{r^5} + 15 \frac{r_\alpha r_\beta (\hat{\mathbf{d}}_1 \cdot \hat{\mathbf{d}}_2)}{r^7} + 15 \frac{\delta_{\alpha\beta} (\mathbf{r} \cdot \hat{\mathbf{d}}_1)(\mathbf{r} \cdot \hat{\mathbf{d}}_2)}{r^7} - 15 \cdot 7 \frac{r_\alpha r_\beta (\mathbf{r} \cdot \hat{\mathbf{d}}_1)(\mathbf{r} \cdot \hat{\mathbf{d}}_2)}{r^9} \\ &\quad + 15 \left[\frac{d_{1,\alpha} r_\beta + d_{1,\beta} r_\alpha}{r^7} (\mathbf{r} \cdot \hat{\mathbf{d}}_2) + \frac{d_{2,\alpha} r_\beta + d_{2,\beta} r_\alpha}{r^7} (\mathbf{r} \cdot \hat{\mathbf{d}}_1) \right] - 3 \frac{d_{1,\alpha} d_{2,\beta} + d_{2,\alpha} d_{1,\beta}}{r^5} \\ &= \delta_{\alpha\beta} \left(-3 \frac{\hat{\mathbf{d}}_1 \cdot \hat{\mathbf{d}}_2}{r^5} + 15 \frac{(\mathbf{r} \cdot \hat{\mathbf{d}}_1)(\mathbf{r} \cdot \hat{\mathbf{d}}_2)}{r^7} \right) - 3 \frac{d_{1,\alpha} d_{2,\beta} + d_{2,\alpha} d_{1,\beta}}{r^5} \\ &\quad + 15 \left[\frac{d_{1,\alpha} r_\beta + d_{1,\beta} r_\alpha}{r^7} (\mathbf{r} \cdot \hat{\mathbf{d}}_2) + \frac{d_{2,\alpha} r_\beta + d_{2,\beta} r_\alpha}{r^7} (\mathbf{r} \cdot \hat{\mathbf{d}}_1) \right] \\ &\quad + 15 r_\alpha r_\beta \left(\frac{\hat{\mathbf{d}}_1 \cdot \hat{\mathbf{d}}_2}{r^7} - 7 \frac{(\mathbf{r} \cdot \hat{\mathbf{d}}_1)(\mathbf{r} \cdot \hat{\mathbf{d}}_2)}{r^9} \right). \end{aligned}$$

For the the Van der Waals interaction the potential has to be squared. Therefore the second derivative reads

$$\begin{aligned} \frac{\partial V_{dd}^2}{\partial r_\alpha} &= 2 V_{dd} \frac{\partial V_{dd}}{\partial r_\alpha} \\ \frac{\partial^2 V_{dd}^2}{\partial r_\alpha \partial r_\beta} &= 2 \frac{\partial V_{dd}}{\partial r_\alpha} \frac{\partial V_{dd}}{\partial r_\beta} + 2 V_{dd} \frac{\partial^2 V_{dd}}{\partial r_\alpha \partial r_\beta} \end{aligned}$$

Winding Number

A chiral Hamiltonian is defined through the property, that it anticommutes with a unitary hermitian operator Γ

$$\Gamma H \Gamma = -H \quad \text{with} \quad \Gamma^2 = 1. \quad (0.1)$$

From this property follows that the Hamiltonian can be written in block off-diagonal form

$$H = \begin{pmatrix} 0 & h \\ h^\dagger & 0 \end{pmatrix}, \quad (0.2)$$

where h is, in general, a non-Hermitian matrix. This structure also applies for the matrix in momentum space and allows the definition of a winding number, which characterizes the topological properties of the system. If the Hamiltonian is 2×2 dimensional, the winding number can be defined through the decomposition of H in terms of the Pauli matrices

$$H = \mathbf{n} \cdot \boldsymbol{\sigma}, \quad (0.3)$$

where $\mathbf{n} = (n_x, n_y, 0)$ is a vector in the xy -plane. The winding number is then given by the winding of the vector $\mathbf{n}$ around the origin when traversing the Brillouin zone. Identifying the xy -plane of $\mathbf{n}$ with the complex plane, which defines a path $\mathbf{n}(k) \rightarrow \gamma(k)$, the winding number can be calculated through

$$w = \frac{1}{2\pi i} \oint_{\gamma_k} \frac{1}{z} dz = \frac{1}{2\pi i} \int_{\text{BZ}} dk \frac{1}{\gamma(k)} \frac{d\gamma(k)}{dk}.$$

In terms of the Hamiltonian above this reads

$$w = \frac{1}{2\pi i} \int_{\text{BZ}} dk \frac{h'(k)}{h(k)}.$$

Generalizing the winding number to higher dimensional Hamiltonians is possible through the use of the block off-diagonal form. The winding number then reads [1, 2]

$$w = \frac{1}{2\pi i} \int_{\text{BZ}} dk \text{Tr} [h^{-1}(k) \partial_k h(k)].$$

Diagonalizing h as $h = S\Lambda S^{-1}$ with the diagonal matrix $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots)$ and the invertible matrix S , the trace can be rewritten as

$$\begin{aligned} \text{Tr} [h^{-1} \partial_k h] &= \text{Tr} [S\Lambda^{-1} S^{-1} \partial_k (S\Lambda S^{-1})] \\ &= \text{Tr} [\Lambda^{-1} \partial_k \Lambda + S^{-1} \partial_k S + (\partial_k S^{-1}) S] \\ \text{Tr} [S^{-1} \partial_k S + S \partial_k S^{-1}] &= \text{Tr} [\partial_k (SS^{-1})] = 0, \\ \Rightarrow \text{Tr} [h^{-1} \partial_k h] &= \text{Tr} [\Lambda^{-1} \partial_k \Lambda] = \sum_i \frac{\partial_k \lambda_i}{\lambda_i} = \partial_k \left(\sum_i \ln(\lambda_i) \right) = \partial_k \ln(\det[h]). \end{aligned}$$

Another useful definition of the winding number is through the so-called Q -matrix, which is defined through the projectors onto the eigenstates of the Hamiltonian. As the Hamiltonian is chiral, the eigenvalues come in pairs $\pm E$. We introduce a labeling of the eigenvalues, such that $E_j = -E_{-j}$ with $j = 1, 2, \dots, N$ and N being half the dimension of the Hamiltonian. The corresponding eigenstates are labeled accordingly ($|\psi_j\rangle$) and posses the property $\Gamma |\psi_j\rangle = |\psi_{-j}\rangle$. The Q -matrix is defined as

$$Q = \sum_{j=1}^N (|\psi_j\rangle \langle \psi_j| - |\psi_{-j}\rangle \langle \psi_{-j}|).$$

As a consequence of the chiral symmetry, the Q -matrix anticommutes with Γ and can therefore be written in block off-diagonal form in the basis, where Γ is diagonal, which is also the case in which H is block off-diagonal.

$$\begin{aligned} \Gamma Q \Gamma &= \sum_{j=1}^N (|\psi_{-j}\rangle \langle \psi_{-j}| - |\psi_j\rangle \langle \psi_j|) = -Q \\ \Rightarrow Q &= \begin{pmatrix} 0 & q \\ q^\dagger & 0 \end{pmatrix} \end{aligned}$$

The fact that the winding number of the q -matrix is the same as that of h can be shown with the help of the following two properties.

$$\begin{aligned} Q^2 = 1 &\Rightarrow qq^\dagger = \mathbb{1}, \quad q^\dagger q = \mathbb{1} \\ QHQ = H &\Rightarrow qh^\dagger q = h^\dagger \end{aligned}$$

From the second property follows that $q^\dagger h = h^\dagger q$. Thus $h^\dagger q$ is a hermitian matrix and therefore has no winding number, since the eigenvalues are real. Hence

$$\begin{aligned} 0 &= \text{Tr} [q^\dagger (h^\dagger)^{-1} \partial_k (h^\dagger q)] \\ &= \text{Tr} [q q^\dagger (h^\dagger)^{-1} \partial_k h^\dagger + h^\dagger (h^\dagger)^{-1} q^\dagger \partial_k q] \\ \Rightarrow \text{Tr} [q^{-1} \partial_k q] &= -\text{Tr} [(h^\dagger)^{-1} \partial_k h^\dagger] \\ &= \text{Tr} [h^{-1} \partial_k h] \end{aligned}$$

There is another useful property, for which we consider the basis in which $\Gamma = \sigma_z \otimes \mathbb{1}_N$. This is the same diagonal basis as previously used. In this basis we can represent all eigenvalues of H as a stack of to N -component vectors. The eigenvector for negative energies is written as

$$|\psi_i^-\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} |h_{1i}\rangle \\ |h_{2i}\rangle \end{pmatrix} \quad i \in \{1, \dots, N\}$$

Because of the chirality of the Hamiltonian the eigenvector with positive energy is already determined by this choice and reads

$$|\psi_i^+\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} |h_{1i}\rangle \\ -|h_{2i}\rangle \end{pmatrix}$$

The matrix Q is represented in the following form

$$Q = \sum_i \begin{pmatrix} 0 & |h_{1i}\rangle \langle h_{2i}| \\ |h_{2i}\rangle \langle h_{1i}| & 0 \end{pmatrix} \\ \Rightarrow q = \sum_i |h_{1i}\rangle \langle h_{2i}|$$

If we now compute the kernel of the winding number with respect to q we get

$$\begin{aligned} \text{Tr}[q^\dagger \partial_k q] &= \text{Tr} \left[\sum_{i,j} |h_{2i}\rangle \langle h_{1i}| \partial_k (|h_{1j}\rangle \langle h_{2j}|) \right] \\ &= \sum_{i,j} \text{Tr} [|h_{2i}\rangle \langle h_{1i}| (\partial_k |h_{1j}\rangle) \langle h_{2j}|] + \sum_{i,j} \text{Tr} [|h_{2i}\rangle \langle h_{1i}| |h_{1j}\rangle (\partial_k \langle h_{2j}|)] \\ &= \sum_i \langle h_{1i}| (\partial_k |h_{1i}\rangle) + \sum_i (\partial_k \langle h_{2i}|) |h_{2i}\rangle \end{aligned}$$

When computing the winding number the second term contributes with a minus sign compared to its complex conjugated self, i.e.

$$\int_{\text{BZ}} dk (\partial_k \langle h_{2i}|) |h_{2i}\rangle = - \int_{\text{BZ}} dk \langle h_{2i}| (\partial_k |h_{2i}\rangle)$$

Thus the winding number reads

$$\begin{aligned} w &= \frac{1}{2\pi i} \int_{\text{BZ}} dk \text{Tr}[q^\dagger \partial_k q] \\ &= \frac{1}{2\pi i} \sum_i \int_{\text{BZ}} dk (\langle h_{1i}| \partial_k |h_{1i}\rangle - \langle h_{2i}| \partial_k |h_{2i}\rangle) \end{aligned}$$

This can be expressed with respect to the full eigenvectors of the Hamiltonian via

$$\begin{aligned} w &= \frac{1}{\pi i} \sum_i \int_{\text{BZ}} dk \langle \Gamma \psi_i^- | \partial_k | \psi_i^- \rangle \\ &= \frac{1}{\pi i} \sum_i \int_{\text{BZ}} dk \langle \psi_i^+ | \partial_k | \psi_i^- \rangle \\ &= \frac{1}{2\pi i} \sum_i \int_{\text{BZ}} dk (\langle h_{1i}| \partial_k |h_{1i}\rangle - \langle h_{2i}| \partial_k |h_{2i}\rangle) \end{aligned}$$

The quantity $\sum_i \langle \Gamma \psi_i^- | \partial_k | \psi_i^- \rangle$ can be related to a polarisation [3].

1D analysis of topological states

In order to have a chiral Hamiltonian I will choose the orientation of the dipoles in a way, such that the chirality remains. This will be done for each lattice configuration separately, i.e. the dipoles depend on the lattice geometry.

In order to make the achievement of chirality more feasible I only consider nearest neighbour interactions. When talking of chirality the chirality of the matrix U , defined above, is in mind. Recall that

$$U_{n,m} = -\frac{\partial^2 V_{n,m}(\mathbf{r}_{nm})}{\partial \mathbf{r}_{nm}^2} \quad \text{if } n \neq m \\ U_{n,n} = \sum_{j; j \neq n} \frac{\partial^2 V_{n,j}(\mathbf{r}_{nj})}{\partial \mathbf{r}_{nj}^2}$$

Chirality implies that all couplings between sites on the same sublattice vanish. The nearest neighbour restriction already assures that couplings between different sites of the same sublattice vanish. Left is thus only the onsite potential due to the Taylor expansion of the dipole-dipole potential. The aim is now to choose the orientation of the dipoles in such a way that the sum in $U_{n,n}$ is equal to zero.

Be aware that so far I didn't write explicitly the dependence of U from the direction of atom movement x, y . Hence $U_{n,m}, U_{n,n}$ are themselves two times two matrixes $(U_{n,m}^{\alpha,\beta})_{\alpha,\beta \in \{x,y\}}$ and the Hamiltonian expressed explicitly with respect to the x, y -phonon modes reads

$$H = \sum_{n,\alpha} \omega a_{n,\alpha}^\dagger a_{n,\alpha} + \frac{2}{m\omega} \sum_{n,m,\alpha,\beta} U_{n,m}^{\alpha,\beta} a_{n,\alpha}^\dagger a_{m,\beta}$$

The terms with $a_{n,\alpha}^\dagger a_{n,\alpha}$ can thus be eliminated by redefining the trap frequency.

$$\omega_\alpha = \omega + \frac{2}{m\omega} U_{n,n}^{\alpha,\alpha}$$

Now only the terms coupling the two direction of phonon modes on the same side have to be tuned to zero. Because of the nearest neighbour nature of the consideration only to distances play a role, the intracell distance $\mathbf{r}_I$ and the intercell distance $\mathbf{r}_O$.

$$U_{n,n}^{x,y} = \frac{\partial^2 V(\mathbf{r}_I)}{\partial r_{I,x} \partial r_{I,y}} + \frac{\partial^2 V(\mathbf{r}_O)}{\partial r_{O,x} \partial r_{O,y}}$$

The dipoles are then chosen in such a way that this sum vanishes.

Aligned dipoles in plane

I consider a geometry with two sublattices repeating itself in the x -axis and separated by the distance h from each other into the y -axis. Different configurations of this geometry are realized by moving the sublattices against each other in the x -direction. I consider an even amount of lattice, with N atoms in each sublattice. The n th atom of each sublattice A and B takes the position.

$$\begin{aligned} \mathbf{r}_n^A &= n \cdot a \mathbf{e}_x \\ \mathbf{r}_n^B &= n \cdot a \mathbf{e}_x + b \mathbf{e}_x + h \mathbf{e}_y, \end{aligned}$$

where a is the lattice constant, and the parameters b and h can be chosen to realize different lattice configurations. b and h have to be chosen in a way, such that the nearest neighbour restriction makes sense. For any of these geometries there exists a unique orientation of the dipoles which makes the Hamiltonian with periodic boundary conditions chiral.










Figure 1: The diagram was created with the parameters $a = 3$, $h = 1.1$. Right to the red dashed line topological phonon states can be found and the winding number is 2.




phonon direction x

phonon direction y












Figure 2: The diagram was created with the parameters $a = 5$, $h = 1.1$. Between the red and the olive dashed line, i.e. between b -values of 2.25-2.73 the winding number is 1, whereas right to the olive dashed line the winding number is 2.




open boundary




periodic boundary



phonon direction x





phonon direction y



$b = 1.00; E = 1.4029$







Bibliography

- [1] M. Maffei, A. Dauphin, F. Cardano, M. Lewenstein, and P. Massignan, “Topological characterization of chiral models through their long time dynamics”, [New Journal of Physics](#) **20**, 013023 (2018).
- [2] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, “Classification of topological insulators and superconductors in three spatial dimensions”, [Physical Review B](#) **78**, 195125 (2008).
- [3] I. Mondragon-Shem, T. L. Hughes, J. Song, and E. Prodan, “Topological criticality in the chiral-symmetric AIII class at strong disorder”, [Physical Review Letters](#) **113**, 046802 (2014).