


Test problems

Branin (heteroscedastic, 5 levels)

$$f(x_1, \ell) = \left(x_2(\ell) - \frac{5.1}{4\pi^2} x_1^2 + \frac{5}{\pi} x_1 - 6 \right)^2 + 10 \left(1 - \frac{1}{8\pi} \right) \cos x_1 + 10, \quad x_1 \in [-5, 10], \quad x_2(\ell) \in \{15, 2, 8, 0, 10\}$$



$$\sigma(x_1, \ell) = \left[0.5 + 0.15(x_1 + 5) + \frac{2.5}{1 + e^{-1.2(x_1 - 2)}} \right] m(\ell), \quad m(\ell) \in \{0.5, 1.0, 1.8, 2.5, 3.5\}$$



Six-hump camel (5 levels: two pairs + bridge)

$$f(x_1, \ell) = \left(4 - 2.1x_1^2 + \frac{x_1^4}{3} \right) x_1^2 + x_1 x_2(\ell) + (-4 + 4x_2(\ell)^2) x_2(\ell)^2, \quad x_1 \in [-2, 2], \quad x_2(\ell) \in \{-1.0, -0.85, 0.0, 0.85, 1.0\}$$

$$\sigma(x_1, \ell) = 0.05 e^{(0.4x_1)^2} m(\ell), \quad m(\ell) \in \{2.0, 3.5, 1.5, 5.0, 2.5\}$$



Griewank 6-D (level-shifted pairs)

$$f(\mathbf{x}_q, \ell) = \sum_{i=1}^6 \frac{x s_i^2}{4000} - \prod_{i=1}^6 \cos\left(\frac{x s_i}{\sqrt{i}}\right) + 1 + b(\ell), \quad x s = [\mathbf{x}_q, v(\ell)], \quad \mathbf{x}_q \in [-3, 3]^5$$

$$v(\ell) \in \{0.5, 1.3, 2.7, 3.1\}, \quad b(\ell) \in \{0, 0.15, 0.6, 0.75\}, \quad \sigma(\mathbf{x}_q, \ell) = 0.02\sqrt{1 + 0.1\|\mathbf{x}_q\|^2/5} m(\ell), \quad m(\ell) \in \{1.5, 1.0, 3.0, 2.0\}$$





Table 1: Mean-over-levels metrics for every problem $\times$ budget $\times$ model (seed 1). Bold on green = best model in each (problem, budget, n_{rep}) block per metric (MAE/RRMSE/IS: lower; coverage: closest to 0.95).

Problem	n	Model	$n_{\text{rep}} = 10$			
			MAE $\downarrow$	RRMSE $\downarrow$	IS $\downarrow$	Cov. $\rightarrow .95$
Branin	25	Separate GPs	11.24	0.4550	125.4	0.7340
		Categorical GP	4.757	0.1821	41.85	0.9340
		Standard LVGP	3.932	0.1763	63.11	0.3710
		H-LVGP	2.554	0.0835	28.91	1.000
	40	Separate GPs	4.295	0.2002	41.91	0.9360
		Categorical GP	1.879	0.0652	14.08	1.000
		Standard LVGP	2.706	0.1087	55.61	0.4250
		H-LVGP	1.350	0.0546	30.60	1.000
Six-hump camel	25	Separate GPs	0.2291	0.3243	2.127	0.9550
		Categorical GP	0.1606	0.2654	1.809	0.9260
		Standard LVGP	0.1704	0.2768	3.481	0.4740
		H-LVGP	0.1801	0.2942	2.264	0.9070
	40	Separate GPs	0.2006	0.2924	2.332	0.8850
		Categorical GP	0.1008	0.1432	1.001	0.9210
		Standard LVGP	0.1618	0.2567	3.070	0.5180
		H-LVGP	0.0743	0.1101	0.7647	0.9980
Griewank 6-D	40	Separate GPs	0.0669	0.9651	0.7438	0.8606
		Categorical GP	0.0647	0.9497	0.8928	0.8145
		Standard LVGP	0.1003	1.266	1.821	0.6140
		H-LVGP	0.0819	1.148	0.9839	0.7612
	80	Separate GPs	0.0703	1.039	0.7977	0.8450
		Categorical GP	0.0587	0.9623	0.7190	0.8899
		Standard LVGP	0.0975	1.215	0.8763	0.8708
		H-LVGP	0.0757	1.035	1.006	0.7766
	160	Separate GPs	0.0553	0.8212	0.6258	0.8604
		Categorical GP	0.0445	0.6732	0.5228	0.9343
		Standard LVGP	0.0666	0.8666	0.5618	0.8921
		H-LVGP	0.0695	0.9592	1.232	0.6562

Table 2: Mean-over-levels metrics for every problem $\times$ budget $\times$ model (without the Categorical GP) (seed 1). Bold on green = best model in each (problem, budget, n_{rep}) block per metric (MAE/RRMSE/IS: lower; coverage: closest to 0.95).

Problem	n	Model	$n_{\text{rep}} = 10$			
			MAE $\downarrow$	RRMSE $\downarrow$	IS $\downarrow$	Cov. $\rightarrow .95$
Branin	25	Separate GPs	11.24	0.4550	125.4	0.7340
		Standard LVGP	3.932	0.1763	63.11	0.3710
		H-LVGP	2.554	0.0835	28.91	1.000
	40	Separate GPs	4.295	0.2002	41.91	0.9360
		Standard LVGP	2.706	0.1087	55.61	0.4250
		H-LVGP	1.350	0.0546	30.60	1.000
Six-hump camel	25	Separate GPs	0.2291	0.3243	2.127	0.9550
		Standard LVGP	0.1704	0.2768	3.481	0.4740
		H-LVGP	0.1801	0.2942	2.264	0.9070
	40	Separate GPs	0.2006	0.2924	2.332	0.8850
		Standard LVGP	0.1618	0.2567	3.070	0.5180
		H-LVGP	0.0743	0.1101	0.7647	0.9980
Griewank 6-D	40	Separate GPs	0.0669	0.9651	0.7438	0.8606
		Standard LVGP	0.1003	1.266	1.821	0.6140
		H-LVGP	0.0819	1.148	0.9839	0.7612
	80	Separate GPs	0.0703	1.039	0.7977	0.8450
		Standard LVGP	0.0975	1.215	0.8763	0.8708
		H-LVGP	0.0757	1.035	1.006	0.7766
	160	Separate GPs	0.0553	0.8212	0.6258	0.8604
		Standard LVGP	0.0666	0.8666	0.5618	0.8921
		H-LVGP	0.0695	0.9592	1.232	0.6562

Table 3: 30-seed replication (fixed design, fresh noise replicates per seed): mean $\pm$ std over seeds. Bold = best mean per (problem, budget) block (coverage: closest to 0.95).

	Problem	n	Model	MAE	RRMSE	IS	Coverage
Branin		25	Separate GPs	10.908 ± 0.466	0.434 ± 0.022	113.7 ± 11.604	0.775 ± 0.028
			Categorical GP	4.529 ± 0.241	0.159 ± 0.011	38.874 ± 2.018	0.945 ± 0.010
			Standard LVGP	4.419 ± 1.063	0.212 ± 0.051	74.800 ± 30.689	0.442 ± 0.143
			H-LVGP	2.520 ± 0.723	0.096 ± 0.037	37.281 ± 20.109	0.996 ± 0.016
		40	Separate GPs	4.454 ± 0.361	0.208 ± 0.017	42.877 ± 1.784	0.931 ± 0.011
			Categorical GP	1.585 ± 0.268	0.054 ± 0.012	14.864 ± 0.896	0.996 ± 0.010
			Standard LVGP	3.109 ± 0.705	0.122 ± 0.030	62.823 ± 20.089	0.410 ± 0.145
			H-LVGP	1.722 ± 0.425	0.064 ± 0.019	35.613 ± 10.826	0.998 ± 0.008
Six-hump camel		25	Separate GPs	0.231 ± 0.009	0.319 ± 0.007	2.072 ± 0.061	0.951 ± 0.008
			Categorical GP	0.168 ± 0.009	0.262 ± 0.010	1.778 ± 0.136	0.923 ± 0.012
			Standard LVGP	0.152 ± 0.018	0.242 ± 0.029	3.405 ± 0.521	0.417 ± 0.097
			H-LVGP	0.178 ± 0.012	0.287 ± 0.016	2.153 ± 0.231	0.905 ± 0.017
		40	Separate GPs	0.196 ± 0.008	0.295 ± 0.006	2.313 ± 0.111	0.888 ± 0.015
			Categorical GP	0.114 ± 0.010	0.160 ± 0.015	1.127 ± 0.164	0.918 ± 0.013
			Standard LVGP	0.145 ± 0.034	0.212 ± 0.066	3.240 ± 1.092	0.324 ± 0.106
			H-LVGP	0.143 ± 0.024	0.226 ± 0.042	1.147 ± 0.318	0.953 ± 0.029

Table 4: Branin — per-level metrics, $n_{\text{rep}} = 10$ (seed 1; MAE/RRMSE/IS lower is better, coverage target 0.95). Bold = best model in the row; the green highlight marks the winners of the *Mean* row.

n	Level	Separate GPs				Categorical GP				Standard LVGP				H-LVGP			
		MAE $\downarrow$	RRMSE $\downarrow$	IS $\downarrow$	Cov. $\rightarrow$.95	MAE $\downarrow$	RRMSE $\downarrow$	IS $\downarrow$	Cov. $\rightarrow$.95	MAE $\downarrow$	RRMSE $\downarrow$	IS $\downarrow$	Cov. $\rightarrow$.95	MAE $\downarrow$	RRMSE $\downarrow$	IS $\downarrow$	Cov. $\rightarrow$.95
25	$x_2 = 15$	5.902	0.1174	35.77	0.6200	4.435	0.0967	20.37	0.9350	6.136	0.1679	96.80	0.4100	3.339	0.0917	32.68	1.000
	$x_2 = 2$	14.17	0.4118	162.3	0.7950	4.714	0.1723	49.29	0.9150	2.032	0.0488	31.46	0.4300	2.327	0.0531	19.96	1.000
	$x_2 = 8$	7.824	0.8427	130.3	0.8600	3.366	0.3004	29.88	1.000	3.853	0.3730	41.36	0.2900	1.691	0.1164	30.41	1.000
	$x_2 = 0$	16.19	0.2789	124.3	0.6850	6.556	0.1302	72.62	0.8200	3.392	0.0622	79.81	0.3450	3.565	0.0615	28.74	1.000
	$x_2 = 10$	12.14	0.6243	174.6	0.7100	4.714	0.2111	37.07	1.000	4.247	0.2297	66.10	0.3800	1.847	0.0949	32.76	1.000
	<i>Mean</i>	11.24	0.4550	125.4	0.7340	4.757	0.1821	41.85	0.9340	3.932	0.1763	63.11	0.3710	2.554	0.0835	28.91	1.000
40	$x_2 = 15$	0.5195	0.0128	8.882	1.000	0.8563	0.0214	7.364	1.000	0.7171	0.0222	6.774	0.8400	1.016	0.0368	30.29	1.000
	$x_2 = 2$	6.850	0.2434	85.51	0.9100	1.591	0.0525	10.31	1.000	3.729	0.1288	92.07	0.1650	0.7563	0.0280	25.59	1.000
	$x_2 = 8$	4.018	0.3915	38.87	0.9050	0.7749	0.0480	13.18	1.000	2.240	0.1564	27.19	0.4800	0.8975	0.0521	27.75	1.000
	$x_2 = 0$	3.333	0.0713	40.92	0.9000	2.576	0.0453	17.40	1.000	2.888	0.0597	45.91	0.3400	1.096	0.0212	34.81	1.000
	$x_2 = 10$	6.753	0.2818	35.39	0.9650	3.599	0.1588	22.14	1.000	3.958	0.1764	106.1	0.3000	2.984	0.1347	34.58	1.000
	<i>Mean</i>	4.295	0.2002	41.91	0.9360	1.879	0.0652	14.08	1.000	2.706	0.1087	55.61	0.4250	1.350	0.0546	30.60	1.000

Table 5: Branin — per-level metrics (without the Categorical GP), $n_{\text{rep}} = 10$ (seed 1; MAE/RRMSE/IS lower is better, coverage target 0.95). Bold = best model in the row; the green highlight marks the winners of the *Mean* row.

n	Level	Separate GPs				Standard LVGP				H-LVGP			
		MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95
25	$x_2 = 15$	5.902	0.1174	35.77	0.6200	6.136	0.1679	96.80	0.4100	3.339	0.0917	32.68	1.000
	$x_2 = 2$	14.17	0.4118	162.3	0.7950	2.032	0.0488	31.46	0.4300	2.327	0.0531	19.96	1.000
	$x_2 = 8$	7.824	0.8427	130.3	0.8600	3.853	0.3730	41.36	0.2900	1.691	0.1164	30.41	1.000
	$x_2 = 0$	16.19	0.2789	124.3	0.6850	3.392	0.0622	79.81	0.3450	3.565	0.0615	28.74	1.000
	$x_2 = 10$	12.14	0.6243	174.6	0.7100	4.247	0.2297	66.10	0.3800	1.847	0.0949	32.76	1.000
	<i>Mean</i>	11.24	0.4550	125.4	0.7340	3.932	0.1763	63.11	0.3710	2.554	0.0835	28.91	1.000
40	$x_2 = 15$	0.5195	0.0128	8.882	1.000	0.7171	0.0222	6.774	0.8400	1.016	0.0368	30.29	1.000
	$x_2 = 2$	6.850	0.2434	85.51	0.9100	3.729	0.1288	92.07	0.1650	0.7563	0.0280	25.59	1.000
	$x_2 = 8$	4.018	0.3915	38.87	0.9050	2.240	0.1564	27.19	0.4800	0.8975	0.0521	27.75	1.000
	$x_2 = 0$	3.333	0.0713	40.92	0.9000	2.888	0.0597	45.91	0.3400	1.096	0.0212	34.81	1.000
	$x_2 = 10$	6.753	0.2818	35.39	0.9650	3.958	0.1764	106.1	0.3000	2.984	0.1347	34.58	1.000
	<i>Mean</i>	4.295	0.2002	41.91	0.9360	2.706	0.1087	55.61	0.4250	1.350	0.0546	30.60	1.000

Table 6: Six-hump camel — per-level metrics, $n_{\text{rep}} = 10$ (seed 1; MAE/RRMSE/IS lower is better, coverage target 0.95). Bold = best model in the row; the green highlight marks the winners of the *Mean* row.

n	Level	Separate GPs				Categorical GP				Standard LVGP				H-LVGP			
		MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95
25	$x_2 = -1$	0.1576	0.2399	2.147	0.9500	0.1375	0.2240	2.087	0.9250	0.1597	0.2405	2.904	0.5250	0.1513	0.2686	2.655	0.9050
	$x_2 = -0.85$	0.2272	0.3401	3.011	0.9250	0.1391	0.2381	1.936	0.9100	0.1641	0.2773	3.695	0.6600	0.1938	0.2606	2.273	0.8700
	$x_2 = 0$	0.1736	0.3946	1.491	0.9700	0.1434	0.3738	1.718	0.9300	0.1410	0.3669	2.630	0.5000	0.1474	0.4058	2.109	0.9200
	$x_2 = 0.85$	0.3342	0.3871	1.770	0.9800	0.2369	0.2755	1.398	0.9350	0.2241	0.2874	5.190	0.2700	0.2301	0.3005	2.306	0.8900
	$x_2 = 1$	0.2530	0.2595	2.216	0.9500	0.1460	0.2154	1.906	0.9300	0.1632	0.2122	2.988	0.4150	0.1781	0.2352	1.976	0.9500
	<i>Mean</i>	0.2291	0.3243	2.127	0.9550	0.1606	0.2654	1.809	0.9260	0.1704	0.2768	3.481	0.4740	0.1801	0.2942	2.264	0.9070
40	$x_2 = -1$	0.1813	0.3345	3.313	0.8900	0.0908	0.1229	0.8693	0.9150	0.1615	0.2556	3.114	0.3650	0.0650	0.0951	0.6201	1.000
	$x_2 = -0.85$	0.1926	0.3230	2.123	0.9400	0.0960	0.1475	0.9524	0.9600	0.1697	0.2842	3.947	0.4450	0.0711	0.1088	0.9587	1.000
	$x_2 = 0$	0.2048	0.3579	3.210	0.7350	0.1250	0.2128	1.531	0.7900	0.1858	0.3612	3.382	0.5850	0.0754	0.1491	0.5218	0.9900
	$x_2 = 0.85$	0.2345	0.2416	1.292	0.9200	0.0907	0.1141	0.7689	0.9850	0.1486	0.1978	2.761	0.6150	0.0810	0.1011	1.021	1.000
	$x_2 = 1$	0.1897	0.2051	1.722	0.9400	0.1013	0.1188	0.8838	0.9550	0.1433	0.1848	2.144	0.5800	0.0790	0.0963	0.7024	1.000
	<i>Mean</i>	0.2006	0.2924	2.332	0.8850	0.1008	0.1432	1.001	0.9210	0.1618	0.2567	3.070	0.5180	0.0743	0.1101	0.7647	0.9980

Table 7: Six-hump camel — per-level metrics (without the Categorical GP), $n_{\text{rep}} = 10$ (seed 1; MAE/RRMSE/IS lower is better, coverage target 0.95). Bold = best model in the row; the green highlight marks the winners of the *Mean* row.

n	Level	Separate GPs				Standard LVGP				H-LVGP			
		MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95
25	$x_2 = -1$	0.1576	0.2399	2.147	0.9500	0.1597	0.2405	2.904	0.5250	0.1513	0.2686	2.655	0.9050
	$x_2 = -0.85$	0.2272	0.3401	3.011	0.9250	0.1641	0.2773	3.695	0.6600	0.1938	0.2606	2.273	0.8700
	$x_2 = 0$	0.1736	0.3946	1.491	0.9700	0.1410	0.3669	2.630	0.5000	0.1474	0.4058	2.109	0.9200
	$x_2 = 0.85$	0.3342	0.3871	1.770	0.9800	0.2241	0.2874	5.190	0.2700	0.2301	0.3005	2.306	0.8900
	$x_2 = 1$	0.2530	0.2595	2.216	0.9500	0.1632	0.2122	2.988	0.4150	0.1781	0.2352	1.976	0.9500
	<i>Mean</i>	0.2291	0.3243	2.127	0.9550	0.1704	0.2768	3.481	0.4740	0.1801	0.2942	2.264	0.9070
40	$x_2 = -1$	0.1813	0.3345	3.313	0.8900	0.1615	0.2556	3.114	0.3650	0.0650	0.0951	0.6201	1.000
	$x_2 = -0.85$	0.1926	0.3230	2.123	0.9400	0.1697	0.2842	3.947	0.4450	0.0711	0.1088	0.9587	1.000
	$x_2 = 0$	0.2048	0.3579	3.210	0.7350	0.1858	0.3612	3.382	0.5850	0.0754	0.1491	0.5218	0.9900
	$x_2 = 0.85$	0.2345	0.2416	1.292	0.9200	0.1486	0.1978	2.761	0.6150	0.0810	0.1011	1.021	1.000
	$x_2 = 1$	0.1897	0.2051	1.722	0.9400	0.1433	0.1848	2.144	0.5800	0.0790	0.0963	0.7024	1.000
	<i>Mean</i>	0.2006	0.2924	2.332	0.8850	0.1618	0.2567	3.070	0.5180	0.0743	0.1101	0.7647	0.9980

Table 8: Griewank 6-D — per-level metrics, $n_{\text{rep}} = 10$ (seed 1; MAE/RRMSE/IS lower is better, coverage target 0.95). Bold = best model in the row; the green highlight marks the winners of the *Mean* row.

n	Level	Separate GPs				Categorical GP				Standard LVGP				H-LVGP			
		MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95
40	$x_6 = 0.5$	0.0862	0.8571	1.198	0.8154	0.0917	0.8883	1.577	0.7051	0.1465	1.183	2.838	0.4160	0.1047	0.9267	1.456	0.6592
	$x_6 = 1.3$	0.0930	0.9010	0.8643	0.8740	0.0780	0.8692	1.254	0.7402	0.1420	1.379	3.012	0.4863	0.1034	0.9944	1.687	0.5801
	$x_6 = 2.7$	0.0564	1.066	0.5188	0.8926	0.0500	0.9440	0.4207	0.8916	0.0665	1.216	0.9542	0.6934	0.0648	1.188	0.4539	0.8750
	$x_6 = 3.1$	0.0322	1.036	0.3945	0.8604	0.0390	1.097	0.3191	0.9209	0.0462	1.287	0.4803	0.8604	0.0546	1.481	0.3397	0.9307
	<i>Mean</i>	0.0669	0.9651	0.7438	0.8606	0.0647	0.9497	0.8928	0.8145	0.1003	1.266	1.821	0.6140	0.0819	1.148	0.9839	0.7612
80	$x_6 = 0.5$	0.1109	1.035	1.111	0.8867	0.0822	0.8085	1.088	0.8145	0.1585	1.249	1.647	0.7188	0.1140	0.9631	1.756	0.6074
	$x_6 = 1.3$	0.0835	0.8720	0.9051	0.8398	0.0712	0.8004	0.8857	0.8506	0.1148	1.065	1.014	0.8311	0.0913	0.9432	1.439	0.6836
	$x_6 = 2.7$	0.0455	1.006	0.7002	0.8037	0.0422	0.9101	0.4403	0.9453	0.0500	0.8773	0.4351	0.9609	0.0534	1.022	0.5069	0.8877
	$x_6 = 3.1$	0.0413	1.241	0.4745	0.8496	0.0392	1.330	0.4623	0.9492	0.0666	1.669	0.4085	0.9727	0.0439	1.213	0.3215	0.9277
	<i>Mean</i>	0.0703	1.039	0.7977	0.8450	0.0587	0.9623	0.7190	0.8899	0.0975	1.215	0.8763	0.8708	0.0757	1.035	1.006	0.7766
160	$x_6 = 0.5$	0.0745	0.7178	0.7606	0.9229	0.0629	0.6550	0.8499	0.8691	0.0968	0.7871	0.8105	0.8125	0.0978	0.8982	2.038	0.5254
	$x_6 = 1.3$	0.0714	0.7728	0.8774	0.8447	0.0553	0.6459	0.6958	0.8926	0.0807	0.7628	0.6706	0.8721	0.0852	0.8944	1.784	0.5430
	$x_6 = 2.7$	0.0455	0.9164	0.4389	0.9150	0.0311	0.5816	0.2776	0.9854	0.0499	0.8979	0.4515	0.9150	0.0575	1.030	0.7724	0.6738
	$x_6 = 3.1$	0.0300	0.8778	0.4265	0.7588	0.0287	0.8104	0.2681	0.9902	0.0389	1.019	0.3144	0.9688	0.0374	1.014	0.3331	0.8828
	<i>Mean</i>	0.0553	0.8212	0.6258	0.8604	0.0445	0.6732	0.5228	0.9343	0.0666	0.8666	0.5618	0.8921	0.0695	0.9592	1.232	0.6562

Table 9: Griewank 6-D — per-level metrics (without the Categorical GP), $n_{\text{rep}} = 10$ (seed 1; MAE/RRMSE/IS lower is better, coverage target 0.95). Bold = best model in the row; the green highlight marks the winners of the *Mean* row.

n	Level	Separate GPs				Standard LVGP				H-LVGP			
		MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95	MAE ↓	RRMSE ↓	IS ↓	Cov. → .95
40	x6 = 0.5	0.0862	0.8571	1.198	0.8154	0.1465	1.183	2.838	0.4160	0.1047	0.9267	1.456	0.6592
	x6 = 1.3	0.0930	0.9010	0.8643	0.8740	0.1420	1.379	3.012	0.4863	0.1034	0.9944	1.687	0.5801
	x6 = 2.7	0.0564	1.066	0.5188	0.8926	0.0665	1.216	0.9542	0.6934	0.0648	1.188	0.4539	0.8750
	x6 = 3.1	0.0322	1.036	0.3945	0.8604	0.0462	1.287	0.4803	0.8604	0.0546	1.481	0.3397	0.9307
	<i>Mean</i>	0.0669	0.9651	0.7438	0.8606	0.1003	1.266	1.821	0.6140	0.0819	1.148	0.9839	0.7612
80	x6 = 0.5	0.1109	1.035	1.111	0.8867	0.1585	1.249	1.647	0.7188	0.1140	0.9631	1.756	0.6074
	x6 = 1.3	0.0835	0.8720	0.9051	0.8398	0.1148	1.065	1.014	0.8311	0.0913	0.9432	1.439	0.6836
	x6 = 2.7	0.0455	1.006	0.7002	0.8037	0.0500	0.8773	0.4351	0.9609	0.0534	1.022	0.5069	0.8877
	x6 = 3.1	0.0413	1.241	0.4745	0.8496	0.0666	1.669	0.4085	0.9727	0.0439	1.213	0.3215	0.9277
	<i>Mean</i>	0.0703	1.039	0.7977	0.8450	0.0975	1.215	0.8763	0.8708	0.0757	1.035	1.006	0.7766
160	x6 = 0.5	0.0745	0.7178	0.7606	0.9229	0.0968	0.7871	0.8105	0.8125	0.0978	0.8982	2.038	0.5254
	x6 = 1.3	0.0714	0.7728	0.8774	0.8447	0.0807	0.7628	0.6706	0.8721	0.0852	0.8944	1.784	0.5430
	x6 = 2.7	0.0455	0.9164	0.4389	0.9150	0.0499	0.8979	0.4515	0.9150	0.0575	1.030	0.7724	0.6738
	x6 = 3.1	0.0300	0.8778	0.4265	0.7588	0.0389	1.019	0.3144	0.9688	0.0374	1.014	0.3331	0.8828
	<i>Mean</i>	0.0553	0.8212	0.6258	0.8604	0.0666	0.8666	0.5618	0.8921	0.0695	0.9592	1.232	0.6562