


## Stat GR5205 Lecture 2

Jingchen Liu

Department of Statistics  
Columbia University


## Least squares estimator



- ▶ Fitting a straight line  $y = \beta_0 + \beta_1 x$
- ▶ Least squares estimate

$$(\hat{\beta}_0, \hat{\beta}_1) = \arg \min_{\beta_0, \beta_1} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2.$$

## Least squares estimator



- ▶ Fitting a straight line  $y = \beta_0 + \beta_1 x$
- ▶ Least squares estimate

$$(\hat{\beta}_0, \hat{\beta}_1) = \arg \min_{\beta_0, \beta_1} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2.$$

## Least squares estimator for simple linear regression

- ▶ The slope

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- ▶ The intercept

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

- ▶ Slope

$$\hat{\beta}_1 = \rho_{x,y} \frac{s_y}{s_x}.$$

- ▶ The fitted regression line

$$(x - \bar{x}) = \rho_{x,y} \frac{s_y}{s_x} (y - \bar{y})$$

## Least squares estimator for simple linear regression

- ▶ The slope

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- ▶ The intercept

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

- ▶ Slope

$$\hat{\beta}_1 = \rho_{x,y} \frac{s_y}{s_x}.$$

- ▶ The fitted regression line

$$(x - \bar{x}) = \rho_{x,y} \frac{s_y}{s_x} (y - \bar{y})$$

## Least squares estimator for simple linear regression

- ▶ The slope

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- ▶ The intercept

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

- ▶ Slope

$$\hat{\beta}_1 = \rho_{x,y} \frac{s_y}{s_x}.$$

- ▶ The fitted regression line

$$(x - \bar{x}) = \rho_{x,y} \frac{s_y}{s_x} (y - \bar{y})$$

## Least squares estimator for simple linear regression

- ▶ The slope

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- ▶ The intercept

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$


- ▶ Slope

$$\hat{\beta}_1 = \rho_{x,y} \frac{s_y}{s_x}.$$

- ▶ The fitted regression line

$$(x - \bar{x}) = \rho_{x,y} \frac{s_y}{s_x} (y - \bar{y})$$

## On the decomposition of the $y$



- ▶ The fitted values – predictable

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$$

- ▶ The residuals – unpredictable


$$\hat{\varepsilon}_i = y_i - \hat{y}_i$$

- ▶ The decomposition

$$y_i = \hat{y}_i + \hat{\varepsilon}_i$$



## On the decomposition of the $y$



- ▶ The fitted values – predictable

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$$


- ▶ The residuals – unpredictable

$$\hat{\varepsilon}_i = y_i - \hat{y}_i$$

- ▶ The decomposition

$$y_i = \hat{y}_i + \hat{\varepsilon}_i$$

## On the decomposition of the $y$



- ▶ The fitted values – predictable

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$$

- ▶ The residuals – unpredictable

$$\hat{\varepsilon}_i = y_i - \hat{y}_i$$

- ▶ The decomposition

$$y_i = \hat{y}_i + \hat{\varepsilon}_i$$

## On the properties of the fitted regression line

- ▶  $\sum_{i=1}^n \hat{\epsilon}_i = 0$
- ▶  $\sum_{i=1}^n y_i = \sum_{i=1}^n \hat{y}_i$
- ▶  $\sum_{i=1}^n x_i \hat{\epsilon}_i = 0$
- ▶  $\sum_{i=1}^n \hat{y}_i \hat{\epsilon}_i = 0$

## On the properties of the fitted regression line

- ▶  $\sum_{i=1}^n \hat{\epsilon}_i = 0$
- ▶  $\sum_{i=1}^n y_i = \sum_{i=1}^n \hat{y}_i$
- ▶  $\sum_{i=1}^n x_i \hat{\epsilon}_i = 0$
- ▶  $\sum_{i=1}^n \hat{y}_i \hat{\epsilon}_i = 0$

## On the properties of the fitted regression line

- ▶  $\sum_{i=1}^n \hat{\epsilon}_i = 0$
- ▶  $\sum_{i=1}^n y_i = \sum_{i=1}^n \hat{y}_i$
- ▶  $\sum_{i=1}^n x_i \hat{\epsilon}_i = 0$
- ▶  $\sum_{i=1}^n \hat{y}_i \hat{\epsilon}_i = 0$

## On the properties of the fitted regression line

- ▶  $\sum_{i=1}^n \hat{\epsilon}_i = 0$
- ▶  $\sum_{i=1}^n y_i = \sum_{i=1}^n \hat{y}_i$
- ▶  $\sum_{i=1}^n x_i \hat{\epsilon}_i = 0$
- ▶  $\sum_{i=1}^n \hat{y}_i \hat{\epsilon}_i = 0$

# Expectation

- Expectation

$$E(X) = \int xf(x)dx$$

- Linear operator

$$E(aX + bY) = aE(x) + bE(Y)$$

- Long run average

$$\frac{X_1 + \dots + X_n}{n} \rightarrow E(x) \text{ as } n \rightarrow \infty \text{ as most surely}$$

# Expectation

- ▶ Expectation

$$E(X) = \int xf(x)dx$$

- ▶ Linear operator

$$E(aX + bY) = aE(x) + bE(Y)$$

- ▶ Long run average

$$\frac{X_1 + \dots + X_n}{n} \rightarrow E(x) \text{ as } n \rightarrow \infty \text{ as most surely}$$



# Expectation

- ▶ Expectation

$$E(X) = \int xf(x)dx$$

- ▶ Linear operator

$$E(aX + bY) = aE(x) + bE(Y)$$

- ▶ Long run average

$$\frac{X_1 + \dots + X_n}{n} \rightarrow E(x) \text{ as } n \rightarrow \infty \text{ as most surely}$$

# Expectation

- ▶ Variance:  $\text{Var}(X) = E\{X - E(X)\}^2$
- ▶ Covariance:  $\text{Cov}(X, Y) = E\{X - E(X)\}\{Y - E(Y)\}$

# Expectation

- ▶ Variance:  $\text{Var}(X) = E\{X - E(X)\}^2$
- ▶ Covariance:  $\text{Cov}(X, Y) = E\{X - E(X)\}\{Y - E(Y)\}$

## Probability model

- ▶ The model setup

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$

where  $E(\varepsilon_i | x_i) = 0$ .

- ▶ The conditional expectation

$$E(y_i | x_i) = \beta_0 + \beta_1 x_i.$$

- ▶ The term “regression.”

## Probability model

- ▶ The model setup

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$

where  $E(\varepsilon_i | x_i) = 0$ .

- ▶ The conditional expectation

$$E(y_i | x_i) = \beta_0 + \beta_1 x_i.$$

- ▶ The term “regression.”

## Probability model

- ▶ The model setup

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$



where  $E(\varepsilon_i | x_i) = 0$ .

- ▶ The conditional expectation

$$E(y_i | x_i) = \beta_0 + \beta_1 x_i.$$



- ▶ The term “regression.”

## Additional assumption



- ▶ Common variance  $\text{Var}(\varepsilon_i|x_i) = \sigma^2$
- ▶ Quantifying the uncertainty

## Additional assumption



- ▶ Uncorrelated errors  $E(\varepsilon_i \varepsilon_j | x_i, x_j) = 0$
- ▶ Quantifying the uncertainty



## Scope of statistical inference

- ▶ Point estimate (Frequentist distribution)
- ▶ Interval estimate
- ▶ Hypothesis testing

## Scope of statistical inference

- ▶ Point estimate (Frequentist distribution)
- ▶ Interval estimate
- ▶ Hypothesis testing

## Scope of statistical inference

- ▶ Point estimate (Frequentist distribution)
- ▶ Interval estimate
- ▶ Hypothesis testing

## The four assumptions of linear model

- ▶ Linearity
- ▶ Equal variance
- ▶ Independence
- ▶ Normality

## The four assumptions of linear model

- ▶ Linearity
- ▶ Equal variance
- ▶ Independence
- ▶ Normality

## The four assumptions of linear model

- ▶ Linearity
- ▶ Equal variance
- ▶ Independence
- ▶ Normality

## The four assumptions of linear model

- ▶ Linearity
- ▶ Equal variance
- ▶ Independence
- ▶ Normality

## The four assumptions of linear model

- ▶  $y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$
- ▶  $\varepsilon_i$ 's are independently and identically distributed as  $N(0, \sigma^2)$



## The parameters

- ▶ Regression coefficients:  $\beta_0$  and  $\beta_1$
- ▶ Variance:  $\sigma^2$

## The parameters

- ▶ Regression coefficients:  $\beta_0$  and  $\beta_1$
- ▶ Variance:  $\sigma^2$

## The sampling distribution

- ▶ Sampling distribution
- ▶ On the sampling distribution of  $(\hat{\beta}_0, \hat{\beta}_1)$ .

## Additional assumption




Figure:  $y = x + N(0, 0.04)$ ,  $n = 50$ , data 1

## Additional assumption




Figure:  $y = x + N(0, 0.04)$ ,  $n = 50$ , data 2

## Additional assumption




Figure:  $y = x + N(0, 0.04)$ ,  $n = 50$ , data 3

## Additional assumption




Figure:  $y = x + N(0, 0.04)$ ,  $n = 50$ , data 4

## Additional assumption




Figure:  $y = x + N(0, 0.04)$ ,  $n = 50$ , data 5



## Frequentist distribution

- ▶ The slope:

$$\hat{\beta}_1 = \sum_{i=1}^n \frac{(x_i - \bar{x})}{\sum_{i=1}^n (x_i - \bar{x})^2} (y_i - \bar{y})$$

- ▶ The intercept

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

## Frequentist distribution

- ▶ The slope:

$$\hat{\beta}_1 = \sum_{i=1}^n \frac{(x_i - \bar{x})}{\sum_{i=1}^n (x_i - \bar{x})^2} (y_i - \bar{y})$$

- ▶ The intercept

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

## Expectation

►  $E(\hat{\beta}_0) = \beta_0 \quad E(\hat{\beta}_1) = \beta_1$

## Probabilistic properties of the least squares estimate

- ▶ A note on variance calculation
- ▶ The variances are

$$\text{Var}(\hat{\beta}_1) = \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \quad \text{Var}(\hat{\beta}_0) = \sigma^2 \left[ \frac{1}{n} + \frac{\bar{x}^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \right]$$

## Probabilistic properties of the least squares estimate

- ▶ A note on variance calculation
- ▶ The variances are

$$\text{Var}(\hat{\beta}_1) = \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \quad \text{Var}(\hat{\beta}_0) = \sigma^2 \left[ \frac{1}{n} + \frac{\bar{x}^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \right]$$

## About normal distribution

- ▶ Standard normal distribution

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

- ▶  $N(\mu, \sigma^2)$

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

## About normal distribution




Figure: Density function of standard normal distribution

## About normal (Gaussian) distribution

- ▶ The most natural distribution
- ▶ Stable distribution
- ▶ If  $Z_1$  and  $Z_2$  are independent normal random variables, then  $Z_1 + Z_2$  is also a normal random variable.
- ▶ The distribution of  $Z_1 + Z_2$  is ...