

Stat GR5205 Lecture 4

Jingchen Liu

Department of Statistics
Columbia University

Hypothesis testing

- ▶ The rationale
- ▶ Test statistic
- ▶ Reference distribution
- ▶ p -value: the probability of observing a test statistic that is as extreme as or more extreme than the observed one.
- ▶ The null hypothesis is rejected if the p -value is less than a threshold, such as 5%.

Hypothesis testing

- ▶ The rationale
- ▶ Test statistic
- ▶ Reference distribution
- ▶ p -value: the probability of observing a test statistic that is as extreme as or more extreme than the observed one.
- ▶ The null hypothesis is rejected if the p -value is less than a threshold, such as 5%.

Hypothesis testing

- ▶ The rationale
- ▶ Test statistic
- ▶ Reference distribution
- ▶ p -value: the probability of observing a test statistic that is as extreme as or more extreme than the observed one.
- ▶ The null hypothesis is rejected if the p -value is less than a threshold, such as 5%.

Hypothesis testing

- ▶ The rationale
- ▶ Test statistic
- ▶ Reference distribution
- ▶ p -value: the probability of observing a test statistic that is **as extreme as or more extreme** than the observed one.
- ▶ The null hypothesis is rejected if the p -value is less than a threshold, such as 5%.

Hypothesis testing

- ▶ The rationale
- ▶ Test statistic
- ▶ Reference distribution
- ▶ p -value: the probability of observing a test statistic that is **as extreme as or more extreme** than the observed one.
- ▶ The null hypothesis is rejected if the p -value is less than a threshold, such as 5%.

Hypothesis testing

- ▶ Size of a test: the probability of making type I error.
- ▶ Power of a test: the probability of rejecting the null under the alternative hypothesis

Hypothesis testing

- ▶ Size of a test: the probability of making type I error.
- ▶ Power of a test: the probability of rejecting the null under the alternative hypothesis

Hypothesis testing

- ▶ The Z-test

$$z = \frac{\hat{\beta}_1}{\sigma \sqrt{\frac{1}{\sum (x_i - \bar{x})}}} \sim N(0, 1)$$

- ▶ The t -test

$$t = \frac{\hat{\beta}_1}{\hat{\sigma} \sqrt{\frac{1}{\sum (x_i - \bar{x})}}} \sim t_{n-2}$$

Hypothesis testing

- ▶ The Z-test

$$z = \frac{\hat{\beta}_1}{\sigma \sqrt{\frac{1}{\sum (x_i - \bar{x})}}} \sim N(0, 1)$$

- ▶ The t -test

$$t = \frac{\hat{\beta}_1}{\hat{\sigma} \sqrt{\frac{1}{\sum (x_i - \bar{x})}}} \sim t_{n-2}$$

Hypothesis testing – p -value

- ▶ Two-sided p -value: $H_0 : \beta_1 = 0$ $H_1 : \beta_1 \neq 0$

$$P(|Z| \geq |z|) \quad \text{or} \quad P(|t_{n-2}| \geq |t|)$$

- ▶ One-sided p -value: $H_0 : \beta_1 \leq 0$ $H_1 : \beta_1 > 0$

$$P(Z \geq z) \quad \text{or} \quad P(t_{n-2} \geq t)$$

Hypothesis testing – p -value

- ▶ Two-sided p -value: $H_0 : \beta_1 = 0$ $H_1 : \beta_1 \neq 0$

$$P(|Z| \geq |z|) \quad \text{or} \quad P(|t_{n-2}| \geq |t|)$$

- ▶ One-sided p -value: $H_0 : \beta_1 \leq 0$ $H_1 : \beta_1 > 0$


$$P(Z \geq z) \quad \text{or} \quad P(t_{n-2} \geq t)$$

One-sample and two-sample t-test

Display 1.4

p. 5

Frequency histograms for male and female starting salaries



One-sample and two-sample t-test

- ▶ $\bar{Y}_{MALE} = 5957, \bar{Y}_{FEMALE} = 5139$
- ▶ $\bar{Y}_{MALE} - \bar{Y}_{FEMALE} = 818, \hat{\sigma} = 596$



$$t - statistic = \frac{\bar{Y}_{MALE} - \bar{Y}_{FEMALE}}{\hat{\sigma} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} = 6.2926.$$

Hypothesis testing – Duality

- ▶ Hypothesis testing and confidence interval
- ▶ The null is rejected if the confidence interval does not cover the hypothesized value. Justification!

Hypothesis testing – Duality

- ▶ Hypothesis testing and confidence interval
- ▶ The null is rejected if the confidence interval does not cover the hypothesized value. Justification!

Analysis of variance

- The decomposition

$$\sum (y_i - \bar{y})^2 = \sum (y_i - \hat{y}_i)^2 + \sum (\hat{y}_i - \bar{y})^2$$

$$SS_{total} = SS_{error} + SS_{regression}$$

- The analysis of variance table

	d.f.	Sum Sq	Mean Sq	F-value	p-value
x	$p - 1$	$\sum (\hat{y}_i - \bar{y})^2$	$\frac{\sum (\hat{y}_i - \bar{y})^2}{p - 1}$	$\frac{(n - p) \sum (\hat{y}_i - \bar{y})^2}{(p - 1) \sum (y_i - \hat{y}_i)^2}$	*
Residuals	$n - p$	$\sum (y_i - \hat{y}_i)^2$	$\frac{\sum (y_i - \hat{y}_i)^2}{n - p}$		
Total	$n - 1$	$\sum (y_i - \bar{y})^2$			

Analysis of variance

- The decomposition

$$\sum (y_i - \bar{y})^2 = \sum (y_i - \hat{y}_i)^2 + \sum (\hat{y}_i - \bar{y})^2$$

$$SS_{total} = SS_{error} + SS_{regression}$$

- The analysis of variance table

	d.f.	Sum Sq	Mean Sq	F-value	p-value
x	$p - 1$	$\sum (\hat{y}_i - \bar{y})^2$	$\frac{\sum (\hat{y}_i - \bar{y})^2}{p - 1}$	$\frac{(n - p) \sum (\hat{y}_i - \bar{y})^2}{(p - 1) \sum (y_i - \hat{y}_i)^2}$	*
Residuals	$n - p$	$\sum (y_i - \hat{y}_i)^2$	$\frac{\sum (y_i - \hat{y}_i)^2}{n - p}$		
Total	$n - 1$	$\sum (y_i - \bar{y})^2$			

Analysis of variance

- ▶ The analysis of variance table of the Iris setosa data

	d.f.	Sum Sq	Mean Sq	<i>F</i> -value	<i>p</i> -value
x	1	3.9	3.2	59.0	7×10^{-10}
Residuals	48	3.2	0.066		
Total	49	7.1			

- ▶ *F*-test and *t*-test

Analysis of variance

- ▶ The analysis of variance table of the Iris setosa data

	d.f.	Sum Sq	Mean Sq	<i>F</i> -value	<i>p</i> -value
x	1	3.9	3.2	59.0	7×10^{-10}
Residuals	48	3.2	0.066		
Total	49	7.1			

- ▶ *F*-test and *t*-test

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ Z-test
 - ▶ t-test: special case – two sample test
 - ▶ F-test: analysis of variance, R^2

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ Z-test
 - ▶ *t*-test: special case – two sample test
 - ▶ *F*-test: analysis of variance, R^2

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ Z-test
 - ▶ t-test: special case – two sample test
 - ▶ F-test: analysis of variance, R^2

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ Z-test
 - ▶ t-test: special case – two sample test
 - ▶ F-test: analysis of variance, R^2

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ Z-test
 - ▶ t-test: special case – two sample test
 - ▶ F-test: analysis of variance, R^2

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ *Z*-test
 - ▶ *t*-test: special case – two sample test
 - ▶ *F*-test: analysis of variance, R^2

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ *Z*-test
 - ▶ *t*-test: special case – two sample test
 - ▶ *F*-test: analysis of variance, R^2

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ *Z*-test
 - ▶ *t*-test: special case – two sample test
 - ▶ *F*-test: analysis of variance, R^2

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ Z -test
 - ▶ t -test: special case – two sample test
 - ▶ F -test: analysis of variance, R^2

Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ Z-test
 - ▶ *t*-test: special case – two sample test
 - ▶ *F*-test: analysis of variance, R^2


Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ Z-test
 - ▶ t -test: special case – two sample test
 - ▶ F -test: analysis of variance, R^2


Inferential tools of simple linear model

- ▶ The four assumptions
- ▶ Point estimate
 - ▶ Understanding the least squares estimate
 - ▶ variance estimate
 - ▶ Frequentist's distribution
- ▶ Interval estimate
 - ▶ Regression coefficients
 - ▶ Prediction: conditional mean, future observation, simultaneous confidence band
- ▶ Hypothesis testing
 - ▶ Z-test
 - ▶ t -test: special case – two sample test
 - ▶ F -test: analysis of variance, R^2



Diagnosis – linearity






Some possible deviation away from the assumption




Diagnosis – graphical analysis




Diagnosis – box plot




Diagnosis – quantile-quantile plot




Diagnosis – quantile-quantile plot



Diagnosis – residuals



Some possible deviation away from the assumption




Diagnostic test

- ▶ Pure significant test
- ▶ Equal variance test: Brown-Forsythe test and Levene's test

Diagnostic test

- ▶ Pure significant test
- ▶ Equal variance test: Brown-Forsythe test and Levene's test


Howard Levene




Transformation 101

- ▶ Logarithm!


Logarithm




Logarithm




Logarithm



Logarithm



Logarithm



Box-Cox tranformation

$$f_{\lambda}(y)$$

- ▶ $f_{\lambda}(y) = (y^{\lambda} - 1)/\lambda$ if $\lambda \neq 0$; $f_{\lambda}(y) = \log y$, if $\lambda = 0$.
- ▶ Choice of λ : maximum likelihood estimate
- ▶ `library{car}`: `box.cox.powers`

Box-Cox tranformation

$$f_{\lambda}(y)$$

- ▶ $f_{\lambda}(y) = (y^{\lambda} - 1)/\lambda$ if $\lambda \neq 0$; $f_{\lambda}(y) = \log y$, if $\lambda = 0$.
- ▶ Choice of λ : maximum likelihood estimate
- ▶ `library{car}: box.cox.powers`

Box-Cox tranformation

$$f_{\lambda}(y)$$

- ▶ $f_{\lambda}(y) = (y^{\lambda} - 1)/\lambda$ if $\lambda \neq 0$; $f_{\lambda}(y) = \log y$, if $\lambda = 0$.
- ▶ Choice of λ : maximum likelihood estimate
- ▶ `library{car}`: `box.cox.powers`

Some other transformations

$$y \in [0, 1]$$

- ▶ Logit transform: $\log \frac{y}{1-y}$
- ▶ Probit transform: $F^{-1}(y)$ where $F(x) = P(Z \leq x)$
- ▶ etc.

Some other transformations

$$y \in [0, 1]$$

- ▶ Logit transform: $\log \frac{y}{1-y}$
- ▶ Probit transform: $F^{-1}(y)$ where $F(x) = P(Z \leq x)$
- ▶ etc.

Some other transformations

$$y \in [0, 1]$$

- ▶ Logit transform: $\log \frac{y}{1-y}$
- ▶ Probit transform: $F^{-1}(y)$ where $F(x) = P(Z \leq x)$
- ▶ etc.