#### Spring 2018 STAT5261 Midterm Practice Questions

No Submission is required. Solution will be posted later.

### 1 Fixed Income Securities

- 1. Suppose that the yearly compounded rate is 3%. There is a 5-year coupon bond with face value (PAR) \$1000 and yearly coupon payment \$30. Suppose that this coupon bond may default and the default time X follows a geometric distribution:  $P(X = t) = (1 p)^{t-1}p$ , t = 1, 2, ... and p = 0.05. Find the expected present value of the coupon. (if the default happens at year 1, no coupon or the premium will be paid; if the default happens at year 2, only the coupon \$30 for year 1 will be paid...)
- 2. Suzanne paid \$800 for a 1-year zero-coupon bond with face value (PAR) \$1000. The current 1-year interest rate (yearly compounded) is r = 5%. Suppose there is a 10% probability that the bond will default and the recovery rate is 0 (i.e. no recovery).
  - (a) For Suzanne's purchase, the (1-year) net return is a random variable due to the possibility of default. Compute this (1-year) net return and find its mean and standard deviation.
  - (b) Suppose that Suzanne's portfolio contains, in addition to the above bond (A), another 1-year zero-coupon bond (B) for which she also paid \$800. Suppose that bond B has face value (PAR) of \$1000 and that its default probability is 10%. If the two default events are mutually exclusive, find the mean and standard deviation of the (1-year) net return of this portfolio.
- 3. Suppose that the forward rate is  $r(t) = 0.03 + 0.001t + 0.0002t^2$ 
  - (a) What is the 5-year spot rate?
  - (b) What is the price of a zero-coupon bond that matures in 5 years?
- 4. Suppose that the continuous forward rate is r(t) = 0.04 + 0.001t when a 8-year zero coupon bond is purchased. Six months later the forward rate is r(t) = 0.03 + 0.0013t and the bond is sold. What is the return?

# 2 Portfolio Theory

- 1. Suppose that the market consists of two risky assets. The mean and standard deviation of the return for asset 1 are  $\mu_1 = 10\%$  and  $\sigma_1 = 20\%$ , respectively and those for asset 2 are  $\mu_2 = 4\%$  and  $\sigma_2 = 10\%$ .
  - (a) Assume that the two returns are uncorrelated. Find the minimum variance point,  $(\sigma_{mv}, \mu_{mv})$ , on the  $\mu \sigma$  diagram for this market.
  - (b) What if the correlation between the two returns is  $\rho = 0.3$  or  $\rho = -0.3$ ?

- 2. Suppose that a market consists of one risk-free asset with rate of return  $\mu_f = 1\%$ , and two risky assets with means and standard deviations of returns being  $\mu_1 = 3\%$ ,  $\mu_2 = 6\%$ ,  $\sigma_1 = 20\%$  and  $\sigma_2 = 30\%$ .
  - (a) Find the Sharpe ratios of the tangency portfolios for different  $\rho = 0.2, 0.4, 0.8$  and write down the equations for the capital market lines.
  - (b) Suppose instead we have  $\sigma_1 = 40\%$  and  $\sigma_2 = 60\%$ . How do Sharpe ratios change and the portfolio weights change corresponding to the changes in  $\sigma$ 's?
- 3. Suppose that the risk-free interest rate is 0.023, that the expected return on the market portfolio is  $\mu_M = 0.10$ , and that the volatility of the market portfolio is  $\sigma_M = 0.12$ .
  - (a) What is the expected return on an efficient portfolio with  $\sigma_R = 0.05$ ?
  - (b) Stock A returns have a covariance of 0.004 with market returns. What is the beta of Stock A?
  - (c) Stock B has beta equal to 1.5 and  $\sigma_{\epsilon} = 0.08$ . Stock C has beta equal to 1.8 and  $\sigma_{\epsilon} = 0.10$ . What is the expected return of a portfolio that is one-half Stock B and one-half Stock C? What is the volatility of a portfolio that is one-half Stock B and one-half Stock C? Assume that the  $\epsilon$ 's of Stocks B and C are independent.

## 3 Rank Correlation and Copula

- 1. U is a uniform random variable on the interval [0,1], V is a Bernoulli random variable with P(V=1)=p, where  $0 \le p \le 1$ . Suppose U and V are independent, Compute Kendall's  $\tau$ , Spearman's  $\rho$  and Pearson's correlation coefficient for (U, U+V) as a function of p.
- 2. Let (X,Y) has the bivariate distribution  $F_{\theta}(x,y)$  of the form

$$F_{\theta}(x,y) = 1 - e^{-x} - e^{-y} - e^{-(x+y+\theta xy)}$$

if  $x \ge 0, y \ge 0$  and  $F_{\theta}(x, y) = 0$  otherwise. Here  $\theta \in [0, 1]$  is a parameter.

- (a) What are the marginal distributions of X and Y?
- (b) What is the copula function associated with (X,Y)?
- 3. Suppose that we have two bonds A and B. Denote by  $T_A$  and  $T_B$  their respective default times (in year). Suppose that  $T_A$  follows exponential distribution with hazard  $\lambda_A = 0.03$  (i.e.  $P(T_A \ge t) = e^{-\lambda_A t}$ ) and  $T_B$  follows exponential distribution with hazard  $\lambda_B = 0.02$ . Suppose that jointly they satisfy the Gumbel copula with  $\alpha = 3$ . Find the probabilities that
  - (a) Both will default by the end of the second year;
  - (b) At least one will default by the end of the second year.
  - (c) Only Bond A defaults by the end of the second year.

## 4 Risk Management

1. Suppose that  $\mathcal{L}$  is the loss over 1 year. Suppose that  $\mathcal{L}$  follows a double exponential distribution, whose density  $f(\cdot)$  has the form

$$f(x) = \frac{1}{Z} \exp(-\lambda |x|), \quad x \in \mathbb{R}$$

where Z is the normalizing constant and  $\lambda > 0$  is a parameter. Find

- (a) VaR(0.05) as a function of  $\lambda$
- (b) ES(0.05) as a function of  $\lambda$ .
- 2. Suppose the yearly returns of stock A, B and C have a multivariate normal distribution with mean (0.03, 0.04, 0.05) and covariance matrix

$$\begin{pmatrix} 0.4 & 0.3 & -0.1 \\ 0.3 & 0.5 & 0.2 \\ -0.1 & 0.2 & 0.8 \end{pmatrix}.$$

Suppose we have a portfolio that invest \$50,000, \$100,000 and \$200,000 to stock A, B and C respectively. Compute the following quantities for the portfolio:

- (a) VaR(0.05).
- (b) ES(0.05).
- 3. For a loss random variable L, recall that expected shortfall is defined as

$$ES(\alpha) := \frac{\int_0^\alpha VaR(s)ds}{\alpha}.$$

Another risk measure, the conditional tail expectation (CTE) is defined as

$$CTE(\alpha) = \mathbb{E}(L|L > VaR(\alpha)). \tag{1}$$

Show that  $ES(\alpha) = CTE(\alpha)$  when L is a continuous random variable. Note that (1) can be used to derive the expected shortfall formula when the loss follows a normal distribution (see HW6).