

# Stat GU4205/5205 Lecture 1

Jingchen Liu

Department of Statistics  
Columbia University

- ▶ Textbook: *Applied Linear Regression Models*, fourth edition, Kutner, Nachtsheim, and Neter
- ▶ Teaching assistants: Guanhua Fang
- ▶ Course design: statistics MA
- ▶ Office hours: TBD

# Syllabus

- ▶ Prerequisites: multivariate calculus, linear algebra, Stat GU4203, and GU4204
- ▶ Course material: simple linear regression, multiple linear regression, model diagnostics, model selection, and all other related statistical issues.
- ▶ Statistical inference: point estimation, hypothesis testing, confidence interval
- ▶ Basic statistical concepts
- ▶ Software: R <http://cran.r-project.org/>

# Syllabus

- ▶ Prerequisites: multivariate calculus, linear algebra, Stat GU4203, and GU4204
- ▶ Course material: simple linear regression, multiple linear regression, model diagnostics, model selection, and all other related statistical issues.
- ▶ Statistical inference: point estimation, hypothesis testing, confidence interval
- ▶ Basic statistical concepts
- ▶ Software: R <http://cran.r-project.org/>

# Syllabus

- ▶ Prerequisites: multivariate calculus, linear algebra, Stat GU4203, and GU4204
- ▶ Course material: simple linear regression, multiple linear regression, model diagnostics, model selection, and all other related statistical issues.
- ▶ Statistical inference: point estimation, hypothesis testing, confidence interval
- ▶ Basic statistical concepts
- ▶ Software: R <http://cran.r-project.org/>

# Syllabus

- ▶ Prerequisites: multivariate calculus, linear algebra, Stat GU4203, and GU4204
- ▶ Course material: simple linear regression, multiple linear regression, model diagnostics, model selection, and all other related statistical issues.
- ▶ Statistical inference: point estimation, hypothesis testing, confidence interval
- ▶ Basic statistical concepts
- ▶ Software: R <http://cran.r-project.org/>

# Syllabus

- ▶ Prerequisites: multivariate calculus, linear algebra, Stat GU4203, and GU4204
- ▶ Course material: simple linear regression, multiple linear regression, model diagnostics, model selection, and all other related statistical issues.
- ▶ Statistical inference: point estimation, hypothesis testing, confidence interval
- ▶ Basic statistical concepts
- ▶ Software: R <http://cran.r-project.org/>

# Grading

- ▶ Homework: 30%
- ▶ Midterm: 30%
- ▶ Final: 40%



## About statistics

- ▶ Statistics is “a mathematical science pertaining to the collection, analysis, interpretation or explanation, and presentation of data” – Wikipedia
- ▶ Statistical modeling: capturing the pattern of data for interpretation and prediction

## About statistics

- ▶ Statistics is “a mathematical science pertaining to the collection, analysis, interpretation or explanation, and presentation of data” – Wikipedia
- ▶ Statistical modeling: capturing the pattern of data for interpretation and prediction

## Single variable

- ▶ Random versus deterministic
- ▶ Probability: the chance of raining tomorrow is 40%.
- ▶ **Distribution**: characterizing the behavior of a random variable (object).

## Single variable

- ▶ Random versus deterministic
- ▶ Probability: the chance of raining tomorrow is 40%.
- ▶ **Distribution**: characterizing the behavior of a random variable (object).

## Single variable


- ▶ Random versus deterministic
- ▶ Probability: the chance of raining tomorrow is 40%.
- ▶ **Distribution**: characterizing the behavior of a random variable (object).

## Daily log-return, December 31, 2013

$$\log(S_{\text{today}} / S_{\text{yesterday}})$$

IBM: 0.6%, AAPL: 0.6%, GS: 0.8%, BAC: 0.1% ...

## Graphical illustration



**Figure:** Histogram of a single random variable

## Basic statistics

$$x_1, \dots, x_n$$

- ▶ Sample mean:

$$\bar{x} = \frac{x_1 + \dots + x_n}{n} = \frac{\sum_{i=1}^n x_i}{n}$$

- ▶ Sample variance:

$$s^2 = \frac{(x_1 - \bar{x})^2 + \dots + (x_n - \bar{x})^2}{n - 1} = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}$$

- ▶ Sample standard deviation:  $s$



## Basic statistics

$$x_1, \dots, x_n$$

- ▶ Sample mean:

$$\bar{x} = \frac{x_1 + \dots + x_n}{n} = \frac{\sum_{i=1}^n x_i}{n}$$

- ▶ Sample variance:

$$s^2 = \frac{(x_1 - \bar{x})^2 + \dots + (x_n - \bar{x})^2}{n - 1} = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}$$

- ▶ Sample standard deviation:  $s$

## Basic statistics

$$x_1, \dots, x_n$$

- ▶ Sample mean:

$$\bar{x} = \frac{x_1 + \dots + x_n}{n} = \frac{\sum_{i=1}^n x_i}{n}$$

- ▶ Sample variance:

$$s^2 = \frac{(x_1 - \bar{x})^2 + \dots + (x_n - \bar{x})^2}{n - 1} = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}$$

- ▶ Sample standard deviation:  $s$

# Basic statistics

- ▶ Median

- ▶  $x_1, \dots, x_{2n+1}$ :  $m = x_n$
- ▶  $x_1, \dots, x_{2n}$ :  $m = \frac{x_n + x_{n+1}}{2}$

- ▶ Quantile

## Basic statistics

- ▶ Median

- ▶  $x_1, \dots, x_{2n+1}$ :  $m = x_n$
- ▶  $x_1, \dots, x_{2n}$ :  $m = \frac{x_n + x_{n+1}}{2}$

- ▶ Quantile

## Basic statistics

- ▶ Median

- ▶  $x_1, \dots, x_{2n+1}$ :  $m = x_n$
- ▶  $x_1, \dots, x_{2n}$ :  $m = \frac{x_n + x_{n+1}}{2}$

- ▶ Quantile

## Basic statistics

- ▶ Median

- ▶  $x_1, \dots, x_{2n+1}$ :  $m = x_n$
- ▶  $x_1, \dots, x_{2n}$ :  $m = \frac{x_n + x_{n+1}}{2}$

- ▶ Quantile

## Two variables




Figure: 2013 Bank of America versus Goldman Sachs

- Deterministic versus random
- Predictability

## Two variables




Figure: 2013 Bank of America versus Goldman Sachs

- ▶ Deterministic versus random
- ▶ Predictability



## Two variables

$$(x_1, y_1), \dots, (x_n, y_n)$$

► Covariance

$$C_{x,y} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{n - 1}$$

► Correlation

$$\rho_{x,y} = \frac{C_{x,y}}{s_x s_y}$$

## Two variables

$$(x_1, y_1), \dots, (x_n, y_n)$$

- Covariance

$$C_{x,y} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{n - 1}$$

- Correlation

$$\rho_{x,y} = \frac{C_{x,y}}{s_x s_y}$$

## Two variables




Figure: 2013 Bank of America versus Apple

## A nonlinear example




Figure: Major depression prevalence rate against age

## Another nonlinear example




Figure: Corn yield against rain fall

## Regression models

- ▶ Separating the predictable part from the random part
- ▶ Basic setting:
  - ▶ Predictor (covariates, independent variable):  $X$
  - ▶ Response variable (dependent variable):  $Y$

## Regression models

- ▶ Separating the predictable part from the random part
- ▶ Basic setting:
  - ▶ Predictor (covariates, independent variable):  $X$
  - ▶ Response variable (dependent variable):  $Y$

## Regression models

- ▶ Separating the predictable part from the random part
- ▶ Basic setting:
  - ▶ Predictor (covariates, independent variable):  $X$
  - ▶ Response variable (dependent variable):  $Y$



## Regression models

- ▶ Separating the predictable part from the random part
- ▶ Basic setting:
  - ▶ Predictor (covariates, independent variable):  $X$
  - ▶ Response variable (dependent variable):  $Y$

# The use of regression models

- ▶ Causality versus association
  - ▶ Clinical trial
  - ▶ Genetic association study
  - ▶ Economics
- ▶ Prediction
- ▶ Signal detection, variable selection

# The use of regression models

- ▶ Causality versus association
  - ▶ Clinical trial
  - ▶ Genetic association study
  - ▶ Economics
- ▶ Prediction
- ▶ Signal detection, variable selection

# The use of regression models

- ▶ Causality versus association
  - ▶ Clinical trial
  - ▶ Genetic association study
  - ▶ Economics
- ▶ Prediction
- ▶ Signal detection, variable selection

# The use of regression models

- ▶ Causality versus association
  - ▶ Clinical trial
  - ▶ Genetic association study
  - ▶ Economics
- ▶ Prediction
- ▶ Signal detection, variable selection

# The use of regression models

- ▶ Causality versus association
  - ▶ Clinical trial
  - ▶ Genetic association study
  - ▶ Economics
- ▶ Prediction
- ▶ Signal detection, variable selection

# The use of regression models

- ▶ Causality versus association
  - ▶ Clinical trial
  - ▶ Genetic association study
  - ▶ Economics
- ▶ Prediction
- ▶ Signal detection, variable selection

## About causality



- ▶ Experimental study
- ▶ Observational study





## About causality

- ▶ Experimental study
- ▶ Observational study


## Experimental study versus observational study



# Line fitting



## Simple example



## Least squares estimator

$$(x_1, y_1), \dots, (x_n, y_n)$$

- ▶ Fitting a straight line  $y = \beta_0 + \beta_1 x$
- ▶ Sum of squares of residuals

$$SS = \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

- ▶ Least squares estimate

$$(\hat{\beta}_0, \hat{\beta}_1) = \arg \min_{\beta_0, \beta_1} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2.$$

## Least squares estimator

$$(x_1, y_1), \dots, (x_n, y_n)$$

- ▶ Fitting a straight line  $y = \beta_0 + \beta_1 x$
- ▶ Sum of squares of residuals

$$SS = \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

- ▶ Least squares estimate

$$(\hat{\beta}_0, \hat{\beta}_1) = \arg \min_{\beta_0, \beta_1} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2.$$

## Least squares estimator

$$(x_1, y_1), \dots, (x_n, y_n)$$

- ▶ Fitting a straight line  $y = \beta_0 + \beta_1 x$
- ▶ Sum of squares of residuals

$$SS = \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

- ▶ Least squares estimate

$$(\hat{\beta}_0, \hat{\beta}_1) = \arg \min_{\beta_0, \beta_1} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2.$$

## Another estimator

- ▶ Least squares estimate

$$(\hat{\beta}_0, \hat{\beta}_1) = \arg \min_{\beta_0, \beta_1} \sum_{i=1}^n |y_i - \beta_0 - \beta_1 x_i|.$$

- ▶ Sample mean and median.



## Another estimator

- ▶ Least squares estimate

$$(\hat{\beta}_0, \hat{\beta}_1) = \arg \min_{\beta_0, \beta_1} \sum_{i=1}^n |y_i - \beta_0 - \beta_1 x_i|.$$

- ▶ Sample mean and median.

# The rationale

- ▶ Prediction
- ▶ From the viewpoint of design,  $x$  is preset and  $y$  is observed.
- ▶ Probabilistic interpretation

## The rationale

- ▶ Prediction
- ▶ From the viewpoint of design,  $x$  is preset and  $y$  is observed.
- ▶ Probabilistic interpretation

## The rationale

- ▶ Prediction
- ▶ From the viewpoint of design,  $x$  is preset and  $y$  is observed.
- ▶ Probabilistic interpretation

- ▶ Derivation of least squares estimator

## Least squares estimator for simple linear regression

- ▶ The slope

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- ▶ The intercept

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

- ▶ Interpretation

## Least squares estimator for simple linear regression

- ▶ The slope

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- ▶ The intercept

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

- ▶ Interpretation

## Least squares estimator for simple linear regression

- ▶ The slope

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- ▶ The intercept

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

- ▶ Interpretation



## Least squares estimator for simple linear regression

- ▶ Another representation

$$\hat{\beta}_1 = \rho_{x,y} \frac{s_y}{s_x}.$$

- ▶ The fitted regression line

$$(x - \bar{x}) = \rho_{x,y} \frac{s_y}{s_x} (y - \bar{y})$$

## Least squares estimator for simple linear regression

- ▶ Another representation

$$\hat{\beta}_1 = \rho_{x,y} \frac{s_y}{s_x}.$$

- ▶ The fitted regression line

$$(x - \bar{x}) = \rho_{x,y} \frac{s_y}{s_x} (y - \bar{y})$$