

Advanced Mathematical Statistics: Assignment 2

Chris Hayduk

October 3, 2019

Problem 4.2.

a) $f(x_1, x_2) = \frac{1}{2\pi\sqrt{1.5}} \times \exp\left\{-\frac{2}{3}\left[\left(\frac{x_1}{\sqrt{2}}\right)^2 + (x_2 - 2)^2 - \left(\frac{x_1}{\sqrt{2}}\right)(x_2 - 2)\right]\right\}$


b) We have that $\rho_{12} = 0.5 \implies \sigma_{12} = 0.5(\sqrt{\sigma_{11}\sigma_{22}}) = 0.5(\sqrt{2})$. So,

$$\begin{aligned} & (\mathbf{x} - \mu)' \Sigma^{-1} (\mathbf{x} - \mu) \\ &= \begin{bmatrix} x_1 - \mu_1 & x_2 - \mu_2 \end{bmatrix} \frac{1}{\sigma_{11}\sigma_{22} - \sigma_{12}^2} \begin{bmatrix} \sigma_{22} & -\sigma_{12} \\ -\sigma_{12} & \sigma_{11} \end{bmatrix} \begin{bmatrix} x_1 - \mu_1 \\ x_2 - \mu_2 \end{bmatrix} \\ &= \begin{bmatrix} x_1 & x_2 - 2 \end{bmatrix} \frac{1}{2 - (0.5\sqrt{2})^2} \begin{bmatrix} 1 & -0.5\sqrt{2} \\ -0.5\sqrt{2} & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 - 2 \end{bmatrix} \\ &= \begin{bmatrix} \frac{2}{3}x_1 + (-\frac{\sqrt{2}}{3})(x_2 - 2) & (-\frac{\sqrt{2}}{3})(x_1) + \frac{4}{3}(x_2 - 2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 - 2 \end{bmatrix} \\ &= x_1 \left(\frac{2}{3}x_1 + \left(\frac{-\sqrt{2}}{3} \right) (x_2 - 2) \right) + (x_2 - 2) \left(\left(\frac{-\sqrt{2}}{3} \right) x_1 + \frac{4}{3}(x_2 - 2) \right) \\ &= \frac{2}{3}x_1^2 + \left(\frac{-\sqrt{2}}{3} \right) (x_2 - 2)x_1 + (x_2 - 2) \left(\frac{-\sqrt{2}}{3} \right) x_1 + \frac{4}{3}(x_2 - 2)^2 \\ &= \frac{2}{3}x_1^2 + \frac{-2\sqrt{2}(x_2x_1 - 2x_1)}{3} + \frac{4}{3}x_2^2 - \frac{16}{3}x_2 + \frac{16}{3} \end{aligned}$$

c) $c^2 = \chi_2^2(0.5) \approx 1.386294$. So we take

$$\frac{2}{3}x_1^2 + \frac{-2\sqrt{2}(x_2x_1 - 2x_1)}{3} + \frac{4}{3}x_2^2 - \frac{16}{3}x_2 + \frac{16}{3} = 1.386294$$

to be the surface of the ellipsoid containing 50% of the probability. The graph for this can be seen below.



Problem 4.3.

- a) X_1 and X_2 are not independent because $\sigma_{12} = \sigma_{21} = -2 \neq 0$.
- b) X_2 and X_3 are independent because $\sigma_{23} = \sigma_{32} = 0$.
- c) If we partition the covariance matrix into (X_1, X_2) and X_3 partitions, we get

$$\left[\begin{array}{cc|c} 1 & -2 & 0 \\ -2 & 5 & 0 \\ \hline 0 & 0 & 2 \end{array} \right]$$

Thus, we can see that the two diagonal sections of the matrix have the forms $\mathbf{0}, \mathbf{0}'$. As a result, (X_1, X_2) and X_3 are independent.

- d) Let $\mathbf{A} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix}$. From Result 4.3, we know \mathbf{AX} is distributed as $N_q(\mathbf{A}\mu, \mathbf{A}\Sigma\mathbf{A}')$ with $q = 2$ in this case.

So we have,

$$\begin{aligned}\mathbf{A}\Sigma\mathbf{A}' &= \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 0 \\ -2 & 5 & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & 0 \\ \frac{1}{2} & 0 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} -\frac{1}{2} & \frac{3}{2} & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & 0 \\ \frac{1}{2} & 0 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & 2 \end{bmatrix}\end{aligned}$$

As can be clearly seen from the above matrix, $\mathbf{A}\Sigma\mathbf{A}'$, the covariance between $\frac{X_1+X_2}{2}$ and X_3 is 0. As a result, $\frac{X_1+X_2}{2}$ and X_3 are independent.

- e) Let $\mathbf{A} = \begin{bmatrix} 0 & 1 & 0 \\ -\frac{5}{2} & 1 & -1 \end{bmatrix}$. From Result 4.3, we know \mathbf{AX} is distributed as $N_q(\mathbf{A}\mu, \mathbf{A}\Sigma\mathbf{A}')$ with $q = 2$ in this case.

So we have,

$$\begin{aligned}\mathbf{A}\Sigma\mathbf{A}' &= \begin{bmatrix} 0 & 1 & 0 \\ -\frac{5}{2} & 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & -2 & 0 \\ -2 & 5 & 0 \\ 0 & 0 & 2 \end{bmatrix} \begin{bmatrix} 0 & -\frac{5}{2} \\ 1 & 1 \\ 0 & -1 \end{bmatrix} \\ &= \begin{bmatrix} -2 & 5 & 0 \\ -\frac{9}{2} & 10 & -2 \end{bmatrix} \begin{bmatrix} 0 & -\frac{5}{2} \\ 1 & 1 \\ 0 & -1 \end{bmatrix} \\ &= \begin{bmatrix} 5 & 10 \\ 10 & \frac{93}{4} \end{bmatrix}\end{aligned}$$

We can see from the above matrix that the covariance between the random variables is not 0. Thus, X_2 and $X_2 - \frac{5}{2}X_1 - X_3$ are not independent.

Problem 4.4.

- a) Let $A = \begin{bmatrix} 3 & -2 & 1 \end{bmatrix}$. By Result 4.3, $3X_1 - 2X_2 + X_3$ is distributed as $N_1(\mathbf{A}\mu, \mathbf{A}\Sigma\mathbf{A}')$ with mean vector and covariance matrix,

$$\mathbf{A}\mu = 6 + 6 + 1 = 13$$

$$\mathbf{A}\Sigma\mathbf{A}' = \begin{bmatrix} 2 & -1 & 1 \end{bmatrix} \mathbf{A}' = 6 + 2 + 1 = 9$$

- b) Let $A = \begin{bmatrix} 0 & 1 & 0 \\ -a_1 & 1 & -a_2 \end{bmatrix}$.

Now find $\mathbf{A}\Sigma\mathbf{A}'$:

$$\begin{aligned}\mathbf{A}\Sigma\mathbf{A}' &= \begin{bmatrix} 1 & 3 & 2 \\ -a_1 + 1 - a_2 & -a_1 + 3 - 2a_2 & a_1 + 2 - 2a_2 \end{bmatrix} \begin{bmatrix} 0 & -a_1 \\ 1 & 1 \\ 0 & -a_2 \end{bmatrix} \\ &= \begin{bmatrix} 3 & -a_1 + 3 - 2a_2 \\ -a_1 + 3 - 2a_2 & (-a_1)^2 - 2a_1 - 2a_1a_2 + 3 - 4a_2 + 2(-a_2)^2 \end{bmatrix}\end{aligned}$$

In order for the covariance to be 0 (ie. X_2 and $-a_1X_1 + X_2 - a_3X_3$ independent), we need $-a_1 + 3 - 2a_2 = 0$. That is, $a_1 + 2a_2 = 3$. For instance, take $a_1 = 1$ and $a_2 = 1$. Then we have,

$$\mathbf{A}\Sigma\mathbf{A}' = \begin{bmatrix} 3 & 0 \\ 0 & -2 \end{bmatrix}$$

Thus, if $\mathbf{a} = \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, then we have that X_2 and $X_2 - \mathbf{a}' \begin{bmatrix} X_1 \\ X_3 \end{bmatrix}$ are independent.

Problem 4.5.

a) By Result 4.6, we have

$$\begin{aligned}\mu &= \mu_1 + \sigma_{12}(\sigma_{22})^{-1}(x_2 - \mu_2) \\ &= 0 + 0.5(\sqrt{2})(1)^{-1}(x_2 - 2) \\ &= 0.5(\sqrt{2})x_2 - \sqrt{2}\end{aligned}$$

In addition, we have

$$\begin{aligned}\sigma &= \sigma_{11} - \sigma_{12}(\sigma_{22})^{-1}\sigma_{21} \\ &= 2 - 0.5(\sqrt{2})(1)^{-1}0.5(\sqrt{2}) \\ &= 2 - 0.25(2) = 1.5\end{aligned}$$

b) Let's rearrange the Σ matrix,

$$\begin{aligned}\Sigma &= \begin{bmatrix} 1 & 0 & -2 \\ 0 & 2 & 0 \\ -2 & 0 & 5 \end{bmatrix} \\ &= \left[\begin{array}{cc|c} 1 & 0 & -2 \\ 0 & 2 & 0 \\ -2 & 0 & 5 \end{array} \right]\end{aligned}$$

and the μ vector,

$$\begin{aligned}\mu &= \begin{bmatrix} -3 \\ 4 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} -3 \\ 4 \\ 1 \end{bmatrix}\end{aligned}$$

Then we have,

$$\begin{aligned}\mu &= \mu_1 + \Sigma_{12}\Sigma_{22}^{-1}(x_2 - \mu_2) \\ &= \begin{bmatrix} -3 \\ 4 \end{bmatrix} + \begin{bmatrix} -2 \\ 0 \end{bmatrix} \left(\frac{1}{5}\right)(x_2 - 1) \\ &= \begin{bmatrix} -3 \\ 4 \end{bmatrix} + \begin{bmatrix} -2 \\ 0 \end{bmatrix} \left(\frac{1}{5}x_2 - \frac{1}{5}\right)\end{aligned}$$

and,

$$\begin{aligned}\Sigma &= \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} - \begin{bmatrix} -2 \\ 0 \end{bmatrix} \left(\frac{1}{5}\right) \begin{bmatrix} -2 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} - \begin{bmatrix} -2 \\ 0 \end{bmatrix} \begin{bmatrix} -\frac{2}{5} & 0 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} - \begin{bmatrix} \frac{4}{5} & 0 \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{5} & 0 \\ 0 & 2 \end{bmatrix}\end{aligned}$$

c) If we partition the covariance matrix into (X_1, X_2) and X_3 partitions, we get

$$\begin{aligned}\Sigma &= \left[\begin{array}{cc|c} 1 & 1 & 1 \\ 1 & 3 & 2 \\ \hline 1 & 2 & 2 \end{array} \right] \\ &= \left[\begin{array}{c|c} \Sigma_{11} & \Sigma_{12} \\ \hline \Sigma_{21} & \Sigma_{22} \end{array} \right]\end{aligned}$$

and,

$$\begin{aligned}\mu &= \begin{bmatrix} 2 \\ -3 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}\end{aligned}$$

So we have,

$$\begin{aligned}
\mu &= \mu_2 + \Sigma_{21}(\Sigma_{11})^{-1}(x_1 - \mu_1) \\
&= 1 + \begin{bmatrix} 1 & 2 \end{bmatrix} \left(\begin{bmatrix} \frac{3}{2} & -\frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{bmatrix} \right) (x_1 - \begin{bmatrix} 2 \\ -3 \end{bmatrix}) \\
&= 1 + \left(\begin{bmatrix} \frac{1}{2} & \frac{1}{2} \end{bmatrix} \right) (x_1 - \begin{bmatrix} 2 \\ -3 \end{bmatrix}) \\
&= 1 + \left(\begin{bmatrix} \frac{1}{2} & \frac{1}{2} \end{bmatrix} \right) \left(\begin{bmatrix} X_1 \\ X_2 \end{bmatrix} - \begin{bmatrix} 2 \\ -3 \end{bmatrix} \right) \\
&= 1 + \left(\begin{bmatrix} \frac{1}{2} & \frac{1}{2} \end{bmatrix} \right) \left(\begin{bmatrix} X_1 - 2 \\ X_2 + 3 \end{bmatrix} \right) \\
&= 1 + \frac{1}{2} (X_1 - 2 + X_2 + 3) \\
&= \frac{1}{2}X_1 + \frac{1}{2}X_2 + \frac{3}{2}
\end{aligned}$$

and,

$$\begin{aligned}
\Sigma &= \Sigma_{22} - \Sigma_{21}(\Sigma_{11})^{-1}\Sigma_{12} \\
&= 2 - \begin{bmatrix} 1 & 2 \end{bmatrix} \left(\begin{bmatrix} \frac{3}{2} & -\frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{bmatrix} \right) \begin{bmatrix} 1 \\ 2 \end{bmatrix} \\
&= 2 - \left(\begin{bmatrix} \frac{1}{2} & \frac{1}{2} \end{bmatrix} \right) \begin{bmatrix} 1 \\ 2 \end{bmatrix} \\
&= 2 - \frac{3}{2} = \frac{1}{2}
\end{aligned}$$

Problem 4.6.

- a) We can see that $\sigma_{12} = \sigma_{21} = 0$. Thus, X_1 and X_2 are independent.
- b) We have that $\sigma_{13} = \sigma_{31} = -1 \neq 0$. Thus, X_1 and X_3 are not independent.
- c) We have that $\sigma_{23} = \sigma_{32} = 0$. Thus, X_2 and X_3 are independent.
- d) Let's rearrange the covariance matrix,

$$\begin{bmatrix} 4 & -1 & 0 \\ -1 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

We can partition it now as well,

$$\left[\begin{array}{cc|c} 4 & -1 & 0 \\ -1 & 2 & 0 \\ \hline 0 & 0 & 2 \end{array} \right]$$

So now we have the covariance matrix partitioned into (X_1, X_3) and X_2 blocks, along with their covariances. It is clear from this partitioning that (X_1, X_3) and X_2 have zero covariance. Thus, they are independent.

- e) Let $\mathbf{A} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 3 & -2 \end{bmatrix}$. From Result 4.3, we know \mathbf{AX} is distributed as $N_q(\mathbf{A}\mu, \mathbf{A}\Sigma\mathbf{A}')$ with $q = 2$ in this case.

So we have,

$$\begin{aligned} \mathbf{A}\Sigma\mathbf{A}' &= \begin{bmatrix} 1 & 0 & 0 \\ 1 & 3 & -2 \end{bmatrix} \begin{bmatrix} 4 & 0 & -1 \\ 0 & 5 & 0 \\ -1 & 0 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 3 \\ 0 & -2 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 0 & -1 \\ 6 & 6 & -5 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 3 \\ 0 & -2 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 6 \\ 6 & 34 \end{bmatrix} \end{aligned}$$

We can see from the above matrix that the covariance between the random variables is not 0. Thus, X_1 and $X_1 + 3X_2 - 2X_3$ are not independent.

Problem 4.7.

- a) Let's define the covariance matrix,

$$\Sigma = \begin{bmatrix} 4 & -1 \\ -1 & 2 \end{bmatrix}$$

By Result 4.6, we have

$$\begin{aligned} \mu &= \mu_1 + \sigma_{12}(\sigma_{22})^{-1}(x_2 - \mu_2) \\ &= 1 + -1 \left(\frac{1}{2} \right) (x_2 - 2) \\ &= 1 - \frac{1}{2}x_2 + 1 = \frac{1}{2} + 2 \end{aligned}$$

and,

$$\begin{aligned} \sigma &= \sigma_{11} - \sigma_{12}(\sigma_{22})^{-1}\sigma_{21} \\ &= 4 - (-1) \left(\frac{1}{2} \right) (-1) \\ &= 4 - \frac{1}{2} = \frac{7}{2} \end{aligned}$$

b) Let's partition the Σ matrix,

$$\left[\begin{array}{cc|c} 4 & 0 & -1 \\ 0 & 5 & 0 \\ -1 & 0 & 2 \end{array} \right] \quad (1)$$

and the μ vector,

$$\begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$$

Then we have,

$$\begin{aligned} \mu &= \mu_1 + \Sigma_{12}\Sigma_{22}^{-1}(\mathbf{x}_2 - \mu_2) \\ &= 1 + -1 \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix} \left(\begin{bmatrix} x_2 \\ x_3 \end{bmatrix} - \begin{bmatrix} -1 \\ 2 \end{bmatrix} \right) \\ &= 1 + -1 \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix} \left(\begin{bmatrix} x_2 + 1 \\ x_3 - 2 \end{bmatrix} \right) \end{aligned}$$

and,

$$\begin{aligned} \Sigma &= \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21} \\ &= \begin{bmatrix} 4 & 0 \end{bmatrix} - \begin{bmatrix} 0 \\ -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 0 & 5 \\ -1 & 0 \end{bmatrix} \\ &= \begin{bmatrix} 4 & 0 \end{bmatrix} - \begin{bmatrix} 0 & -\frac{5}{2} \end{bmatrix} \\ &= \begin{bmatrix} 4 & \frac{5}{2} \end{bmatrix} \end{aligned}$$

Problem 4.10.

Problem 4.11.

Problem 4.12.

Problem 4.13.

Problem 4.14.

Problem 4.15.

Problem 4.16.

Problem 4.17.

Problem 4.18.

Problem 4.19.

Problem 4.20.

Problem 4.21.