

# Stochastic Differential Equations: Final Project

Chris Hayduk

December 15, 2019

**NOTE: Still need to complete Problems 2, 7, 8, 15**

## **Problem 1.**

We know that the probability density of the sum of two independent random variables can be computed by the convolutions of  $f_X$  and  $f_Y$ . Thus, the probability density of  $X + Y$ , denoted by  $f_Z$ , is

$$f_Z(z) = (f_Y * f_X) = \int_{-\infty}^{\infty} f_Y(z - x) f_X(x) dx \quad (1)$$

We know that the normal density function is given by,

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

Plugging this density function into equation (1) and letting  $\sigma_Z = \sqrt{\sigma_X^2 + \sigma_Y^2}$  yields,

$$\begin{aligned}
f_Z(z) &= \int_{-\infty}^{\infty} \left[ \frac{1}{\sigma_Y \sqrt{2\pi}} e^{-\frac{1}{2} \left( \frac{z-x-\mu_Y}{\sigma_Y} \right)^2} \frac{1}{\sigma_X \sqrt{2\pi}} e^{-\frac{1}{2} \left( \frac{x-\mu_X}{\sigma_X} \right)^2} \right] dx \\
&= \int_{-\infty}^{\infty} \left[ \frac{1}{\sigma_Y \sigma_X (2\pi)} e^{-\frac{1}{2} \left[ \left( \frac{z-x-\mu_Y}{\sigma_Y} \right)^2 + \left( \frac{x-\mu_X}{\sigma_X} \right)^2 \right]} \right] dx \\
&= \int_{-\infty}^{\infty} \left[ \frac{1}{\sigma_Y \sigma_X (2\pi)} e^{-\frac{\sigma_X^2 (z-x-\mu_Y)^2 + \sigma_Y^2 (x-\mu_X)^2}{2\sigma_Y^2 \sigma_X^2}} \right] dx \\
&= \int_{-\infty}^{\infty} \left[ \frac{1}{(2\pi) \sigma_X \sigma_Y} e^{-\frac{x^2 (\sigma_X^2 + \sigma_Y^2) - 2x (\sigma_X^2 (z-\mu_Y) + \sigma_Y^2 \mu_X) + \sigma_X^2 (z^2 + \mu_Y^2 - 2z\mu_Y) + \sigma_Y^2 \mu_X^2}{2\sigma_Y^2 \sigma_X^2}} \right] dx \\
&= \int_{-\infty}^{\infty} \left[ \frac{1}{\sqrt{2\pi} \sigma_Z} e^{-\frac{\sigma_Z^2 (\sigma_X^2 (z-\mu_Y)^2 + \sigma_Y^2 \mu_X^2) - (\sigma_X^2 (z-\mu_Y) + \sigma_Y^2 \mu_X)^2}{2\sigma_Z^2 (\sigma_X \sigma_Y)^2}} \frac{1}{\sqrt{2\pi} \frac{\sigma_X \sigma_Y}{\sigma_Z}} e^{-\frac{\left( x - \frac{\sigma_X^2 (z-\mu_Y) + \sigma_Y^2 \mu_X}{\sigma_Z^2} \right)^2}{2 \left( \frac{\sigma_X \sigma_Y}{\sigma_Z} \right)^2}} \right] dx \\
&= \frac{1}{\sqrt{2\pi} \sigma_Z} e^{-\frac{(z-(\mu_X+\mu_Y))^2}{2\sigma_Z^2}} \int_{-\infty}^{\infty} \left[ \frac{1}{\sqrt{2\pi} \frac{\sigma_X \sigma_Y}{\sigma_Z}} e^{-\frac{\left( x - \frac{\sigma_X^2 (z-\mu_Y) + \sigma_Y^2 \mu_X}{\sigma_Z^2} \right)^2}{2 \left( \frac{\sigma_X \sigma_Y}{\sigma_Z} \right)^2}} \right] dx
\end{aligned}$$

The equation inside the integral symbol represents a valid normal density function for  $x$ , so we know it integrates to 1. Thus, the probability density function for  $X + Y$  is given by

$$f_Z(z) = \frac{1}{\sqrt{2\pi} \sigma_Z} e^{-\frac{(z-(\mu_X+\mu_Y))^2}{2\sigma_Z^2}}$$

This is precisely a normal density function with mean  $\mu_X + \mu_Y$  and variance  $\sigma_Z^2 = (\sqrt{\sigma_X^2 + \sigma_Y^2})^2 = \sigma_X^2 + \sigma_Y^2$ .

Hence, we have shown that, given  $X \sim N(\mu_X, \sigma_X^2)$  and  $Y \sim N(\mu_Y, \sigma_Y^2)$ ,  $X + Y$  is distributed as  $N(\mu_X + \mu_Y, \sigma_X^2 + \sigma_Y^2)$ .

## Problem 2.

Let  $Z = Y + 1$ . We'll begin by finding the probability density function for  $Z$ . Since  $Y$  is a standard uniform random variable, we know that

$$f(y) = \begin{cases} 1 & 0 \leq y \leq 1 \\ 0 & \text{elsewhere} \end{cases}$$

Thus, the distribution function approach yields,

$$\begin{aligned}
 F_Z(z) &= P(Z \leq z) = P(Y + 1 \leq z) = P(Y \leq z - 1) \\
 &= \int_{-\infty}^{z-1} f(y) dy \\
 &= \int_0^{z-1} 1 dy \\
 &= z - 1
 \end{aligned}$$

So, since as  $Y$  ranges from 0 to 1,  $Z$  ranges from 1 to 2, we have

$$F_Z(z) = \begin{cases} 0 & z < 1 \\ z - 1 & 1 \leq z \leq 2 \\ 1 & \text{elsewhere} \end{cases}$$

And the density function for  $Z$  is

$$f_Z(z) = \frac{dF_Z(z)}{dz} = \begin{cases} 1 & 1 \leq z \leq 2 \\ 0 & \text{elsewhere} \end{cases}$$

Now we can simplify the original function to be  $U = X/Z$ , where we now know the probability density function for both random variables ( $X \sim \text{Unif}(0, 1)$  and  $Z \sim \text{Unif}(1, 2)$ ).

So the distribution function is given by,

$$\begin{aligned}
 F_U(u) &= P(U \leq u) = P(X/Z \leq u) = P(X \leq uZ) \\
 &= \int_{-\infty}^{uz} f(x) dx \\
 &= \int_0^{uz} 1 dx \\
 &= uz
 \end{aligned}$$

FINISH THIS LATER.

### Problem 3.

Let  $Y$  be a standard normal variable. Then the moment-generating function of  $Y$  is given by,

$$\begin{aligned}
 m(t) &= E(e^{tY}) \\
 &= \int_{-\infty}^{\infty} e^{ty} f(y) dy
 \end{aligned}$$

Since  $Y$  is a standard normal variable, we know that  $\mu = 0$  and  $\sigma = 1$ . Thus, we have

$$f(y) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}y^2}$$

Plugging this into the above equation yields,

$$\begin{aligned}
m(t) &= \int_{-\infty}^{\infty} e^{ty} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}y^2} dy \\
&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2} + ty} dy \\
&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y-t)^2} e^{\frac{1}{2}t^2} dy \\
&= e^{\frac{1}{2}t^2} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(y-t)^2} dy
\end{aligned}$$

We can see that the integral is precisely the integral for a normal random variable with  $\mu = y - t$  and  $\sigma = 1$ . Thus, it integrates to 1, yielding,

$$m(t) = e^{\frac{1}{2}t^2}$$

**Problem 4.**

Suppose  $g(x)$  is a monotone increasing function and  $X$  is a random variable with the probability density function  $f_X$ .

Let  $U = g(X)$  where  $X$  has the above density function. Since  $g(x)$  is an increasing function of  $x$ , then  $g^{-1}(u)$  is an increasing function of  $u$ . Thus,

$$\begin{aligned}
P(U \leq u) &= P[g(X) \leq u] \\
&= P\{g^{-1}[g(X)] \leq g^{-1}(u)\} \\
&= P[X \leq g^{-1}(u)]
\end{aligned}$$

The above sequence of equalities implies that,

$$F_U(u) = F_X[g^{-1}(u)]$$

When we differentiate with respect to  $u$ , we get,

$$f_U(u) = \frac{dF_X[g^{-1}(u)]}{du} = f_X(g^{-1}(u)) \frac{d[g^{-1}(u)]}{du}$$

**Problem 5.**

We know that the moment-generating function of  $X$  is  $\phi(t)$ . Thus,

$$\begin{aligned}
\phi(t) &= E(e^{tX}) \\
&= \int_{-\infty}^{\infty} e^{tx} f(x) dx \\
&= \int_{-\infty}^{\infty} \left(1 + tx + \frac{t^2 x^2}{2!} + \frac{t^3 x^3}{3!} + \cdots\right) f(x) dx \\
&= \int_{-\infty}^{\infty} f(x) dx + t \int_{-\infty}^{\infty} x f(x) dx + \frac{t^2}{2!} \int_{-\infty}^{\infty} x^2 f(x) dx + \cdots
\end{aligned}$$

Now plugging in  $aX + b$  to the above equation yields,

$$\begin{aligned} E(e^{t(aX+b)}) &= E(e^{taX+tb}) \\ &= E[e^{taX}(e^{tb})] \end{aligned}$$

Since expected value is a linear operator and  $e^{tb}$  is a constant, we can pull it out of the expectation operator,

$$\begin{aligned} E[e^{taX}(e^{tb})] &= e^{tb} E(e^{taX}) \\ &= e^{tb} \int_{-\infty}^{\infty} e^{tax} f(ax) dx \end{aligned}$$

We can see from the above equation that this is equal to  $e^{tb}\phi(ta)$ .

Thus, the moment-generating function for  $aX + b$  with constants  $a \neq 0$  and  $b$  is  $e^{tb}\phi(ta)$ .

### Problem 6.

We know by Theorem 2.11.1 that the distribution of the sum of two random variables  $X$  and  $Y$  is given by the convolution of their densities,  $f_X$  and  $f_Y$ .

Thus, with  $Z = X + Y$ , we have,

$$f_Z(z) = (f * g)(z) = \int_{-\infty}^{\infty} f_X(z - y)f_Y(y)dy$$

The mean of this distribution is,

$$E(Z) = \int_{-\infty}^{\infty} z f_Z(z - y)f_Y(y)dy$$

where  $z = x + y$ . Thus, we have,

$$\begin{aligned} E(Z) &= \int_{-\infty}^{\infty} z f_Z(z)dz \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [(x + y)f_X((x + y) - y)f_Y(y)] dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [(x + y)f_X(x)f_Y(y)] dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [(xf_X(x) + yf_X(x))f_Y(y)] dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [xf_X(x)f_Y(y) + yf_X(x)f_Y(y)] dx dy \\ &= \int_{-\infty}^{\infty} [\mu_X f_Y(y) + yf_Y(y)] dy \\ &= \mu_X + \mu_Y \end{aligned}$$

In the above equations, we are using the properties that, for any random variable  $X$  with density function  $f(x)$  and mean  $\mu$ , we have that  $\int_{-\infty}^{\infty} f(x) = 1$  and  $\int_{-\infty}^{\infty} xf(x) = \mu$ .

Now for the variance, we know that for a given random variable  $X$ ,  $\text{Var}(X) = E[X^2] - E[X]^2$ . Plugging in  $Z$  to this formula yields,

$$\begin{aligned}
\text{Var}(Z) &= \sigma_Z^2 = E[Z^2] - E[Z]^2 \\
&= \int_{-\infty}^{\infty} z^2 f_Z(z) dz + (\mu_X + \mu_Y)^2 \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [(x+y)^2 f_X((x+y) - y) f_Y(y)] dx dy - (\mu_X + \mu_Y)^2 \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [(x+y)^2 f_X(x) f_Y(y)] dx dy - (\mu_X + \mu_Y)^2 \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [x^2 + 2xy + y^2] f_X(x) f_Y(y) dx dy - (\mu_X + \mu_Y)^2 \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} [x^2 f_X(x) f_Y(y) + 2xy f_X(x) f_Y(y) + y^2 f_X(x) f_Y(y)] dx dy - (\mu_X + \mu_Y)^2 \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x^2 f_X(x) f_Y(y) dx dy - \mu_X + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} 2xy f_X(x) f_Y(y) dx dy + \\
&\quad \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y^2 f_X(x) f_Y(y) dx dy - \mu_Y \\
&= \left[ \int_{-\infty}^{\infty} x^2 f_X(x) dx - \mu_X \right] + 2\mu_X \mu_Y + \left[ \int_{-\infty}^{\infty} y^2 f_Y(y) dy - \mu_Y \right] \\
&= \sigma_X^2 + 2\mu_X \mu_Y + \sigma_Y^2
\end{aligned}$$

Thus, we have  $\text{Var}(Z) = \sigma_X^2 + 2\mu_X \mu_Y + \sigma_Y^2$ .

### Problem 7.

The definition of the Central Limit Theorem is as follows:

Let  $Y_1, Y_2, \dots, Y_n$  be independent and identically distributed random variables with  $E(Y_i) = \mu$  and  $V(Y_i) = \sigma^2 < \infty$ . Define

$$U_n = \frac{\sum_{i=1}^n Y_i - n\mu}{\sigma\sqrt{n}} = \frac{\bar{Y} - \mu}{\sigma/\sqrt{n}}$$

where  $\bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i$ .

Then the distribution function of  $U_n$  converges to the standard normal distribution function as  $n \rightarrow \infty$ . That is,

$$\lim_{n \rightarrow \infty} P(U_n \leq u) = \int_{-\infty}^u \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt$$

for all  $u$ .

Let  $f_n(x)$  be the probability mass function for a binomial random variable with  $n$  trials and success probability  $p$ .

FINISH THIS LATER.

**Problem 8.**

Let  $X$  and  $Y$  be two independent, continuous random variables described by probability density functions  $f_X$  and  $f_Y$ . Also let  $Z = XY$ . We'll begin by finding the cumulative distribution function for  $Z$ . This yields,

$$\begin{aligned}
 F_Z(z) &= P(Z \leq z) \\
 &= P(XY \leq z) \\
 &= P(XY \leq z, X \geq 0) + P(XY \leq z, X \leq 0) \\
 &= P(Y \leq z/X, X \geq 0) + P(Y \geq z/X, X \leq 0) \\
 &= \int_0^\infty f_X(x) \int_{-\infty}^{z/x} f_Y(y) dy dx + \int_{-\infty}^0 f_X(x) \int_{z/X}^\infty f_Y(y) dy dx
 \end{aligned}$$

Now in order to find the probability density function for  $Z$ , we need to differentiate with respect to  $z$  on both sides of the above equation.

$$\begin{aligned}
 f_Z(z) &= \frac{d}{dz} F_Z(z) \\
 &= \frac{d}{dz} \left[ \int_0^\infty f_X(x) \int_{-\infty}^{z/x} f_Y(y) dy dx + \int_{-\infty}^0 f_X(x) \int_{z/X}^\infty f_Y(y) dy dx \right] \\
 &= \int_0^\infty f_X(x) \left[ f_Y(z/x) \left( \frac{1}{x} \right) - f(-\infty) \right] dx + \int_{-\infty}^0 f_X(x) \left[ f_Y(\infty) - f_Y(z/x) \left( \frac{1}{x} \right) \right] dx
 \end{aligned}$$

We obtained the above equation using the fundamental theorem of calculus and the chain rule. Now, we will use the fact that if  $f(x)$  is a distribution function, as  $x \rightarrow \infty$ ,  $f(x) \rightarrow 0$ . The same holds true as  $x \rightarrow -\infty$ . This yields,

$$\begin{aligned}
 f_Z(z) &= \int_0^\infty f_X(x) \left[ f_Y(z/x) \left( \frac{1}{x} \right) - f(-\infty) \right] dx + \int_{-\infty}^0 f_X(x) \left[ f_Y(\infty) - f_Y(z/x) \left( \frac{1}{x} \right) \right] dx \\
 &= \int_0^\infty f_X(x) f_Y(z/x) \frac{1}{x} dx + \int_{-\infty}^0 f_X(x) (-f_Y(z/x)) \frac{1}{x} dx \\
 &= \int_0^\infty f_X(x) f_Y(z/x) \frac{1}{x} dx - \int_{-\infty}^0 f_X(x) f_Y(z/x) \frac{1}{x} dx \\
 &= \int_0^\infty f_X(x) f_Y(z/x) \frac{1}{|x|} dx + \int_{-\infty}^0 f_X(x) f_Y(z/x) \frac{1}{|x|} dx \\
 &= \int_{-\infty}^\infty f_X(x) f_Y(z/x) \frac{1}{|x|} dx
 \end{aligned}$$

Now we need to find the mean of  $Z$  where  $Z = XY$ ,

$$\begin{aligned}
E[Z] &= \int_{-\infty}^{\infty} z f_Z(z) dz \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (xy) f_X(x) f_Y(y) \frac{1}{|x|} dx dy \\
&= \mu_Y \int_{-\infty}^{\infty} \frac{x}{|x|} f_X(x) dx \\
&= \mu_Y \left[ \int_{-\infty}^0 -f_X(x) dx + \int_0^{\infty} f_X(x) dx \right] \\
&= \mu_Y \left[ - \int_{-\infty}^0 f_X(x) dx + \int_0^{\infty} f_X(x) dx \right]
\end{aligned}$$

FINISH THIS LATER.

### Problem 9.

We have that,

$$\begin{aligned}
t_j - t_{j-1} &= (j/N)(b-a) + a - [(j-1/N)(b-a) + a] \\
&= (jb - ja)/N + a - (jb - ja - b + a)/N - a \\
&= (jb - ja - jb + ja + b - a)/N \\
&= (b-a)/N
\end{aligned}$$

In addition, since the function  $f$  is bounded on  $[a, b]$ , we know that  $\exists M \in \mathbb{N}$  such that  $M \geq |f|$ . These two facts yield the following,

$$\begin{aligned}
\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p &= \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})((b-a)/N)^p \\
&\leq \lim_{N \rightarrow \infty} \sum_{j=1}^N M((b-a)/N)^p \\
&= \lim_{N \rightarrow \infty} MN((b-a)/N)^p \\
&= M \lim_{N \rightarrow \infty} N((b-a)/N)^p \\
&= M(b-a)^p \lim_{N \rightarrow \infty} N/N^p \\
&= M(b-a)^p \lim_{N \rightarrow \infty} N^{1-p}
\end{aligned}$$

Since  $1-p < 0$ , we have that  $\lim_{N \rightarrow \infty} N^{1-p} = 0$ . Thus, this gives us,

$$\begin{aligned}
\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p &\leq M(b-a)^p \lim_{N \rightarrow \infty} N^{1-p} \\
&= [M(b-a)]0 = 0
\end{aligned}$$

Now, if we use  $-M$  for the lower bound, we get,

$$\begin{aligned}
\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p &\geq -M(b-a)^p \lim_{N \rightarrow \infty} N^{1-p} \\
&= [-M(b-a)]0 = 0
\end{aligned}$$



Since we have shown that

$$0 \leq \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p \leq 0$$

we have that  $\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p = 0$ .

**Problem 10.**

We know, by Proposition 4.9.1, that for a given stochastic process  $X_t$ , if  $E[X_t] \rightarrow k$  a constant and  $\text{Var}(X_t) \rightarrow 0$  as  $t \rightarrow \infty$ , then  $\text{ms-}\lim_{t \rightarrow \infty} X_t = k$ .

Thus, we will need to check the mean and variance of the given function. We will begin with the mean. Using the fact that the function  $f$  is a constant in terms of the expected value function, we get

$$E\left(\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(\Delta_{j-1}^j W)^2\right) = \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})E[(\Delta_{j-1}^j W)^2]$$

We know that the expected value of the square of a random variable is its variance, which is  $t_j - t_{j-1}$  in this case. Hence,

$$\begin{aligned} E\left(\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(\Delta_{j-1}^j W)^2\right) &= \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})E[(\Delta_{j-1}^j W)^2] \\ &= \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1}) \end{aligned}$$

This is the equation for the Riemann integral of  $f$  over  $[a, b]$ . Since we already know  $f$  is bounded and defined on a closed interval, as long as  $f$  is continuous almost everywhere on  $[a, b]$  we assert that the above limit exists and is equal to the integral of  $f$ . If that is the case, then we have

$$\begin{aligned} E\left(\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(\Delta_{j-1}^j W)^2\right) &= \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1}) \\ &= \int_a^b f(t)dt \end{aligned}$$

Now we need to show that the variance of the function is 0. Using Problem 8 to show that  $\text{Var}[(\Delta_{j-1}^j W)^2] = 2(t_j - t_{j-1})^2$ , we have

$$\begin{aligned} \text{Var}\left(\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(\Delta_{j-1}^j W)^2\right) &= \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 \text{Var}[(\Delta_{j-1}^j W)^2] \\ &= \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 \cdot 2(t_j - t_{j-1})^2 \\ &= 2 \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 (t_j - t_{j-1})^2 \end{aligned}$$

Now observe that if  $M$  is an upper bound for  $|f|$  on  $[a, b]$  (as in Problem 9), then  $M^2$  is an upper bound for  $|f|^2$  on  $[a, b]$ . This is true because increasing functions preserve inequalities, squaring is an increasing function for  $x \geq 0$ , and  $|f| \geq 0$  for every  $x$ . In addition, observe that  $|f|^2 = f^2$ . Thus, we have that  $M^2$  is an upper bound for  $f^2$  on  $[a, b]$ .

We can now use the above and the result from Problem 9 in order to obtain the following,

$$\begin{aligned}\text{Var}\left(\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(\Delta_{j-1}^j W)^2\right) &= 2 \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 (t_j - t_{j-1})^2 \\ &\leq 2M^2 \lim_{N \rightarrow \infty} \sum_{j=1}^N (t_j - t_{j-1})^2 \\ &= 2M^2(0) = 0\end{aligned}$$

In addition, since we know that variance is always  $\geq 0$ , we have that,

$$\text{Var}\left(\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(\Delta_{j-1}^j W)^2\right) = 0$$

Thus, we have shown that this function satisfies Proposition 4.9.1 and can state that  $\text{ms-}\lim_{t \rightarrow \infty} [\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(\Delta_{j-1}^j W)^2] = \int_a^b f(t) dt$ .

### Problem 11.

We need to show that the mean and variance of this function are 0. We'll begin with the mean,

$$E\left[\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p (\Delta_{j-1}^j W)^q\right] = \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p E[(\Delta_{j-1}^j W)^q]$$

If  $q$  is odd, then  $E[(\Delta_{j-1}^j W)^q] = 0$ . If  $q$  is even, then  $E[(\Delta_{j-1}^j W)^q] = \sigma^q (q-1)!! = (\sqrt{t_j - t_{j-1}})^p (p-1)!!$  where  $!!$  is the double factorial. Since the expected value of the whole function is trivially 0 if  $q$  is odd, we will only consider the situation where  $q$  is even,

$$\begin{aligned}\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p E[(\Delta_{j-1}^j W)^q] &= \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p (\sqrt{t_j - t_{j-1}})^q (q-1)!! \\ &= \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p (t_j - t_{j-1})^{q/2} (q-1)!! \\ &= (q-1)!! \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^{p+q/2}\end{aligned}$$

By the constraints in the problem, we know that  $p + q/2 > 1$ . Hence, we can apply the result of Problem 9. Hence, the limit is equal to 0,

$$\begin{aligned}E\left[\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p (\Delta_{j-1}^j W)^q\right] &= (q-1)!! \lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^{p+q/2} \\ &= (q-1)!!(0) \\ &= 0\end{aligned}$$

Now we must show that the variance is equal to 0,

$$\begin{aligned}\text{Var}\left[\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p (\Delta_{j-1}^j W)^q\right] &= \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 (t_j - t_{j-1})^{2p} \cdot \text{Var}[(\Delta_{j-1}^j W)^q] \\ &= \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 (t_j - t_{j-1})^{2p} \cdot E[(\Delta_{j-1}^j W)^{2q}] \\ &= \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 (t_j - t_{j-1})^{2p} \cdot E[(\Delta_{j-1}^j W)^{2q}] \\ &= \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 (t_j - t_{j-1})^{2p} \cdot (t_j - t_{j-1})^q (2q-1) \\ &= (2q-1) \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 (t_j - t_{j-1})^{2p+q}\end{aligned}$$

By the same reasoning as in Problem 10, this limit is equal to 0. Thus,

$$\begin{aligned} \text{Var}[\lim_{N \rightarrow \infty} \sum_{j=1}^N f(t_{j-1})(t_j - t_{j-1})^p (\Delta_{j-1}^j W)^q] &= (2q-1) \lim_{N \rightarrow \infty} \sum_{j=1}^N (f(t_{j-1}))^2 (t_j - t_{j-1})^{2p+q} \\ &= 0 \end{aligned}$$

Since the mean is equal to a constant (0 in this case) and the variance is equal to 0, we can say that the mean-squared limit of this function is 0.

**Problem 12.**

We have that,

$$\begin{aligned} \sum_{k=m}^n f_k(g_{k+1} - g_k) &= f_m(g_{m+1} - g_m) + f_{m+1}(g_{m+2} - g_{m+1}) + f_{m+2}(g_{m+3} - g_{m+2}) + \cdots \\ &= f_m(g_{m+1}) - f_m(g_m) + f_{m+1}(g_{m+2}) - f_{m+1}(g_{m+1}) + \cdots \\ &= -f_m(g_m) + g_{m+1}(f_m - f_{m+1}) + g_{m+2}(f_{m+1} - f_{m+2}) + \cdots + f_n(g_{n+1}) \\ &= f_n(g_{n+1}) - f_m(g_m) - [g_{m+1}(f_m - f_{m+1}) + g_{m+2}(f_{m+1} - f_{m+2}) + \cdots + g_n(f_{n-1} - f_n)] \\ &= f_n(g_{n+1}) - f_m(g_m) - \sum_{k=m+1}^n g_k(f_{k-1} - f_k) \end{aligned}$$

**Problem 13.**

We have  $F_t = W_t$  with the interval of interest being  $[0, T]$ . The partial sums of the integral  $\int_0^T W_t dW_t$  are given by

$$\begin{aligned} S_n &= \sum_{i=0}^{n-1} F_{t_i}(W_{t_{i+1}} - W_{t_i}) \\ &= \sum_{i=0}^{n-1} W_{t_i}(W_{t_{i+1}} - W_{t_i}) \end{aligned}$$

We also have that

$$W_{t_i}(W_{t_{i+1}} - W_{t_i}) = \frac{1}{2}W_{t_{i+1}}^2 - \frac{1}{2}W_{t_i}^2 - \frac{1}{2}(W_{t_{i+1}} - W_{t_i})^2$$

Plugging this into the partial sums formula yields,

$$\begin{aligned} S_n &= \frac{1}{2} \sum_{i=0}^{n-1} W_{t_{i+1}}^2 - \frac{1}{2} \sum_{i=0}^{n-1} W_{t_i}^2 - \frac{1}{2} \sum_{i=0}^{n-1} (W_{t_{i+1}} - W_{t_i})^2 \\ &= \frac{1}{2} W_{t_n}^2 - \frac{1}{2} \sum_{i=0}^{n-1} (W_{t_{i+1}} - W_{t_i})^2 \end{aligned}$$

If we let  $t_n = T$ , we get

$$S_n = \frac{1}{2} W_T^2 - \frac{1}{2} \sum_{i=0}^{n-1} (W_{t_{i+1}} - W_{t_i})^2$$

Using Proposition 4.11.6 and the fact that  $\frac{1}{2} W_T^2$  does not depend on  $n$ , we have

$$\begin{aligned} \text{ms-}\lim_{n \rightarrow \infty} S_n &= \frac{1}{2} W_T^2 - \text{ms-}\lim_{n \rightarrow \infty} \frac{1}{2} \sum_{i=0}^{n-1} (W_{t_{i+1}} - W_{t_i})^2 \\ &= \frac{1}{2} W_T^2 - \frac{1}{2} T \end{aligned}$$

Since the Ito integral is defined as the mean-squared limit of the partial sums  $S_n$ , we have that

$$\begin{aligned}\int_0^T W_t dW_t &= \text{ms-}\lim_{n \rightarrow \infty} S_n \\ &= \frac{1}{2}W_T^2 - \frac{1}{2}T\end{aligned}$$

as required.

**Problem 14.**

We are given the Ito integral,

$$\int_0^T f(t) dW_t$$

where  $f(t)$  is an arbitrary bounded and continuous function. Each increment is given by,

$$f(t_i)(W_{t_{i+1}} - W_{t_i})$$

So the partial sums are given by

$$S_n = \sum_{i=0}^{n-1} f(t_i)(W_{t_{i+1}} - W_{t_i})$$

We need to find the mean-squared limit of these partial sums. We'll start by finding the expected value,

$$\begin{aligned}E[\sum_{i=0}^{n-1} f(t_i)(W_{t_{i+1}} - W_{t_i})] &= \sum_{i=0}^{n-1} f(t_i)E[(W_{t_{i+1}} - W_{t_i})] \\ &= \sum_{i=0}^{n-1} f(t_i)(0) = 0\end{aligned}$$

Thus, the mean is 0. Now to find the variance,

$$\begin{aligned}\text{Var}[\sum_{i=0}^{n-1} f(t_i)(W_{t_{i+1}} - W_{t_i})] &= \sum_{i=0}^{n-1} f(t_i)^2 \text{Var}[(W_{t_{i+1}} - W_{t_i})] \\ &= \sum_{i=0}^{n-1} f(t_i)^2 (t_{i+1} - t_i)\end{aligned}$$

Now when we take the limit as  $n \rightarrow \infty$  of the variance, we get,

$$\lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} f(t_i)^2 (t_{i+1} - t_i) = \int_0^T f(t)^2$$

Thus, if  $Y_t = \int_0^T f(t) dW_t$ , then  $Y_t \sim N(0, \int_0^T f(t)^2)$ .

**Problem 15.**

Let  $Z_t = \int_0^t W_s ds$ . We will use integration by parts to solve this integral.

Let  $u = W_s$  and  $dv = ds$ . This implies that  $du = dW_s$  and  $v = t - s$ . Then using the formula,

$$\int u dv = uv - \int v du$$

we get,

$$\int_0^t W_s ds = W_s(t - s) - \int (t - s) dW_s$$

FINISH THIS LATER.

**Problem 16.**

We have  $X_t = \int_0^t (a + \frac{bu}{t}) dW_u$ . Since the non-infinitesimal part of the integral does not depend on a random variable, this is a Wiener integral.

By Proposition 5.6.1 , we know that Wiener integrals are normal random variables with mean 0 and variance,

$$\begin{aligned} \int_0^t (a + \frac{bu}{t})^2 du &= \int_0^t (a^2 + a\frac{bu}{t} + \frac{b^2u^2}{t^2}) du \\ &= a^2t + a\frac{bt^2}{2t} + \frac{b^2t^3}{3t^2} \\ &= a^2t + a\frac{bt}{2} + \frac{b^2t}{3} \\ &= (a^2 + \frac{ab}{2} + \frac{b^2}{3})t \end{aligned}$$

So, in order for the variance to equal 1, we require,

$$\begin{aligned} (a^2 + \frac{ab}{2} + \frac{b^2}{3})t &= 1 \\ \implies a^2 + \frac{ab}{2} + \frac{b^2}{3} &= \frac{1}{t} \end{aligned}$$

**Problem 17.**