

# Chapter 8 - Vector and Cartesian Spaces

Chris Hayduk

February 13, 2020

## 1 Exercises

**Problem 8.A.** If  $V$  is a vector space and if  $x + z = x$  for some  $x$  and  $z$  in  $V$ , show that  $z = 0$ . Hence, the zero element in  $V$  is unique.

Suppose  $x + z = x$ . Then by the properties of addition on a vector space, we have,

$$\begin{aligned}x + z &= x \\ \iff x + z + (-x) &= x + (-x) \\ \iff x + (-x) + z &= 0 \\ \iff 0 + z &= 0 \\ \iff z &= 0\end{aligned}$$

All of these statements follow directly from the definition of a vector space (Definition 8.1 in the text).

**Problem 8.B.** If  $x + y = 0$  for some  $x$  and  $y$  in  $V$ , show that  $y = -x$ .

Again using the properties of addition on a vector space, we have,

$$\begin{aligned}x + y &= 0 \\ \iff x + y + (-x) &= 0 + (-x) \\ \iff x + (-x) + y &= -x \\ \iff 0 + y &= -x \\ \iff y &= -x\end{aligned}$$

**Problem 8.C.** Let  $S = \{1, 2, \dots, p\}$  for some  $p \in \mathbb{N}$ . Show that the vector space  $\mathbb{R}^S$  is “essentially the same” as the space  $\mathbb{R}^p$ .

We know from Example 8.2d in the text that  $\mathbb{R}^S$  denotes the collection of all functions  $u$  with domain  $S$  and range in  $\mathbb{R}$ .

We assert that there is a bijection between  $\mathbb{R}^p$  and  $\mathbb{R}^S$  defined using the functions in  $\mathbb{R}^S$ . Namely,  $f : \mathbb{R}^S \rightarrow \mathbb{R}^p$  with  $f(u) = (u(1), u(2), \dots, u(k))$ .

First we will show that  $f$  is injective. So suppose  $u, v \in \mathbb{R}^S$  and  $x \in \mathbb{R}^p$  with  $f(u) = x$  and  $f(v) = x$ . That is,  $u(k) = x_k$  and  $v(k) = x_k$  for every  $k$ . Then clearly  $u(k) = v(k)$  for every  $k$  and hence the functions are equal.

Now to show that  $f$  is surjective. Let  $x \in \mathbb{R}^p$ . Then  $x = (x_1, x_2, \dots, x_p)$ .

We know that  $\mathbb{R}^S$  denotes the collection of all functions  $u$  with domain  $S$  and range in  $\mathbb{R}$ , and we know that  $x_k \in \mathbb{R}$  for every  $k$ . Hence, there clearly exists a function  $u \in \mathbb{R}^S$  such that  $u(k) = x_k$  for every  $k$ . Thus,  $f$  is surjective.

Now we need to show that  $f$  preserves addition and scalar multiplication. That is,  $f(u + v) = f(u) + f(v)$  and  $f(cu) = cf(u)$ .

Let  $u, v \in \mathbb{R}^S$  with  $f(u) = x$  and  $f(v) = y$ . Then,

$$\begin{aligned} f(u + v) &= (u(1) + v(1), u(2) + v(2), \dots, u(p) + v(p)) \\ &= (x_1 + y_1, \dots, x_p + y_p) \\ &= (x_1, \dots, x_p) + (y_1, \dots, y_p) \\ &= x + y \\ &= f(u) + f(v) \end{aligned}$$

So  $f$  preserves addition. Now to check scalar multiplication:

$$\begin{aligned} f(cu) &= (cu(1), \dots, cu(p)) \\ &= c(u(1), \dots, u(p)) \\ &= cf(u) \end{aligned}$$

So  $f$  preserves scalar multiplication.

Since  $f$  is a bijection between the elements of  $\mathbb{R}^S$  and  $\mathbb{R}^p$  which preserves addition and scalar multiplication, these two vector spaces are isomorphic (ie. “essentially the same”).

**Problem 8.D.** If  $w_1$  and  $w_2$  are strictly positive, show that the definition  $(x_1, x_2) \cdot (y_1, y_2) = x_1y_1w_1 + x_2y_2w_2$  yields an inner product on  $\mathbb{R}^2$ . Generalize this to  $\mathbb{R}^p$ .

We need to show that this potential norm satisfies the five conditions of Definition 8.3 in the text:

1.

$$\begin{aligned} x \cdot x &= (x_1, x_2) \cdot (x_1, x_2) = x_1x_1w_1 + x_2x_2w_2 \\ &= x_1^2w_1 + x_2^2w_2 \\ &\geq 0 \end{aligned}$$

2.

$$\begin{aligned}
& (x_1, x_2) \cdot (x_1, x_2) = 0 \\
& \iff x_1^2 w_1 + x_2^2 w_2 = 0 \\
& \iff x_1^2 w_1 = -x_2^2 w_2
\end{aligned}$$

We know that  $x_1^2 w_1 \geq 0$  and  $x_2^2 w_2 \geq 0$ . Hence,  $x_1^2 w_1 = -x_2^2 w_2$  iff  $x_1^2 w_1 = 0 = x_2^2 w_2$ .

3.

$$\begin{aligned}
x \cdot y &= (x_1, x_2) \cdot (y_1, y_2) = x_1 y_1 w_1 + x_2 y_2 w_2 \\
&= y_1 x_1 w_1 + y_2 x_2 w_2 \\
&= (y_1, y_2) \cdot (x_1, x_2) \\
&= y \cdot x
\end{aligned}$$

4.

$$\begin{aligned}
x \cdot (y + z) &= (x_1, x_2) \cdot (y_1 + z_1, y_2 + z_2) = x_1(y_1 + z_1)w_1 + x_2(y_2 + z_2)w_2 \\
&= (x_1 y_1 + x_1 z_1)w_1 + (x_2 y_2 + x_2 z_2)w_2 \\
&= x_1 y_1 w_1 + x_1 z_1 w_1 + x_2 y_2 w_2 + x_2 z_2 w_2 \\
&= (x_1 y_1 w_1 + x_2 y_2 w_2) + (x_1 z_1 w_1 + x_2 z_2 w_2) \\
&= (x_1, x_2) \cdot (y_1, y_2) + (x_1, x_2) \cdot (z_1, z_2) \\
&= x \cdot y + x \cdot z
\end{aligned}$$

and,

$$\begin{aligned}
(x + y) \cdot z &= (x_1 + y_1, x_2 + y_2) \cdot (z_1, z_2) = (x_1 + y_1)z_1 w_1 + (x_2 + y_2)z_2 w_2 \\
&= (x_1 z_1 + y_1 z_1)w_1 + (x_2 z_2 + y_2 z_2)w_2 \\
&= x_1 z_1 w_1 + y_1 z_1 w_1 + x_2 z_2 w_2 + y_2 z_2 w_2 \\
&= (x_1 z_1 w_1 + x_2 z_2 w_2) + (y_1 z_1 w_1 + y_2 z_2 w_2) \\
&= (x_1, x_2) \cdot (z_1, z_2) + (y_1, y_2) \cdot (z_1, z_2) \\
&= x \cdot z + y \cdot z
\end{aligned}$$

5.

$$\begin{aligned}
(ax) \cdot y &= (ax_1, ax_2) \cdot (y_1, y_2) \\
&= ax_1 y_1 w_1 + ax_2 y_2 w_2 \\
&= a(x_1 y_1 w_1 + x_2 y_2 w_2) = a(x \cdot y) \\
&= x_1 (ay_1) w_1 + x_2 (ay_2) w_2 \\
&= x \cdot (ay)
\end{aligned}$$

Hence, the proposed norm satisfies all of the necessary properties and thus defines a valid norm on  $\mathbb{R}^2$ .

In order to generalize this norm to  $\mathbb{R}^p$ , we need to fix  $w \in \mathbb{R}^p$  such that each component of  $w$  is strictly positive. Then the corresponding norm is given by,

$$x \cdot y = \sum_{i=1}^p x_i y_i w_i$$

**Problem 8.E.** The definition  $(x_1, x_2) \cdot (y_1, y_2) = x_1 y_1$  is *not* an inner product on  $\mathbb{R}^2$ . Why?

Let  $x = (0, 1)$ . Then  $x \cdot x = 0(0) = 0$ , but  $x \neq (0, 0)$ . Hence, property 2 of Definition 8.3 is violated and this does not define a valid norm.

**Problem 8.F.** If  $x = (x_1, x_2, \dots, x_p) \in \mathbb{R}^p$ , define  $\|x\|_1$  by  $\|x\|_1 = |x_1| + |x_2| + \dots + |x_p|$ . Prove that  $x \rightarrow \|x\|_1$  is a norm on  $\mathbb{R}^p$ .

We need to show that this proposed norm satisfies all four properties of Definition 8.5 in the text.

1. We have that  $\|x\|_1 = |x_1| + |x_2| + \dots + |x_p|$  for every  $x \in \mathbb{R}^p$ . Note that, for each  $x_k \in \mathbb{R}$ ,  $|x_k| \geq 0$ . Hence, the sum of these terms must also be greater than or equal to 0.

As a result, we have that  $\|x\| \geq 0$  for every  $x \in \mathbb{R}^p$ .

2. Suppose  $\|x\|_1 = 0$ . Then  $|x_1| + |x_2| + \dots + |x_p| = 0$ . Since each  $|x_k| \geq 0$  for every  $k$ , the sum of the terms is 0 iff each term is 0. Hence,  $x = 0$ .

Now suppose  $x = 0$ . Then  $\|x\|_1 = |x_1| + |x_2| + \dots + |x_p| = |0| + \dots + |0| = 0$ .

3. Let  $a \in \mathbb{R}$  and  $x \in \mathbb{R}^p$ . Then,

$$\begin{aligned} \|ax\|_1 &= |ax_1| + |ax_2| + \dots + |ax_p| \\ &= |a| \cdot |x_1| + |a| \cdot |x_2| + \dots + |a| \cdot |x_p| \\ &= |a| \cdot (|x_1| + \dots + |x_p|) \\ &= |a| \cdot \|x\|_1 \end{aligned}$$

4. Let  $x, y \in \mathbb{R}^p$ . We have that,

$$\|x + y\| = |x_1 + y_1| + |x_2 + y_2| + \dots + |x_p + y_p|$$

By Theorem 5.12 in the text, we know that  $|x_k + y_k| \leq |x_k| + |y_k|$  since  $x_k, y_k \in \mathbb{R}$  for every  $k$ . Hence, we have,

$$\begin{aligned} \|x + y\| &\leq |x_1| + |y_1| + |x_2| + |y_2| + \dots + |x_p| + |y_p| \\ &= |x_1| + \dots + |x_p| + |y_1| + \dots + |y_p| \\ &= \|x\| + \|y\| \end{aligned}$$

Thus, all of the properties hold and this defines a valid norm on  $\mathbb{R}^p$ .

**Problem 8.G.** If  $x = (x_1, x_2, \dots, x_p) \in \mathbb{R}^p$ , define  $\|x\|_\infty$  by  $\|x\|_\infty = \sup\{|x_1|, |x_2|, \dots, |x_p|\}$ . Prove that  $x \rightarrow \|x\|_\infty$  is a norm on  $\mathbb{R}^p$ .

Once again, we need to show that this proposed norm satisfies all four properties of Definition 8.5 in the text.

- 1.
- 2.
- 3.
- 4.

## 2 Project