

# Probability and Statistics (EPRST)

## Lecture 6

# Transformations of random variables - an example

## Example

*Say  $X \sim U(0, 1)$ . What is the distribution of  $Y = X^2$ ? Is it still uniform? What is its density?*

# Transformations of random variables - an example

## Example

*Let  $X \sim \mathcal{N}(0, 1)$ . What is the distribution of  $Y = X^2$ ?*

# Expectation

## Definition

*The expected value (the expectation or mean) of a random variable  $X$  is a number (denoted  $\mathbb{E}X$ ) given by*

$$\mathbb{E}X = \begin{cases} \sum_{x_i \in S} x_i \mathbb{P}(X = x_i), & \text{if } X \text{ is a discrete random variable,} \\ \int_{\mathbb{R}} x \cdot f(x) dx, & \text{if } X \text{ is continuous random variable} \end{cases}$$

*( $S$  denotes the set of values of  $X$  in the discrete case, and  $f$  stands for the density in the continuous case).*

*The expectation is well-defined if the series (or the integral) are absolutely convergent. If  $X \geq 0$  (that is,  $\mathbb{P}(X \geq 0) = 1$ ), then if the series (or the integral) diverge to  $\infty$  - then we define  $\mathbb{E}X := \infty$ .*

# The expectation - some examples

## Example

- If  $\mathbb{P}(X = -1) = 0.2$ ,  $\mathbb{P}(X = 0) = 0.4$ ,  $\mathbb{P}(X = 1) = 0.4$ , then  $\mathbb{E}X =$
- If  $\mathbb{P}(X = -1) = 0.4$ ,  $\mathbb{P}(X = 0) = 0.2$ ,  $\mathbb{P}(X = 1) = 0.4$ , then  $\mathbb{E}X =$
- If  $\mathbb{P}(X = -1) = 0.2$ ,  $\mathbb{P}(X = 0) = 0.4$ ,  $\mathbb{P}(X = 100) = 0.4$ , then  $\mathbb{E}X =$

## Example

Compute the mean of  $X \sim \mathcal{U}[0, 1]$ .

# The expectation - some examples

The expected value is not always finite:

## Example

*Let  $\mathbb{P}(X = k) = c/k^2$  for  $k \in \mathbb{N}$ . What is the value of  $c$ ? Does  $\mathbb{E}X$  exist?*

The expected value does not always exist:

## Example

*Let  $X$  - a random variable with a Cauchy distribution, that is, with the density*

$$f(x) = \frac{1}{\pi(1 + x^2)}, \quad x \in \mathbb{R}.$$

*Does  $\mathbb{E}X$  exist?*

## The expectation - some properties

- $\mathbb{E}c = c$  for every  $c \in \mathbb{R}$ ,
- $\mathbb{E}(c \cdot X) = c \cdot \mathbb{E}X$  for every  $c \in \mathbb{R}$ ,
- $\mathbb{E}(X_1 + \dots + X_n) = \mathbb{E}X_1 + \dots + \mathbb{E}X_n$
- if  $X \geq 0$ , then  $\mathbb{E}X \geq 0$ .

In particular, for all  $a, b \in \mathbb{R}$

$$\mathbb{E}(aX + b) = a\mathbb{E}X + b.$$

# The expectations of some important distributions

- if  $X \sim \text{bin}(n, p)$ , then  $\mathbb{E}X = ?$ ,
- if  $X \sim \text{geom}(p)$ , then  $\mathbb{E}X = ?$ ,
- if  $X \sim \text{Pois}(\lambda)$ , then  $\mathbb{E}X = ?$ ,
- if  $X \sim U(a, b)$ , then  $\mathbb{E}X = ?$ ,
- if  $X \sim \text{Exp}(\lambda)$ , then  $\mathbb{E}X = ?$ ,
- if  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\mathbb{E}X = ?$



# Sample mean

In statistics, a central problem is how to use data to estimate unknown parameters of a distribution. It is especially common to want to estimate the mean of a distribution (=the expected value of a random variable).

If the data are  $n$  values of a random variable  $X$  (generated independently), then the most natural way to estimate  $\mathbb{E}X$  is simply to average the values, taking the arithmetic mean. For example, if the observed data are 3, 1, 1, 5, then a simple, natural way to estimate the mean of the distribution that generated the data is to use

$$\frac{3 + 1 + 1 + 5}{4} = 2.5.$$

This is called the **sample mean**. The sample mean is an empirical estimate of the theoretical expected value (sometimes called the **population mean** or **true mean**) .

## Sample mean - cont'd

So if  $x_1, \dots, x_n$  are some (random) values of a random variable  $X$  (independently generated) then the sample mean is

$$\bar{X}_n := \frac{1}{n} \sum_{i=1}^n x_i.$$

### Example 1

We rolled a symmetric die 9 times and got: 3, 1, 2, 5, 1, 4, 3, 2, 6.  
The sample mean, computed from this particular sample is

$$\bar{X}_9 = \frac{1}{9}(3 + 1 + 2 + 5 + 1 + 4 + 3 + 2 + 6) = 3.$$

The population mean  $\mathbb{E}X = 3.5$ .

# The expectation of a function of random variable

If  $Y = g(X)$  then

$$\begin{aligned}\mathbb{E}(Y) &= \mathbb{E}g(X) = \\ &= \begin{cases} \sum_{x_i \in S} g(x_i) \cdot \mathbb{P}(X = x_i), & \text{discrete case,} \\ \int_{\mathbb{R}} g(x) \cdot f(x) dx, & \text{continuous case,} \end{cases}\end{aligned}$$

(if the series or the integral above are absolutely convergent).

As previously, if  $g(X) \geq 0$  (that is  $\mathbb{P}(g(X) \geq 0) = 1$ ), then if the series (or the integral) diverge to  $\infty$  - then we define  $\mathbb{E}g(X) = \infty$ .

# Some examples

## Example

*Suppose  $\mathbb{P}(X = -1) = \mathbb{P}(X = 1) = 1/4$ ,  $\mathbb{P}(X = 0) = 1/2$ . What is  $\mathbb{E}X^2$ ? Does it equal  $(\mathbb{E}X)^2$ ?*

## Example

*Compute  $\mathbb{E}X^2$  for  $X \sim U(0, 1)$ . Is it the same as  $(\mathbb{E}X)^2$ ?*

# The expectation of a function of random variable - some special cases

## Definition

- If  $g(x) = x^k$  for some  $k$ , then

$$\mathbb{E}g(X) = \mathbb{E}X^k$$

is called the  $k$ -th **moment** of  $X$ .

- If  $g(x) = (x - \mathbb{E}X)^2$ , then

$$\mathbb{E}g(X) = \mathbb{E}(X - \mathbb{E}X)^2$$

is called the **variance** of  $X$  (denoted  $\text{Var } X$ ).

- $\text{D } X = \sqrt{\text{Var } X}$  is the **standard deviation** of  $X$ .

# The variance - some properties

So the variance of  $X$  is defined as

$$\text{Var } X = \mathbb{E}(X - \mathbb{E}X)^2.$$

In order for  $\text{Var } X$  to exist, the second moment of  $X$  must be finite:  $\mathbb{E}X^2 < \infty$ .

Properties of the variance:

- $\text{Var } X = \mathbb{E}X^2 - (\mathbb{E}X)^2$ ,
- $\text{Var } X \geq 0$ ,
- $\text{Var } X = 0$  iff  $X$  has a one-point distribution,
- $\text{Var}(X + b) = \text{Var } X$  for every number  $b$ ,
- $\text{Var}(aX) = a^2 \text{Var } X$  for every number  $a$ .

In particular, for all  $a, b \in \mathbb{R}$

$$\text{Var}(aX + b) = a^2 \text{Var}(X).$$

## Computing the variance - some examples

- If  $\mathbb{P}(X = 1) = p = 1 - \mathbb{P}(X = 0)$ , then  $\text{Var } X = ?$
- If  $X \sim \text{U}(0, 1)$ , then  $\text{Var } X = ?$
- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\text{Var } X = ?$

## Computing the variance - some examples

Sort the variances of  $X$ ,  $Y$ ,  $Z$  and  $W$  in the increasing order, if

- $X$ , with the values:  $\{1, 2, 3, 4, 5\}$

$$\mathbb{P}(X = 1) = \dots = \mathbb{P}(X = 5) = \frac{1}{5},$$

- $Y$ , the values  $\{1, 2, 3, 4, 5\}$ :

$$\mathbb{P}(Y = 1) = \mathbb{P}(Y = 5) = \frac{1}{10}, \quad \mathbb{P}(Y = 2) = \mathbb{P}(Y = 4) = \frac{2}{10},$$
$$\mathbb{P}(Y = 3) = \frac{4}{10}$$

- $Z$ , the values  $\{1, 5\}$ :

$$\mathbb{P}(Z = 1) = \mathbb{P}(Z = 5) = \frac{1}{2},$$

- $W$ , the only value  $\{3\}$ :

$$\mathbb{P}(W = 3) = 1.$$



## Variances of the important named distributions

- if  $X \sim \text{bin}(n, p)$ , then  $\text{Var } X = np(1 - p)$
- if  $X \sim \text{geom}(p)$ , then  $\text{Var } X = (1 - p)/p^2$
- if  $X \sim \text{Pois}(\lambda)$ , then  $\text{Var } X = \lambda$
- if  $X \sim \text{U}(a, b)$ , then  $\text{Var } X = (b - a)^2/12$
- if  $X \sim \text{Exp}(\lambda)$ , then  $\text{Var } X = 1/\lambda^2$
- if  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\text{Var } X = \sigma^2$

## Sample variance and sample standard deviation

If  $x_1, \dots, x_n$  are some (random) values of a random variable  $X$  (independently generated) then a natural estimate of  $\mathbb{E}g(X)$  ( $g$  is any function) is the arithmetic mean of the values  $g(x_1), \dots, g(x_n)$ :

$$\frac{1}{n} \sum_{i=1}^n g(x_i).$$

However, the **sample variance** is defined as

$$S_n^2 := \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{X}_n)^2$$

The **sample standard deviation** is the square root of the sample variance.

## Sample variance - cont'd

The idea of the definition is to mimic the formula

$$\text{Var } X = \mathbb{E} (X - \mathbb{E}X)^2$$

by averaging the squared distances of the  $x_i$  from the sample mean, except with  $n - 1$  rather than  $n$  in the denominator.

The motivation for the  $n - 1$  is that this makes  $S_n^2$  **unbiased** for estimating  $\text{Var } X$ , that is it is correct on average.