

Numerical Methods

Laboratory no. 1

March 27, 2023 Krzysztof Watras

Computer Science

Warsaw University of Technology

Documentation of laboratory work results

Task 1

Given a function

$$y = \frac{\cos(x)}{x^3} - x^2$$

we want to calculate it's T(x) coefficient. We will use the property

$$\sigma[\tilde{v}^k] = T^k \cdot \sigma[\tilde{x}^{k-1}] + \nu^k$$

In our function we use:

$$\begin{aligned}v_1 &= \cos(x) \\v_2 &= x^3 \\v_4 &= \frac{v_1}{v_2} \\v_4 &= x^2 \\y &= v_4 - v_4\end{aligned}$$

Therefore we obtain:

$$\begin{aligned}\tilde{v}_1 &= \cos(x(1 + \epsilon_1)) \\ \tilde{v}_2 &= (x(1 + \epsilon))^3 = x^3(1 + 3\epsilon_2) \\ \tilde{v}_3 &= \frac{\tilde{v}_1}{\tilde{v}_2}(1 + \epsilon_3) \\ \tilde{v}_4 &= (x(1 + \epsilon))^2 = x^2(1 + 2\epsilon_4) \\ \tilde{y} &= \tilde{v}_3 - \tilde{v}_4\end{aligned}$$

Let us focus on \tilde{v}_1 :

$$\cos(x(1 + \epsilon)) = \cos(x + x\epsilon) = \cos(x) \cos(x\epsilon) - \sin(x) \sin(x\epsilon)$$

Now we need to use property of trigonometric functions:

$$For \alpha \rightarrow 0, \sin(\alpha) \approx \alpha, \cos(\alpha) \approx 1 - \frac{\alpha^2}{2}$$

Using those properties:

$$\begin{aligned}
\cos(x(1+\epsilon)) &= \cos(x)\cos(x\epsilon) - \sin(x)\sin(x\epsilon) \\
&= \cos(x)\left(1 - \frac{(x\epsilon)^2}{2}\right) - \sin(x)x\epsilon \\
&= \cos(x) - \sin(x)x\epsilon \\
&= \cos(x) \cdot (1 - \tan(x)x\epsilon)
\end{aligned}$$

Now solve v_3 :

$$\begin{aligned}
\tilde{v}_3 &= \frac{\cos(x) \cdot (1 - x \tan(x)\epsilon_1)}{x^3(1 + 3\epsilon_2)}(1 + \epsilon_3) \\
&= \frac{\cos(x)}{x^3}(1 - x \tan(x)\epsilon_1) \cdot (1 + 3\epsilon_2)^{-1}(1 + \epsilon_3) \\
&= \frac{\cos(x)}{x^3}(1 - x \tan(x)\epsilon_1) \cdot (1 - 3\epsilon_2)(1 + \epsilon_3) \\
&= \frac{\cos(x)}{x^3}(1 - x \tan(x)\epsilon_1 - 3\epsilon_2)(1 + \epsilon_3) \\
&= \frac{\cos(x)}{x^3}(1 - x \tan(x)\epsilon_1 - 3\epsilon_2 + \epsilon_3)
\end{aligned}$$

Finally, we get:

$$\begin{aligned}
\tilde{y} &= \frac{\cos(x)}{x^3}(1 - x \tan(x)\epsilon_1 - 3\epsilon_2 + \epsilon_3) - x^2(1 + 2\epsilon_4) \\
&= \frac{\cos(x)}{x^3} - \frac{\cos(x)}{x^3}(x \tan(x)\epsilon_1 + 3\epsilon_2 - \epsilon_3) - x^2 - 2x^2\epsilon_4 \\
&= y - \frac{\cos(x)}{x^3}(x \tan(x)\epsilon_1 + 3\epsilon_2 - \epsilon_3) - 2x^2\epsilon_4 \\
&= y\left(1 + \left(-\frac{\cos(x)}{x^3}(x \tan(x)\epsilon_1 + 3\epsilon_2 - \epsilon_3) - 2x^2\epsilon_4\right)\frac{1}{y}\right) \\
&= y\left(1 + \left(-\frac{\sin(x)}{x^2}\epsilon_1 - 3\frac{\cos(x)}{x^3}\epsilon_2 - \frac{\cos(x)}{x^3}\epsilon_3 - 2x^2\epsilon_4\right)\frac{1}{y}\right)
\end{aligned}$$


Therefore,

$$T(x) = \frac{\left|-\frac{\sin(x)}{x^2}\right| + \left| -3\frac{\cos(x)}{x^3} \right| + \left| -\frac{\cos(x)}{x^3} \right| + \left| -2x^2 \right|}{\frac{\cos(x)}{x^3} + x^2}$$

This yields results different from the ones we got via calculating the error numerically. Function

$$T(x) = \frac{1}{\epsilon_{sim}} \left| \frac{y(\tilde{x}) - y(x)}{y(x)} \right|$$

Differs significantly from the one we got from analitical solution.
Graph that visualises the difference:



1 Task 2

ipsum dolor sit amet, officia excepteur ex fugiat reprehenderit enim labore culpa sint ad nisi Lorem pariatur mollit ex esse exercitation amet. Nisi anim cupidatat excepteur officia. Reprehenderit nostrud nostrud ipsum Lorem est aliquip amet voluptate voluptate dolor minim nulla est proident. Nostrud officia pariatur ut officia. Sit irure elit esse ea nulla sunt ex occaecat reprehenderit commodo officia dolor Lorem duis laboris cupidatat officia voluptate. Culpa proident adipisicing id nulla nisi laboris ex in Lorem sunt duis officia eiusmod. Aliqua reprehenderit commodo ex non excepteur duis sunt velit enim. Voluptate laboris sint cupidatat ullamco ut ea consectetur et est culpa et culpa duis.