


## Problem 1.


A block of mass  $M$  is on a horizontal table. There is a hole in the table, through which a bullet is shot into the block. The bullet is stopped completely and becomes embedded in the block. The bullet has mass  $m$  and speed  $v_1$ . You can assume that the penetration occurs quickly.



- a) Determine the speed of the block immediately after the collision.
- b) How far above the table does the block-and-bullet system travel?

## Problem 2.

A particle with mass  $m = 10.0$  kg has at time  $t = 0$  s a velocity of  $v_0 = 10.0$  m/s. The particle is affected by a time-dependent force  $F$ , shown in the figure below.



What is the particle's speed at time  $t = 10$  s?

- A) The speed is  $v = 7.50$  m/s
- B) The speed is  $v = 10.0$  m/s
- C) The speed is  $v = 15.0$  m/s
- D) The speed is  $v = 17.5$  m/s
- E) The speed is  $v = 20.0$  m/s
- F) The speed is  $v = 32.0$  m/s
- G) Do not know

## Problem 3.


Two blocks with masses  $M$  (small block) and  $2M$  (large block) are at rest on a smooth, horizontal table. The two blocks are connected to a compressed spring. Now, the spring expands and accelerates the two blocks.

Which of the following statements are correct?

- A) The two blocks receive the same amount of momentum from the spring
- B) The large block receives a larger amount of momentum from the spring than the small block.
- C) The small block receives a larger amount of momentum from the spring than the large block.
- D) The two blocks receive the same kinetic energy from the spring.
- E) The small block receives more kinetic energy from the spring than the large block.
- F) The large block receives more kinetic energy from the spring than the small block.
- G) Do not know

## Problem 4.

A ball collides with a stationary wall as shown in the figure. The ball's mass is 0.1 kg, and it arrives with a speed of  $v = 10$  m/s. The collision is elastic, and there is no force of friction between the ball and the wall. A high-speed-camera measures the contact between the ball and the wall to last  $\Delta t = 1.0 \cdot 10^{-3}$  s.



What is the magnitude of the *average* force,  $F_{av}$  on the ball from the wall in this time-frame?

- A)  $F_{av} \approx 5.0 \cdot 10^2$  N
- B)  $F_{av} \approx 8.7 \cdot 10^3$  N
- C)  $F_{av} \approx 1.0 \cdot 10^3$  N
- D)  $F_{av} \approx 1.7 \cdot 10^3$  N
- E)  $F_{av} \approx 2.0 \cdot 10^3$  N
- F) Do not know

## Problem 5.

Two blocks A and B move towards each other in a straight line on a smooth table. The blocks have the same speed  $u$ . Block A has mass  $m$  and moves to the right (positive direction), while block B has mass  $4m$  and moves to the left. There is Velcro at the front of each block, ensuring that the blocks undergo a completely inelastic collision.


- a) Determine the speed of the blocks after the collision.
- b) The total kinetic energy of the blocks after the collision is the following fraction of the total kinetic energy of the blocks before the collision:

- A) 0
- B)  $\frac{9}{25}$
- C) 1
- D)  $\frac{1}{4}$
- E)  $\frac{9}{10}$
- F) Do not know

## Problem 6.

Two blocks with masses  $M_1$  and  $M_2$  lie on a horizontal, smooth table. A bullet with speed  $v$  and mass  $m$  passes completely through block  $M_1$ . Immediately after, the bullet impacts block  $M_2$  and thereafter travels with  $M_2$ . After perforating  $M_1$ , the bullet has a speed of  $v_1$ . After the collisions,  $M_1$  and the body consisting of  $M_2$  and  $m$  both have a speed of  $v_2$ .

- a) Find an expression for  $v_1$  and  $v_2$ .



## Problem 7.

Two cars crash into each other on a road. After the collision, they stick together and move as a single body. The first car has mass 1500 kg and drives to the east with speed 25.0 m/s before the collision. The second car has mass 2500 kg and drives northwards with speed 20.0 m/s before the collision.

- a) Determine the magnitude and direction of the velocity of the cars after the collision.

Problem 8.

Two blocks A and B move towards each other in a straight line on a smooth table. The blocks have the same speed  $u$ . Block A has mass  $m$  and moves to the right (positive direction), while block B has mass  $4m$  and moves to the left. On the front of each block is a spring, which ensures that the blocks undergo an elastic collision.


The velocities of the blocks  $v_A$  and  $v_B$  after the collision are

- A)  $v_A = -\frac{11}{5}u$  and  $v_B = -\frac{1}{5}u$
- B)  $v_A = 0$  and  $v_B = 0$
- C)  $v_A = -\frac{23}{17}u$  and  $v_B = -\frac{7}{17}u$
- D)  $v_A = -u$  and  $v_B = u$
- E) Do not know



## Problem 9.

Three blocks, A, B and C can slide along a horizontal, smooth surface. The blocks have mass  $m$ ,  $2m$ , and  $m$ , respectively. Block A moves towards block B with a speed of  $9.0 \text{ m/s}$  and undergoes a central, elastic collision with B. Afterwards, B undergoes a central, completely inelastic collision with C.



What is the final speed of block C?

- A)  $1.0 \text{ m/s}$
- B)  $2.0 \text{ m/s}$
- C)  $3.0 \text{ m/s}$
- D)  $4.0 \text{ m/s}$
- E)  $5.0 \text{ m/s}$
- F)  $6.0 \text{ m/s}$
- G)  $7.0 \text{ m/s}$
- H)  $8.0 \text{ m/s}$
- I) Do not know