

On the nested p -center problem



Master Thesis Seminar

Christof Brandstetter, Supervisor: Univ.-Prof. DI Markus Sinnl, BSc PhD

2024-01-12

Introduction

- **p -center problem**: Open p facilities such that the maximum distance between any customer and its nearest open facility is minimized
- Related to the set cover problem and the assignment problem
- Nesting: Open additional facilities subsequently
- **Nested p -center problem**: Open p_h facilities in period h such that the sum of the maximum distances between any customer and its nearest open facility in period h is minimized.
- Use case: Ambulance/logistics stations, network design, screening/vaccination stations
- Optimizes expansion or retraction path

Research question

How can the nesting concept be applied to the p -center problem?

Research question

How can the nesting concept be applied to the p -center problem?

- proposed by McGarvey and Thorsen (2022)

Research question

How can the nesting concept be applied to the p -center problem?

- proposed by McGarvey and Thorsen (2022)
- Which formulations can be used for the nesting?

Research question

How can the nesting concept be applied to the p -center problem?

- proposed by McGarvey and Thorsen (2022)
- Which formulations can be used for the nesting?
- Which algorithms perform best in terms of solution time?

Research question

How can the nesting concept be applied to the p -center problem?

- proposed by McGarvey and Thorsen (2022)
- Which formulations can be used for the nesting?
- Which algorithms perform best in terms of solution time?
- How can the nested p -center problem affect managerial decisions?

Research question

How can the nesting concept be applied to the p -center problem?

- proposed by McGarvey and Thorsen (2022)
- Which formulations can be used for the nesting?
- Which algorithms perform best in terms of solution time?
- How can the nested p -center problem affect managerial decisions?

Definition of the nested p -center problem

Definition:

- given a set of customer demand points \mathcal{I} , potential facility locations \mathcal{J} , time periods $\mathcal{H} = \{1, \dots, H\}$, integers $\mathcal{P} = \{p^1, \dots, p^H\}$ where $p^h \leq p^{h+1}$ for $h = 1, \dots, H - 1$ and distances $d_{ij} \geq 0$ between each $i \in \mathcal{I}$ and $j \in \mathcal{J}$

Definition of the nested p -center problem

Definition:

- given a set of customer demand points \mathcal{I} , potential facility locations \mathcal{J} , time periods $\mathcal{H} = \{1, \dots, H\}$, integers $\mathcal{P} = \{p^1, \dots, p^H\}$ where $p^h \leq p^{h+1}$ for $h = 1, \dots, H - 1$ and distances $d_{ij} \geq 0$ between each $i \in \mathcal{I}$ and $j \in \mathcal{J}$
- open p^h -facilities at different locations $j \in \mathcal{J}$
- the sum of maximum distance of any customer demand i point to its closest opened facility in time period h is minimized
- and facilities open in period h has to be open in period $h + 1$ as well for $h = 1, \dots, H - 1$

Related work



p -center problem / TODO

- First introduction of the p -center problem by Hakimi (1964)
- The standard textbook formulation of the p -center problem can be found in Daskin (2013)
- Solution approach based on the set cover problem by Contardo, Iori, and Kramer (2018)
- A compact formulation by Ales and Elloumi (2018) and Elloumi, Labbé, and Pochet (2004)
- A fast projection-based branch-and-cut algorithm by Gaar and Sinnl (2022)

Nested facility location problems

- First introduction of the nesting property and constraint by Roodman and Schwarz (1975)
- Extension of the nesting to a phase-in and phase-out by Roodman and Schwarz (1977)
- Reintroduction of the nesting on the example of the p -median problem by McGarvey and Thorsen (2022)
- Recent work on multi-period facility locations problems mainly focus on varying demand, distances, or cost over time f.e. Calogiuri et al. (2021)

Mixed Integer Linear Programming formulations



First MILP formulation

\mathcal{I} ... set of customer demand points with $i \in \mathcal{I}$

\mathcal{J} ... set of potential facility locations with $j \in \mathcal{J}$

\mathcal{H} ... set of time periods, $\mathcal{H} = \{1, 2, \dots, H\}$

p_h = number of facilities to be opened in time period $h \in \mathcal{H}$ with $p_h > p_{h-1}$

First MILP formulation

\mathcal{I} ... set of customer demand points with $i \in \mathcal{I}$

\mathcal{J} ... set of potential facility locations with $j \in \mathcal{J}$

\mathcal{H} ... set of time periods, $\mathcal{H} = \{1, 2, \dots, H\}$

p_h = number of facilities to be opened in time period $h \in \mathcal{H}$ with $p_h > p_{h-1}$

d_{ij} = distance between customer demand point i and potential facility location j

$x_{ijh} = \begin{cases} 1 & \dots \text{ if customer } i \text{ is assigned to facility } j \text{ in time period } h \\ 0 & \dots \text{ otherwise} \end{cases}$

$y_{jh} = \begin{cases} 1 & \dots \text{ if a facility is opened at location } j \text{ in time period } h \\ 0 & \dots \text{ otherwise} \end{cases}$

z_h = maximum distance between any customer i and its nearest open facility in period h

\mathcal{I} ... set of customer demand points i
 $[-1mm]\mathcal{J}$... set of potential facility locations j
 \mathcal{H} ... set of time periods, $\mathcal{H} = \{1, 2, \dots, H\}$
 \mathcal{P} ... set of integers of facilities to open, $\mathcal{P} = \{p^1, \dots, p^H\}$ where p^h for $h = 1, \dots, H - 1$
 d_{ij} = distance between customer demand point i and potential facility location j
 $x_{ij}^h = \begin{cases} 1 & \dots \text{ if a customer demand point } i \text{ is assigned to facility } j \text{ in period } h \\ 0 & \dots \text{ otherwise} \end{cases}$
 $y_j^h = \begin{cases} 1 & \dots \text{ if a facility is opened at location } j \text{ in period } h \\ 0 & \dots \text{ otherwise} \end{cases}$
 z^h ... maximum distance between any customer i and its nearest facility in period h

First MILP formulation

$$\begin{array}{ll}
 (nPC1) \quad \min & \sum_{h \in \mathcal{H}} z_h \quad \text{(objective)} \\
 \text{s.t.} & \sum_{j \in \mathcal{J}} y_{jh} = p_h \quad \forall h \in \mathcal{H} \quad \text{(facility)} \\
 & \sum_{j \in \mathcal{J}} x_{ijh} = 1 \quad \forall i \in \mathcal{I}, h \in \mathcal{H} \quad \text{(assignment)} \\
 & x_{ijh} \leq y_j h \quad \forall i \in \mathcal{I}, j \in \mathcal{J}, h \in \mathcal{H} \quad \text{(open)} \\
 & \sum_{j \in \mathcal{J}} d_{ij} x_{ijh} \leq z_h \quad \forall i \in \mathcal{I}, h \in \mathcal{H} \quad \text{(push)} \\
 & y_{jh} \geq y_{jh-1} \quad \forall h \in \mathcal{H} \setminus \{1\} \quad \text{(nesting)} \\
 & x_{ijh}, y_{jh} \in \{0, 1\} \quad \forall i \in \mathcal{I}, j \in \mathcal{J}, h \in \mathcal{H} \quad \text{(binary)} \\
 & z_h \in \mathbb{R}_{\geq 0} \quad \forall h \in \mathcal{H} \quad \text{(non-negativity)}
 \end{array}$$

Second MILP formulation

\mathcal{I} ... set of customer demand points i

\mathcal{J} ... set of potential facility locations j

\mathcal{H} ... set of time periods, $\mathcal{H} = \{1, 2, \dots, H\}$

\mathcal{P} ... set of integers of facilities to open, $\mathcal{P} = \{p^1, \dots, p^H\}$ where $p^h > p^{h+1}$
for $h = 1, \dots, H - 1$

d_{ij} = distance between customer demand point i and potential facility location j

$y_j^h = \begin{cases} 1 & \dots \text{ if a facility is opened at location } j \text{ in period } h \\ 0 & \dots \text{ otherwise} \end{cases}$

z^h ... maximum distance between any customer i and its nearest open facility
in period h

Second MILP formulation

$$\begin{aligned}
 (nPC2) \quad & \min \sum_{h \in H} z_h && \text{(objective-y)} \\
 \text{s.t.} \quad & \sum_{j \in J} y_{jh} = p_h && \forall h \in H \quad \text{(facility-y)} \\
 & z_h \geq d_{ij} - \sum_{j': d_{ij'} < d_{ij}} (d_{ij} - d_{ij'}) y_{j'h} && \forall i \in I, j \in J, h \in H \quad \text{(push-y)} \\
 & y_h \geq y_{h-1} && \forall j \in J, h \in H \setminus \{1\} \quad \text{(nesting-y)} \\
 & y_{jh} \in \{0, 1\} && \forall j \in J, h \in H \quad \text{(binary-y)} \\
 & z_h \in \mathbb{R}_{\geq 0} && \forall h \in H \quad \text{(non-negativity-y)}
 \end{aligned}$$

Third formulation

Constraint lifting

Theorem [Brandstetter(2023)]


Let LB_h being a lower bound on the decision variable z_h of (nPC1) for every $i \in I, j \in J, h \in H$ then

$$\sum_{j \in J} \max\{LB_h, d_{ij}\} x_{ijh} \leq z_h \quad \forall i \in I, h \in H \quad (\text{lift})$$


is a valid equality. Theorem is based on Gaar and Sinnl (2022).

- also valid for the y-formulation
- using various techniques to obtain LB_h


Constraint lifting / Delete




$p = 2$
 $d_{ij} = 0$
 $d_{ij} = 4$ if not indicated




Constraint lifting / Delete




$p = 2$
 $d_{ij} = 0$
 $d_{ij} = 4$ if not indicated
 $LB = 2.833$




Constraint lifting / Delete




$p = 2$
 $d_{ij} = 0$
 $d_{ij} = 4$ if not indicated
 $LB = 2.833$




Constraint lifting / Delete




$p = 2$
 $d_{ij} = 0$
 $d_{ij} = 4$ if not indicated
 $LB = 2.877$



Constraint lifting / Delete



$p = 2$
 $d_{ij} = 0$
 $d_{ij} = 4$ if not indicated
 $LB = 2.877$



Branch-and-cut / Delete

- adds the cutting plane method

Branch-and-cut / Delete

- adds the cutting plane method
- adding cut(s) \rightarrow branching \rightarrow adding cut(s) \rightarrow branching

Branch-and-cut / Delete

- adds the cutting plane method
- adding cut(s) \rightarrow branching \rightarrow adding cut(s) \rightarrow branching
- start with lower number of (push) constraints

Branch-and-cut / Delete

- adds the cutting plane method
- adding cut(s) \rightarrow branching \rightarrow adding cut(s) \rightarrow branching
- start with lower number of (push) constraints
- add (lift) inequalities

Branch-and-cut / Delete

- adds the cutting plane method
- adding cut(s) \rightarrow branching \rightarrow adding cut(s) \rightarrow branching
- start with lower number of (push) constraints
- add (lift) inequalities
- allows for shorter solving times

Outline of the results / This should be the main topic for the final presentation



Data

- data set **PMED** from "A note on solving large p -median problems" by Beasley (1985)
 - set of 40 instances
 - between 100 and 900 nodes

Data

- data set **PMED** from "A note on solving large p -median problems" by Beasley (1985)
 - set of 40 instances
 - between 100 and 900 nodes
- data set **TSPLIB** 2D-Euclidean distances from "TSPLIB – A Traveling Salesman Problem Library" by Reinelt (1991)
 - set of 80 instance
 - between 51 and 18512 nodes
 - rounded to the nearest integer value

Comparison

- comparing the results between (nPC1) and (nPC2)

Comparison

- comparing the results between (nPC1) and (nPC2)
- comparing the different lifting methods

Comparison

- comparing the results between (nPC1) and (nPC2)
- comparing the different lifting methods
- analysing the solutions of the nested and comparing it to the standard p -center problem

Comparison

- comparing the results between (nPC1) and (nPC2)
- comparing the different lifting methods
- analysing the solutions of the nested and comparing it to the standard p -center problem
- analysing the results regarding managerial insights

On the nested p -center problem



Master Thesis Seminar

Christof Brandstetter, Supervisor: Univ.-Prof. DI Markus Sinnl, BSc PhD

2024-01-12

Further research topics

- u -space model introduced by Elloumi, Labbé, and Pochet (2004)
- maximal regret as objective function
- trying to improve the performance of the nested p -median problem?

References I

- [1] Zacharie Ales and Sourour Elloumi. Compact MILP Formulations for the p -Center Problem. In: Combinatorial Optimization. Ed. by Jon Lee, Giovanni Rinaldi, and A. Ridha Mahjoub. Cham: Springer International Publishing, 2018, pp. 14–25. ISBN: 978-3-319-96151-4.
- [2] J.E. Beasley. A note on solving large p -median problems. In: European Journal of Operational Research 21.2 (1985), pp. 270–273. ISSN: 0377-2217. DOI: [https://doi.org/10.1016/0377-2217\(85\)90040-2](https://doi.org/10.1016/0377-2217(85)90040-2).

References II

- [3] Tobia Calogiuri et al. The multi-period p -center problem with time-dependent travel times. In: Computers & Operations Research 136 (2021), p. 105487. ISSN: 0305-0548. DOI: <https://doi.org/10.1016/j.cor.2021.105487>. URL: <https://www.sciencedirect.com/science/article/pii/S0305054821002343>.
- [4] Claudio Contardo, Manuel Iori, and Raphael Kramer. A scalable exact algorithm for the vertex p -center problem. In: Computers & Operations Research 103 (Mar. 2018). DOI: 10.1016/j.cor.2018.11.006.

References III

- [5] Mark S. Daskin. Center Problems. In: Network and Discrete Location: Models, Algorithms, and Applications, Second Edition. John Wiley & Sons, Ltd, 2013. Chap. 5, pp. 193–234. ISBN: 9781118537015. DOI: <https://doi.org/10.1002/9781118537015.ch05>.
- [6] Sourour Elloumi, Martine Labbé, and Yves Pochet. A New Formulation and Resolution Method for the p -Center Problem. In: INFORMS Journal on Computing 16 (Feb. 2004), pp. 83–94. DOI: 10.1287/ijoc.1030.0028.

References IV

- [7] Elisabeth Gaar and Markus Sinnl. A scaleable projection-based branch-and-cut algorithm for the p -center problem. In: European Journal of Operational Research 303.1 (2022), pp. 78–98. ISSN: 0377-2217. DOI: <https://doi.org/10.1016/j.ejor.2022.02.016>.
- [8] S. L. Hakimi. Optimum Locations of Switching Centers and the Absolute Centers and Medians of a Graph. In: Operations Research 12.3 (1964), pp. 450–459. DOI: 10.1287/opre.12.3.450.
- [9] Ronald McGarvey and Andreas Thorsen. Nested-Solution Facility Location Models. In: Optimization Letters 16 (Mar. 2022). DOI: 10.1007/s11590-021-01759-4.

References V

- [10] Gerhard Reinelt. TSPLIB — A Traveling Salesman Problem Library. In: ORSA Journal on Computing 3.4 (1991), pp. 376–384. DOI: 10.1287/ijoc.3.4.376.
- [11] Gary M. Roodman and Leroy B. Schwarz. Extensions of the Multi-Period Facility Phase-Out Model: New Procedures and Application to a Phase-In/Phase-Out Problem. In: A I E Transactions 9.1 (1977), pp. 103–107. DOI: 10.1080/05695557708975128.
- [12] Gary M. Roodman and Leroy B. Schwarz. Optimal and Heuristic Facility Phase-out Strategies. In: A I E Transactions 7.2 (1975), pp. 177–184. DOI: 10.1080/05695557508975000.