

# On the nested $p$ -center problem



Christof Brandstetter, Markus Sinnl

Institute of Business Analytics and Technology Transformation / JKU Business School,  
Johannes Kepler University Linz

2024-03-05

# The $p$ -center problem $p$ CP: Definition

## Definition

- Given

# The $p$ -center problem $p$ CP: Definition

## Definition

- Given
  - a set  $V$  of locations,

# The $p$ -center problem $p$ CP: Definition

## Definition

- Given
  - a set  $V$  of locations,
  - $p \in \mathbb{Z}$ , and

# The $p$ -center problem $p$ CP: Definition

## Definition

- Given
  - a set  $V$  of locations,
  - $p \in \mathbb{Z}$ , and
  - distances  $d_{ij}$  from location  $i \in V$  to  $j \in V$

# The $p$ -center problem $p$ CP: Definition

## Definition

- Given
  - a set  $V$  of locations,
  - $p \in \mathbb{Z}$ , and
  - distances  $d_{ij}$  from location  $i \in V$  to  $j \in V$
- we want to
  - open  $p$  locations of  $V$ , such that

# The $p$ -center problem $p$ CP: Definition

## Definition

- Given
  - a set  $V$  of locations,
  - $p \in \mathbb{Z}$ , and
  - distances  $d_{ij}$  from location  $i \in V$  to  $j \in V$
- we want to
  - open  $p$  locations of  $V$ , such that
  - the maximum distance of any location to its closest opened location is minimized.

# The nesting property

## Definition

- Given



# The nesting property

## Definition

- Given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and

# The nesting property

## Definition

- Given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open

# The nesting property

## Definition

- Given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$

# The nesting property

## Definition

- Given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$
- the subsets  $V^h \subseteq V$  with  $|V^h| = p^h$

# The nesting property

## Definition

- Given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$
- the subsets  $V^h \subseteq V$  with  $|V^h| = p^h$
- are called **nested** iff  $V^h \subset V^{h+1}$  for  $h = 1, \dots, H-1$


# The nesting property

## Definition

- Given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$
- the subsets  $V^h \subseteq V$  with  $|V^h| = p^h$
- are called **nested** iff  $V^h \subset V^{h+1}$  for  $h = 1, \dots, H-1$

Introduced by Roodman and Schwarz 1975 and used in e.g., Albareda-Sambola et al. 2009; Bakker and Nickel 2024; Conforti et al. 2014 and reintroduced as nesting by McGarvey and Thorsen 2022

# $p$ -center problem vs nested $p$ -center problem for $\mathcal{P} = \{4, 5, 6\}$



# The nested $p$ -center problem: Definition

## Definition part I

- given a set of locations  $V$ ,



# The nested $p$ -center problem: Definition

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,

# The nested $p$ -center problem: Definition

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$

# The nested $p$ -center problem: Definition

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$   
and
  - $p^H \leq |V|$

# The nested $p$ -center problem: Definition

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$   
and
  - $p^H \leq |V|$
- distances  $d_{ij} \geq 0$  between each  $i \in V$   
and  $j \in V$

# The nested $p$ -center problem: Definition

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$   
and
  - $p^H \leq |V|$
- distances  $d_{ij} \geq 0$  between each  $i \in V$   
and  $j \in V$

## Definition part II

- a feasible solution to the nested  $p$ -center problem consists of a set  $\mathcal{J}^h \subseteq V$

# The nested $p$ -center problem: Definition

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$   
and
  - $p^H \leq |V|$
- distances  $d_{ij} \geq 0$  between each  $i \in V$   
and  $j \in V$

## Definition part II

- a feasible solution to the nested  $p$ -center problem consists of a set  $\mathcal{J}^h \subseteq V$ 
  - with  $|\mathcal{J}^h| = p^h$  for  $h \in \mathcal{H}$ ,
  - for which  $\mathcal{J}^h \subseteq \mathcal{J}^{h+1}$  for  $h = 1, \dots, H-1$  holds

# The nested $p$ -center problem: Definition

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$  and
  - $p^H \leq |V|$
- distances  $d_{ij} \geq 0$  between each  $i \in V$  and  $j \in V$

## Definition part II

- a feasible solution to the nested  $p$ -center problem consists of a set  $\mathcal{J}^h \subseteq V$ 
  - with  $|\mathcal{J}^h| = p^h$  for  $h \in \mathcal{H}$ ,
  - for which  $\mathcal{J}^h \subseteq \mathcal{J}^{h+1}$  for  $h = 1, \dots, H-1$  holds
- the goal is to find a feasible solution which minimizes  $\sum_{h=1}^H d_h(\mathcal{J}^h)$ ,
  - where  $d_h(\mathcal{J}) = \max_{i \in \mathcal{I}} \min_{j \in \mathcal{J}} d_{ij}$  for  $h \in \mathcal{H}$

# The nested $p$ -center problem: Potential applications

- variant of the  $p$ -center problem



# The nested $p$ -center problem: Potential applications

- variant of the  $p$ -center problem
- part of the multi-period facility locations

# The nested $p$ -center problem: Potential applications

- variant of the  $p$ -center problem
- part of the multi-period facility locations
- emergency medical services

# The nested $p$ -center problem: Potential applications

- variant of the  $p$ -center problem
- part of the multi-period facility locations
- emergency medical services
- relief actions in humanitarian crisis

# The nested $p$ -center problem: Potential applications

- variant of the  $p$ -center problem
- part of the multi-period facility locations
- emergency medical services
- relief actions in humanitarian crisis
- expansion into new markets/areas

# Mixed Integer Linear Programming (MILP) Formulations



# The nested $p$ -center problem: MILP formulation

Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

# The nested $p$ -center problem: MILP formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

# The nested $p$ -center problem: MILP formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

## (nPC2) based on Gaar and Sinnl 2022

$$\min \sum_{h \in \mathcal{H}} z^h \quad (1a)$$

$$\text{s.t.} \quad \sum_{j \in V} y_j^h = p^h \quad \forall h \in \mathcal{H} \quad (1b)$$

$$z^h \geq d_{ij} - \sum_{j': d_{ij'} < d_{ij}} (d_{ij} - d_{ij'}) y_{j'}^h \quad \forall i, j \in V, h \in \mathcal{H} \quad (1c)$$

$$y_j^h \geq y_j^{h-1} \quad \forall j \in V, \forall h \in \mathcal{H} \setminus \{1\} \quad (1d)$$

$$(y, z) \in |V| |\mathcal{H}| \times \mathbb{R}_{\geq 0} \quad (1e)$$



# The nested $p$ -center problem: Compact MILP formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

# The nested $p$ -center problem: Compact MILP formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

# The nested $p$ -center problem: Compact MILP formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

$$u_k^h \dots \begin{cases} 1 \dots \text{if } z^h \geq D_k \text{ in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

# The nested $p$ -center problem: Compact MILP formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

$$u_k^h \dots \begin{cases} 1 \dots \text{if } z^h \geq D_k \text{ in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

(nPC3) based on Ales and Elloumi 2018

$$\begin{aligned} \min \quad & \sum_{h \in \mathcal{H}} z^h \\ \text{s.t.} \quad & \sum_{j \in V} y_j^h = p^h \quad \forall h \in \mathcal{H} \\ & D_0 + \sum_{k=1}^K (D_k - D_{k-1}) u_k^h \leq z^h \quad \forall h \in \mathcal{H} \\ & u_k^h + \sum_{j: d_{ij} < D_k} y_j^h \geq 1 \quad \forall i \in V, \forall h \in \mathcal{H}, \forall k \in S_i \cup \{K\} \\ & u_k^h \geq u_{k+1}^h \quad \forall h \in \mathcal{H}, \forall k \in \mathcal{K} \setminus \{K\} \\ & y_j^h \geq y_j^{h-1} \quad \forall h \in \mathcal{H} \setminus \{1\} \\ & (u, y, z) \in |V| \cdot |K| \times |V| \cdot |\mathcal{H}| \times \mathbb{R}_{\geq 0} \end{aligned}$$

# Comparing formulations

Table: Comparison of complexity of the formulations

	(nPC1)	(nPC2)	(nPC3)
Variables	$\mathcal{O}( \mathcal{I}  \mathcal{J}  \mathcal{H} )$	$\mathcal{O}( \mathcal{J}  \mathcal{H} )$	$\mathcal{O}(( \mathcal{I}  +  \mathcal{K} ) \mathcal{H} )$
Constraints	$\mathcal{O}( \mathcal{I}  \mathcal{J}  \mathcal{H} )$	$\mathcal{O}( \mathcal{I}  \mathcal{J}  \mathcal{H} )$	$\mathcal{O}(\min( \mathcal{I}  \mathcal{J} ,  \mathcal{I}  \mathcal{K} ) \mathcal{H} )$

# Comparing formulations

Table: Comparison of complexity of the formulations

	(nPC1)	(nPC2)	(nPC3)
Variables	$\mathcal{O}( \mathcal{I}  \mathcal{J}  \mathcal{H} )$	$\mathcal{O}( \mathcal{J}  \mathcal{H} )$	$\mathcal{O}(( \mathcal{I}  +  \mathcal{K} ) \mathcal{H} )$
Constraints	$\mathcal{O}( \mathcal{I}  \mathcal{J}  \mathcal{H} )$	$\mathcal{O}( \mathcal{I}  \mathcal{J}  \mathcal{H} )$	$\mathcal{O}(\min( \mathcal{I}  \mathcal{J} ,  \mathcal{I}  \mathcal{K} ) \mathcal{H} )$

The  $p$ CP version of formulation (nPC3) has the best known linear programming  $LP$ -bounds for the  $p$ CP, while the  $p$ CP versions of (nPC1) and (nPC2) have equally but worse  $LP$ -bounds.

# Improving the formulations



# Strengthening constraints (nPC2)



# Strengthening constraints (nPC2)

## *Lemma 1*

*Let  $LB_h$  be a lower bound on the decision variable  $z_h$  of (nPC2) for every  $i \in \mathcal{I}, j \in \mathcal{J}, h \in \mathcal{H}$  then*

# Strengthening constraints (nPC2)

## Lemma 1

Let  $LB_h$  be a lower bound on the decision variable  $z_h$  of (nPC2) for every  $i \in \mathcal{I}, j \in \mathcal{J}, h \in \mathcal{H}$  then

$$z^h \geq \max\{LB^h, d_{ij}\} - \sum_{j': d_{ij'} < d_{ij}} \left( \max\{LB^h, d_{ij}\} - \max\{LB^h, d_{ij'}\} \right) y_{j'}^h \quad (\text{nL-OPT})$$

# Strengthening constraints (nPC2)

## Lemma 1

Let  $LB_h$  be a lower bound on the decision variable  $z_h$  of (nPC2) for every  $i \in \mathcal{I}, j \in \mathcal{J}, h \in \mathcal{H}$  then

$$z^h \geq \max\{LB^h, d_{ij}\} - \sum_{j': d_{ij'} < d_{ij}} \left( \max\{LB^h, d_{ij}\} - \max\{LB^h, d_{ij'}\} \right) y_{j'}^h \quad (\text{nL-OPT})$$

is a valid inequality for (nPC2), i.e., every feasible solution of (nPC2) fulfills (nL-OPT). The theorem is based on Lemma 5 in Gaar and Sinnl (2022).

## Reducing set $\mathcal{K}$ in (nPC3)

## Reducing set $\mathcal{K}$ in (nPC3)

### *Lemma 2*

*Let  $\underline{z}^h$  be a valid lower bound and  $\overline{z}^h$  be a valid upper bound on the decision variable  $z^h$  for  $h \in \mathcal{H}$ , then the distinct distance  $D_k$  can only be the optimal distance for  $z^h$  if  $\underline{z}^h \leq D^k \leq \overline{z}^h$  holds.*

## Reducing set $\mathcal{K}$ in (nPC3)

### Lemma 2

*Let  $\underline{z}^h$  be a valid lower bound and  $\overline{z}^h$  be a valid upper bound on the decision variable  $z^h$  for  $h \in \mathcal{H}$ , then the distinct distance  $D_k$  can only be the optimal distance for  $z^h$  if  $\underline{z}^h \leq D^k \leq \overline{z}^h$  holds.*

*Therefore let set  $S_i^h \subseteq S_i$  for  $h \in \mathcal{H}$ , where  $S_i^h$  contains only the indices  $k$  where  $\underline{z}^h \leq D^k \leq \overline{z}^h$  holds.*

Depending on the bounds  $\underline{z}^h$  and  $\overline{z}^h$  the sets  $S_i^h$  can be much smaller than  $S_i$ .

# Reducing set I

## Observation 1

Let  $\underline{z}^h$  be a valid lower bound and  $\overline{z}^h$  be a valid upper bound on the decision variable  $z^h$  for  $h \in \mathcal{H}$ , then the distinct distance  $D_k$  can only be the optimal distance for  $z^h$  if  $\underline{z}^h \leq D^k \leq \overline{z}^h$  holds.

## Observation 2

Following Observation 1 we observe that decision variables  $u_h^k = 0$  for  $k : D_k > \overline{z}^h$  and  $u_h^k = 1$  for  $k : D_k < \underline{z}^h$ .

# Reducing set II

## Observation 3

*Following Observation 2 we observe that for  $k$  where variables  $u_k^h$  have been set to either zero or one, the variables do not have to be created.*



# Obtaining bounds

# Obtaining bounds

## *Lemma 3*

*Let  $z'^{h*}$  be the optimal objective function value of  $pCP$  with  $p = p^h$  for  $h \in \mathcal{H} = \{1, 2, \dots, H\}$  where  $p^h > p^{h+1}$ ,*

# Obtaining bounds

## Lemma 3

*Let  $z'^{h*}$  be the optimal objective function value of  $pCP$  with  $p = p^h$  for  $h \in \mathcal{H} = \{1, 2, \dots, H\}$  where  $p^h > p^{h+1}$ , then  $UB = Hz'^{1*}$  is a valid upper bound on the optimal objective function value of  $n$ - $pCP$ .*

# Obtaining bounds

## Lemma 3

Let  $z'^{h*}$  be the optimal objective function value of pCP with  $p = p^h$  for  $h \in \mathcal{H} = \{1, 2, \dots, H\}$  where  $p^h > p^{h+1}$ , then  $UB = Hz'^{1*}$  is a valid upper bound on the optimal objective function value of n-pCP.

Then

$$\overline{z^h} = \frac{UB - \sum_{x=h+1}^H z'^{x*}}{h}$$

# Obtaining bounds

## Lemma 3

Let  $z'^{h*}$  be the optimal objective function value of pCP with  $p = p^h$  for  $h \in \mathcal{H} = \{1, 2, \dots, H\}$  where  $p^h > p^{h+1}$ , then  $UB = Hz'^{1*}$  is a valid upper bound on the optimal objective function value of n-pCP.

Then

$$\overline{z^h} = \frac{UB - \sum_{x=h+1}^H z'^{x*}}{h}$$

where  $\overline{z^h}$  is a valid upper bound on the decision variable  $z^h$  of the n-pCP for  $h \in \mathcal{H}$ .

# Implementation



# Implementation

- C++, CPLEX 20.1

# Implementation

- C++, CPLEX 20.1
- preprocessing



# Implementation

- C++, CPLEX 20.1
- preprocessing
  - Solving  $p$ -center problem with  $p = p^H$
  - Use optimal solution value for  $p$ -center problem with  $p = p^{H-1}$
  - Repeat for remaining  $p^h$
- (nPC2)

# Implementation

- C++, CPLEX 20.1
- preprocessing
  - Solving  $p$ -center problem with  $p = p^H$
  - Use optimal solution value for  $p$ -center problem with  $p = p^{H-1}$
  - Repeat for remaining  $p^h$
- (nPC2)
  - branch-and-cut algorithm
  - separation similar to fixedCustomer from Gaar and Sinnl 2022

# Implementation

- C++, CPLEX 20.1
- preprocessing
  - Solving  $p$ -center problem with  $p = p^H$
  - Use optimal solution value for  $p$ -center problem with  $p = p^{H-1}$
  - Repeat for remaining  $p^h$
- (nPC2)
  - branch-and-cut algorithm
  - separation similar to fixedCustomer from Gaar and Sinnl 2022
- (nPC3)

# Implementation

- C++, CPLEX 20.1
- preprocessing
  - Solving  $p$ -center problem with  $p = p^H$
  - Use optimal solution value for  $p$ -center problem with  $p = p^{H-1}$
  - Repeat for remaining  $p^h$
- (nPC2)
  - branch-and-cut algorithm
  - separation similar to fixedCustomer from Gaar and Sinnl 2022
- (nPC3)
  - branch-and-bound algorithm
  - starting with the reduced set  $\mathcal{K}$

# Settings

- noPP: without preprocessing or any lifting

# Settings

- noPP: without preprocessing or any lifting
- PP: with preprocessing

# Settings

- noPP: without preprocessing or any lifting
- PP: with preprocessing
  - (nPC2): set upper bounds on  $z^h$  and using lower bounds in (nL-OPT)
  - (nPC3): problem initialized on reduced set  $\mathcal{K}$

# Settings

- noPP: without preprocessing or any lifting
- PP: with preprocessing
  - (nPC2): set upper bounds on  $z^h$  and using lower bounds in (nL-OPT)
  - (nPC3): problem initialized on reduced set  $\mathcal{K}$
- PH: PP with starting heuristic



# Results



# Instance from literature

- computational setup
  - single core of Intel Xeon X5570 with 2.93 GHz and 8 GB RAM
  - **timelimit** of 3600 seconds


# Instance from literature

- computational setup
  - single core of Intel Xeon X5570 with 2.93 GHz and 8 GB RAM
  - **timelimit** of 3600 seconds
- instance set **PMED** Çalık and Tansel 2013; D. Chen and R. Chen 2009; Contardo, Iori, and Kramer 2019
  - 40 instances
  - $\mathcal{P} = \{p, p + 1, p + 2\}$ ,  $p$  from 5 to 200,  $|V|$  from 100 and 900 nodes


# Instance from literature

- computational setup
  - single core of Intel Xeon X5570 with 2.93 GHz and 8 GB RAM
  - **timelimit** of 3600 seconds
- instance set **PMED** Çalık and Tansel 2013; D. Chen and R. Chen 2009; Contardo, Iori, and Kramer 2019
  - 40 instances
  - $\mathcal{P} = \{p, p+1, p+2\}$ ,  $p$  from 5 to 200,  $|V|$  from 100 and 900 nodes
- instance set **TSPLIB** Contardo, Iori, and Kramer 2019; Gaar and Sinnl 2022
  - 80 instances
  - $\mathcal{P} = \{4, 5, 6\}$ ,  $|V|$  from 51 and 1002 nodes

# Setting comparison on formulation (nPC1)




(a) runtime




(b) optimality gap [ $\frac{UB-LB}{UB}$ ]

# Setting comparison on formulation (nPC2)




(c) runtime




(d) optimality gap

# Setting comparison on formulation (nPC3)




(e) runtime




(f) optimality gap

# Formulation comparison on setting PH




(g) runtime



(h) optimality gap




# Managerial insights




(i) Relative regret of the optimal solution value

# Managerial insights



(k) Relative regret of the optimal solution value



(l) Relative regret of # of opened facilities

# Conclusion

- (nPC3) best performance in runtimes and optimality gap
- The (nPC2) outperforms the (nPC1)
- Preprocessing brings a large speed up on all formulations
- Starting heuristic little effect, shows good upper bound obtained in preprocessing
- Maximal relative regret of the optimal objective function value of 15%
- Maximal relative regret of number of opened facilities above 150%

# On the nested $p$ -center problem



Christof Brandstetter, Markus Sinnl

Institute of Business Analytics and Technology Transformation / JKU Business School,

Johannes Kepler University Linz

2024-03-05

# References I

- [1] M. Albareda-Sambola et al. The multi-period incremental service facility location problem. In: Computers & Operations Research 36.5 (2009), pp. 1356–1375.
- [2] Z. Ales and S. Elloumi. Compact MILP formulations for the  $p$ -center problem. In: Combinatorial Optimization: 5th International Symposium, ISCO 2018, Marrakesh, Morocco, April 11–13, 2018, Revised Selected Papers 5. 2018, pp. 14–25.
- [3] H. Bakker and S. Nickel. The Value of the Multi-period Solution revisited: When to model time in capacitated location problems. In: Computers & Operations Research 161 (2024), p. 106428.

## References II

- [4] H. Çalık and B. C. Tansel. Double bound method for solving the  $p$ -center location problem. In: Computers & Operations Research 40.12 (2013), pp. 2991–2999.
- [5] D. Chen and R. Chen. New relaxation-based algorithms for the optimal solution of the continuous and discrete  $p$ -center problems. In: Computers & Operations Research 36.5 (2009), pp. 1646–1655.
- [6] M. Conforti et al. Integer programming models. Springer, 2014.
- [7] C. Contardo, M. Iori, and R. Kramer. A scalable exact algorithm for the vertex  $p$ -center problem. In: Computers & Operations Research 103 (2019), pp. 211–220.

## References III

- [8] M. S. Daskin. Network and Discrete Location: Models, Algorithms, and Applications, Second Edition. John Wiley & Sons, Ltd, 2013.
- [9] E. Gaar and M. Sinnl. A scaleable projection-based branch-and-cut algorithm for the  $p$ -center problem. In: European Journal of Operational Research 303.1 (2022), pp. 78–98.
- [10] R. G. McGarvey and A. Thorsen. Nested-solution facility location models. In: Optimization letters 16.2 (2022), pp. 497–514.
- [11] G. M. Roodman and L. B. Schwarz. Optimal and heuristic facility phase-out strategies. In: AIIE transactions 7.2 (1975), pp. 177–184.

# The nested $p$ -center problem: Classical MILP formulation

## Decision variables

$$x_{ij}^h \dots \begin{cases} 1 \dots \text{if location } i \text{ is assigned to} \\ \quad \text{location } j \text{ in time period } h \\ 0 \dots \text{otherwise} \end{cases}$$



# The nested $p$ -center problem: Classical MILP formulation

## Decision variables

$$x_{ij}^h \dots \begin{cases} 1 \dots \text{if location } i \text{ is assigned to} \\ \quad \text{location } j \text{ in time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \quad \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

# The nested $p$ -center problem: Classical MILP formulation

## Decision variables

$$x_{ij}^h \dots \begin{cases} 1 \dots \text{if location } i \text{ is assigned to} \\ \quad \text{location } j \text{ in time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \quad \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$$z^h \dots \text{maximum distance between any} \\ \text{customer } i \text{ and its nearest open facility} \\ \text{in period } h$$

# The nested $p$ -center problem: Classical MILP formulation

## Decision variables

$$x_{ij}^h \dots \begin{cases} 1 \dots \text{if location } i \text{ is assigned to} \\ \text{location } j \text{ in time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$$z^h \dots \text{maximum distance between any} \\ \text{customer } i \text{ and its nearest open facility} \\ \text{in period } h$$

## (nPC1) based on Daskin 2013

$$\min \sum_{h \in \mathcal{H}} z^h \quad (3a)$$

$$\text{s.t. } \sum_{j \in V} y_j^h = p^h \quad \forall h \in \mathcal{H} \quad (3b)$$

$$\sum_{j \in V} x_{ij}^h = 1 \quad \forall i \in V, h \in \mathcal{H} \quad (3c)$$

$$x_{ij}^h \leq y_j^h \quad \forall i, j \in V, h \in \mathcal{H} \quad (3d)$$

$$\sum_{j \in V} d_{ij} x_{ij}^h \leq z^h \quad \forall i \in V, h \in \mathcal{H} \quad (3e)$$

$$y_j^h \geq y_j^{h-1} \quad \forall h \in \mathcal{H} \setminus \{1\} \quad (3f)$$

$$(x, y, z) \in |V|^2 |\mathcal{H}| \times |V| |\mathcal{H}| \times \mathbb{R}_{\geq 0} \quad (3g)$$