

# On the nested $p$ -center problem



Christof Brandstetter, Markus Sinnl

Institute of Business Analytics and Technology Transformation / JKU Business School,  
Johannes Kepler University Linz

This research was funded in whole, or in part, by the Austrian Science Fund (FWF) [P 35160-N].  
2024-03-06



JOHANNES KEPLER  
UNIVERSITY LINZ  
Altenberger Straße 69  
4040 Linz, Austria  
jku.at

# The $p$ -center problem ( $p$ CP):

## Definition

# The $p$ -center problem ( $p$ CP):

## Definition

- given
  - a set  $V$  of locations,

# The $p$ -center problem ( $p$ CP):

## Definition

- given
  - a set  $V$  of locations,
  - $p \in \mathbb{Z}$ , and

# The $p$ -center problem ( $p$ CP):

## Definition

- given
  - a set  $V$  of locations,
  - $p \in \mathbb{Z}$ , and
  - distances  $d_{ij}$  from location  $i \in V$  to  $j \in V$

# The $p$ -center problem ( $p$ CP):

## Definition

- given
  - a set  $V$  of locations,
  - $p \in \mathbb{Z}$ , and
  - distances  $d_{ij}$  from location  $i \in V$  to  $j \in V$
- we want to
  - open  $p$  locations of  $V$ , such that

# The $p$ -center problem ( $p$ CP):

## Definition

- given
  - a set  $V$  of locations,
  - $p \in \mathbb{Z}$ , and
  - distances  $d_{ij}$  from location  $i \in V$  to  $j \in V$
- we want to
  - open  $p$  locations of  $V$ , such that
  - the maximum distance of any location to its closest opened location is minimized.

# The nesting property

## Definition

- given
  - a set of locations  $V$



# The nesting property

## Definition

- given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and

# The nesting property

## Definition

- given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open

# The nesting property

## Definition

- given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$

# The nesting property

## Definition

- given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$
- the subsets  $V^h \subseteq V$  with  $|V^h| = p^h$

# The nesting property

## Definition


- given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$
- the subsets  $V^h \subseteq V$  with  $|V^h| = p^h$
- are called **nested** iff  $V^h \subseteq V^{h+1}$  for  $h = 1, \dots, H-1$

# The nesting property

## Definition

- given
  - a set of locations  $V$
  - a set of time periods  $\mathcal{H} = \{1, \dots, H\}$  and
  - a set of integers  $\mathcal{P} = \{p^1, \dots, p^H\}$  indicating the number of facilities to open
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$
- the subsets  $V^h \subseteq V$  with  $|V^h| = p^h$
- are called **nested** iff  $V^h \subseteq V^{h+1}$  for  $h = 1, \dots, H-1$

Introduced by Roodman and Schwarz 1975 and used in e.g., Albareda-Sambola et al. 2009; Bakker and Nickel 2024; Conforti et al. 2014 and reintroduced as nesting by McGarvey and Thorsen 2022



# The nested $p$ -center problem

## Definition part I

- given a set of locations  $V$ ,



# The nested $p$ -center problem

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,

# The nested $p$ -center problem

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$

# The nested $p$ -center problem

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$   
and
  - $p^H \leq |V|$

# The nested $p$ -center problem

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$   
and
  - $p^H \leq |V|$
- distances  $d_{ij} \geq 0$  between each  $i \in V$   
and  $j \in V$

# The nested $p$ -center problem

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$   
and
  - $p^H \leq |V|$
- distances  $d_{ij} \geq 0$  between each  $i \in V$   
and  $j \in V$

## Definition part II

- a feasible solution to the nested  $p$ -center problem consists of a set  $\mathcal{J}^h \subseteq V$

# The nested $p$ -center problem

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$   
and
  - $p^H \leq |V|$
- distances  $d_{ij} \geq 0$  between each  $i \in V$   
and  $j \in V$

## Definition part II

- a feasible solution to the nested  $p$ -center problem consists of a set  $\mathcal{J}^h \subseteq V$ 
  - with  $|V^h| = p^h$  for  $h \in \mathcal{H}$ ,
  - for which  $V^h \subseteq V^{h+1}$  for  $h = 1, \dots, H-1$  holds

# The nested $p$ -center problem

## Definition part I

- given a set of locations  $V$ ,
- time periods  $\mathcal{H} = \{1, \dots, H\}$ ,
- integers  $\mathcal{P} = \{p^1, \dots, p^H\}$ 
  - where  $p^h \leq p^{h+1}$  for  $h = 1, \dots, H-1$  and
  - $p^H \leq |V|$
- distances  $d_{ij} \geq 0$  between each  $i \in V$  and  $j \in V$

## Definition part II

- a feasible solution to the nested  $p$ -center problem consists of a set  $\mathcal{J}^h \subseteq V$ 
  - with  $|V^h| = p^h$  for  $h \in \mathcal{H}$ ,
  - for which  $V^h \subseteq V^{h+1}$  for  $h = 1, \dots, H-1$  holds
- the goal is to find a feasible solution which minimizes  $\sum_{h=1}^H d(V^h)$ ,
  - where  $d(V^h) = \max_{i \in V} \min_{j \in V^h} d_{ij}$  for  $h \in \mathcal{H}$

# Mixed Integer Linear Programming (MILP) Formulations





# First formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

# First formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

# First formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

## (nPC2) based on Gaar and Sinnl 2022

$$\min \sum_{h \in \mathcal{H}} z^h \quad (1a)$$

$$\text{s.t.} \quad \sum_{j \in V} y_j^h = p^h \quad \forall h \in \mathcal{H} \quad (1b)$$

$$z^h \geq d_{ij} - \sum_{j': d_{ij'} < d_{ij}} (d_{ij} - d_{ij'}) y_{j'}^h \quad \forall i, j \in V, h \in \mathcal{H} \quad (1c)$$

$$y_j^h \geq y_j^{h-1} \quad \forall j \in V, \forall h \in \mathcal{H} \setminus \{1\} \quad (1d)$$

$$(y, z) \in \mathbb{B}^{|V||\mathcal{H}|} \times \mathbb{R}_{\geq 0}^{|\mathcal{H}|} \quad (1e)$$

# Second formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

# Second formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

$$u_k^h \dots \begin{cases} 1 \dots \text{if } z^h \geq D_k \text{ in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

# Second formulation

## Decision variables

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

$$u_k^h \dots \begin{cases} 1 \dots \text{if } z^h \geq D_k \text{ in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

## (nPC3) based on Ales and Elloumi 2018

$$\begin{aligned} \min \quad & \sum_{h \in \mathcal{H}} z^h \\ \text{s.t.} \quad & \sum_{j \in V} y_j^h = p^h & \forall h \in \mathcal{H} \\ & D_0 + \sum_{k=1}^K (D_k - D_{k-1}) u_k^h \leq z^h \quad \forall h \in \mathcal{H} \\ & u_k^h + \sum_{j: d_{ij} < D_k} y_j^h \geq 1 & \forall i \in V, \forall h \in \mathcal{H}, \forall k \in S_i \cup \{K\} \\ & u_k^h \geq u_{k+1}^h & \forall h \in \mathcal{H}, \forall k \in \mathcal{K} \setminus \{K\} \\ & y_j^h \geq y_j^{h-1} & \forall h \in \mathcal{H} \setminus \{1\} \\ (u, y, z) \quad & \in \mathbb{B}^{|\mathcal{K}| |\mathcal{H}|} \times \mathbb{R}_{\geq 0}^{|\mathcal{H}|} \end{aligned}$$

# Improving the formulations



# Strengthening constraints (nPC2)



# Strengthening constraints (nPC2)

## *Lemma 1*

*Let  $\underline{z}^h$  be a lower bound on the decision variable  $z_h$  of (nPC2) for every  $i \in \mathcal{I}, j \in \mathcal{J}, h \in \mathcal{H}$  then constraints*

# Strengthening constraints (nPC2)

## Lemma 1

Let  $\underline{z}^h$  be a lower bound on the decision variable  $z_h$  of (nPC2) for every  $i \in \mathcal{I}, j \in \mathcal{J}, h \in \mathcal{H}$  then constraints

$$z^h \geq d_{ij} - \sum_{j': d_{ij'} < d_{ij}} (d_{ij} - d_{ij'}) y_{j'}^h$$

can be replaced by

# Strengthening constraints (nPC2)

## Lemma 1

Let  $\underline{z}^h$  be a lower bound on the decision variable  $z_h$  of (nPC2) for every  $i \in \mathcal{I}, j \in \mathcal{J}, h \in \mathcal{H}$  then constraints

$$z^h \geq d_{ij} - \sum_{j': d_{ij'} < d_{ij}} (d_{ij} - d_{ij'}) y_{j'}^h$$

can be replaced by

$$z^h \geq \max\{\underline{z}^h, d_{ij}\} - \sum_{j': d_{ij'} < d_{ij}} \left( \max\{\underline{z}^h, d_{ij}\} - \max\{\underline{z}^h, d_{ij'}\} \right) y_{j'}^h \quad (\text{nL-OPT})$$

The lemma is based on Lemma 5 in Gaar and Sinnl (2022).

# Reducing the number of variables $u_k^h$ in (nPC3)

# Reducing the number of variables $u_k^h$ in (nPC3)

## Observation 1

*Let  $\underline{z}^h$  be a valid lower bound and  $\overline{z}^h$  be a valid upper bound on the decision variable  $z^h$  for  $h \in \mathcal{H}$ , then the distinct distance  $D_k$  can only be the optimal distance for  $z^h$  if  $\underline{z}^h \leq D^k \leq \overline{z}^h$  holds.*

## Observation 2

*Following Observation 1, we observe that decision variables  $u_h^k = 0$  for  $k : D_k > \overline{z}^h$  and  $u_h^k = 1$  for  $k : D_k < \underline{z}^h$  in any optimal solution.*

# Reducing the number of variables $u_k^h$ in (nPC3)

## Observation 1

*Let  $\underline{z}^h$  be a valid lower bound and  $\overline{z}^h$  be a valid upper bound on the decision variable  $z^h$  for  $h \in \mathcal{H}$ , then the distinct distance  $D_k$  can only be the optimal distance for  $z^h$  if  $\underline{z}^h \leq D^k \leq \overline{z}^h$  holds.*

## Observation 2

*Following Observation 1, we observe that decision variables  $u_h^k = 0$  for  $k : D_k > \overline{z}^h$  and  $u_h^k = 1$  for  $k : D_k < \underline{z}^h$  in any optimal solution.*

The decision variables  $u_k^h$  which are zero or one following Observation 2, are redundant.

# Obtaining bounds

# Obtaining bounds

## Observation 3

*For  $\mathcal{H} = \{1\}$  the  $(n\text{-}p\text{CP})$  reduces to the  $(p\text{CP})$  where  $p = p^1$ , so the optimal objective value  $(z'^{h*})$  of the  $(p\text{CP})$ , where  $p = p^h$  is a lower bound  $\underline{z}^h$  on the decision variable  $z^h$  of the  $(n\text{-}p\text{CP})$ .*



# Obtaining bounds

## Observation 3

For  $\mathcal{H} = \{1\}$  the  $(n-pCP)$  reduces to the  $(pCP)$  where  $p = p^1$ , so the optimal objective value ( $z^{h*}$ ) of the  $(pCP)$ , where  $p = p^h$  is a lower bound  $\underline{z}^h$  on the decision variable  $z^h$  of the  $(n-pCP)$ .

## Proposition 1

Given an valid upper bound  $UB$  on the objective value of the  $(n-pCP)$  and valid lower bounds  $\underline{z}^h$  on the variable  $z^h$  can be obtained by the following equation:

$$\overline{z}^h = \frac{UB - \sum_{x=h+1}^H \underline{z}^h}{h} \quad (3)$$

# Implementation



# Implementation

- C++, CPLEX 20.1

# Implementation

- C++, CPLEX 20.1
- preprocessing
  - Solving  $p$ -center problem with  $p = p^H$
  - Use optimal solution value as lower bound for  $p$ -center problem with  $p = p^{H-1}$
  - Repeat for remaining  $p^h$  and calculate the upper bounds  $\overline{z}^h$

# Implementation

- C++, CPLEX 20.1
- preprocessing
  - Solving  $p$ -center problem with  $p = p^H$
  - Use optimal solution value as lower bound for  $p$ -center problem with  $p = p^{H-1}$
  - Repeat for remaining  $p^h$  and calculate the upper bounds  $\overline{z}^h$
- (nPC2)
  - branch-and-cut algorithm
  - separation based on the fixedCustomer separation scheme from Gaar and Sinnl 2022

# Implementation

- C++, CPLEX 20.1
- preprocessing
  - Solving  $p$ -center problem with  $p = p^H$
  - Use optimal solution value as lower bound for  $p$ -center problem with  $p = p^{H-1}$
  - Repeat for remaining  $p^h$  and calculate the upper bounds  $\overline{z}^h$
- (nPC2)
  - branch-and-cut algorithm
  - separation based on the fixedCustomer separation scheme from Gaar and Sinnl 2022
- (nPC3)

# Implementation

- C++, CPLEX 20.1
- preprocessing
  - Solving  $p$ -center problem with  $p = p^H$
  - Use optimal solution value as lower bound for  $p$ -center problem with  $p = p^{H-1}$
  - Repeat for remaining  $p^h$  and calculate the upper bounds  $\overline{z}^h$
- (nPC2)
  - branch-and-cut algorithm
  - separation based on the fixedCustomer separation scheme from Gaar and Sinnl 2022
- (nPC3)
  - branch-and-bound algorithm

# Settings

- noPP: without preprocessing and without lifting



# Settings

- noPP: without preprocessing and without lifting
- PP: with preprocessing

# Settings

- noPP: without preprocessing and without lifting
- PP: with preprocessing
  - (nPC2): set upper bounds on  $z^h$  and using lower bounds in (nL-OPT)
  - (nPC3): problem initialized on reduced number of  $u_k^h$

# Settings

- noPP: without preprocessing and without lifting
- PP: with preprocessing
  - (nPC2): set upper bounds on  $z^h$  and using lower bounds in (nL-OPT)
  - (nPC3): problem initialized on reduced number of  $u_k^h$
- PH: PP with starting heuristic

# Results



# Instance from literature

- instance set **PMED**
  - 40 instances
  - $\mathcal{P} = \{p, p + 1, p + 2\}$ ,  $p$  from 5 to 200,  $|V|$  from 100 and 900 nodes


# Instance from literature

- instance set **PMED**
  - 40 instances
  - $\mathcal{P} = \{p, p+1, p+2\}$ ,  $p$  from 5 to 200,  $|V|$  from 100 and 900 nodes
- instance set **TSPLIB**
  - 50 instances
  - $\mathcal{P} = \{4, 5, 6\}$ ,  $|V|$  from 51 and 1002 nodes


# Instance from literature

- instance set **PMED**
  - 40 instances
  - $\mathcal{P} = \{p, p + 1, p + 2\}$ ,  $p$  from 5 to 200,  $|V|$  from 100 and 900 nodes
- instance set **TSPLIB**
  - 50 instances
  - $\mathcal{P} = \{4, 5, 6\}$ ,  $|V|$  from 51 and 1002 nodes
- computational setup
  - single core of Intel Xeon X5570 with 2.93 GHz and 8 GB RAM
  - **timelimit** of 3600 seconds

# Setting comparison on formulation (nPC2)




(a) runtime




(b) optimality gap



# Setting comparison on formulation (nPC3)




(c) runtime




(d) optimality gap

# Formulation comparison on setting PH




(e) runtime




(f) optimality gap

# Managerial insights




(g) Relative regret of the optimal solution value

# Managerial insights



(i) Relative regret of the optimal solution value



(j) Relative regret of # of opened facilities

# Conclusion

- (nPC3) best performance in runtimes and optimality gap
- The (nPC2) outperforms the (nPC1)
- Preprocessing brings a large speed up on all formulations
- Starting heuristic little effect, shows good upper bound obtained in preprocessing
- Maximal relative regret of the optimal objective function value of 15%
- Maximal relative regret of number of opened facilities above 150%

# On the nested $p$ -center problem



Christof Brandstetter, Markus Sinnl

Institute of Business Analytics and Technology Transformation / JKU Business School,  
Johannes Kepler University Linz

This research was funded in whole, or in part, by the Austrian Science Fund (FWF) [P 35160-N].  
2024-03-06



JOHANNES KEPLER  
UNIVERSITY LINZ  
Altenberger Straße 69  
4040 Linz, Austria  
jku.at

# References I


- [1] M. Albareda-Sambola et al. The multi-period incremental service facility location problem. In: Computers & Operations Research 36.5 (2009), pp. 1356–1375.
- [2] Z. Ales and S. Elloumi. Compact MILP formulations for the  $p$ -center problem. In: Combinatorial Optimization: 5th International Symposium, ISCO 2018, Marrakesh, Morocco, April 11–13, 2018, Revised Selected Papers 5. 2018, pp. 14–25.
- [3] H. Bakker and S. Nickel. The Value of the Multi-period Solution revisited: When to model time in capacitated location problems. In: Computers & Operations Research 161 (2024), p. 106428.
- [4] M. Conforti et al. Integer programming models. Springer, 2014.

## References II


- [5] M. S. Daskin. Network and Discrete Location: Models, Algorithms, and Applications, Second Edition. John Wiley & Sons, Ltd, 2013.
- [6] E. Gaar and M. Sinnl. A scaleable projection-based branch-and-cut algorithm for the  $p$ -center problem. In: European Journal of Operational Research 303.1 (2022), pp. 78–98.
- [7] R. G. McGarvey and A. Thorsen. Nested-solution facility location models. In: Optimization letters 16.2 (2022), pp. 497–514.
- [8] G. M. Roodman and L. B. Schwarz. Optimal and heuristic facility phase-out strategies. In: AIIE transactions 7.2 (1975), pp. 177–184.



# Setting comparison on formulation (nPC1)



(k) runtime



(l) optimality gap  $[\frac{UB-LB}{UB}]$

# The nested $p$ -center problem: Classical MILP formulation

## Decision variables

$$x_{ij}^h \dots \begin{cases} 1 \dots \text{if location } i \text{ is assigned to} \\ \quad \text{location } j \text{ in time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

# The nested $p$ -center problem: Classical MILP formulation

## Decision variables

$$x_{ij}^h \dots \begin{cases} 1 \dots \text{if location } i \text{ is assigned to} \\ \quad \text{location } j \text{ in time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \quad \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

# The nested $p$ -center problem: Classical MILP formulation

## Decision variables

$$x_{ij}^h \dots \begin{cases} 1 \dots \text{if location } i \text{ is assigned to} \\ \quad \text{location } j \text{ in time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$$y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \quad \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$$

$z^h \dots$  maximum distance between any customer  $i$  and its nearest open facility in period  $h$

# The nested $p$ -center problem: Classical MILP formulation

## Decision variables

$x_{ij}^h \dots \begin{cases} 1 \dots \text{if location } i \text{ is assigned to} \\ \text{location } j \text{ in time period } h \\ 0 \dots \text{otherwise} \end{cases}$   
 $y_j^h \dots \begin{cases} 1 \dots \text{if location } j \text{ is open in} \\ \text{time period } h \\ 0 \dots \text{otherwise} \end{cases}$   
 $z^h \dots \text{maximum distance between any} \\ \text{customer } i \text{ and its nearest open facility} \\ \text{in period } h$

## (nPC1) based on Daskin 2013

$$\min \sum_{h \in \mathcal{H}} z^h \quad (4a)$$

$$\text{s.t. } \sum_{j \in V} y_j^h = p^h \quad \forall h \in \mathcal{H} \quad (4b)$$

$$\sum_{j \in V} x_{ij}^h = 1 \quad \forall i \in V, h \in \mathcal{H} \quad (4c)$$

$$x_{ij}^h \leq y_j^h \quad \forall i, j \in V, h \in \mathcal{H} \quad (4d)$$

$$\sum_{j \in V} d_{ij} x_{ij}^h \leq z^h \quad \forall i \in V, h \in \mathcal{H} \quad (4e)$$

$$y_j^h \geq y_j^{h-1} \quad \forall h \in \mathcal{H} \setminus \{1\} \quad (4f)$$

$$(x, y, z) \in |V|^2 |\mathcal{H}| \times |V| |\mathcal{H}| \times \mathbb{R}_{\geq 0} \quad (4g)$$