



UMD - Rulebased prototypes (k=3)


Class 0 - P1




Class 0 - P2




Class 0 - P3




Class 1 - P1




Class 1 - P2




Class 1 - P3




Class 2 - P1



Class 2 - P2




Class 2 - P3




UMD - rulebased support examples #1 (k=3, num_rules=0, acc=0.94)


Class 0 - idx 17




Class 0 - idx 3




Class 0 - idx 2




Class 1 - idx 1




Class 1 - idx 18




Class 1 - idx 13




Class 2 - idx 6



Class 2 - idx 12



Class 2 - idx 5



UMD - rulebased rules #1 (k=3, num_rules=0, acc=0.94)

Class 0:

R1: Contains at least one pronounced narrow upward peak that rises clearly above the surrounding plateau or baseline.

R2: Apart from this peak, the signal is mostly flat in early and late regions, with a contrasting broad high or low plateau occupying much of the middle.

R3: No short, isolated deep downward troughs are present.

Class 1:

R1: Early and late regions are flat and stay close to a common baseline level.

R2: The middle region forms a single wide high plateau that is clearly higher than this baseline and spans most of the middle of the series.

R3: Aside from the two plateau edges, the signal has no strong local spikes or dips.


Class 2:

R1: Contains at least one short, sharp deep trough where values briefly fall far below the surrounding baseline or plateau and then quickly return.


R2: Outside these troughs, the signal is piecewise flat, with early and late regions near the same baseline and at most one broad plateau or valley in the middle.

UMD - rulebased support examples #2 (k=3, num_rules=0, acc=0.91)


Class 0 - idx 17




Class 0 - idx 3




Class 0 - idx 2




Class 1 - idx 1




Class 1 - idx 18




Class 1 - idx 13




Class 2 - idx 6



Class 2 - idx 12



Class 2 - idx 5



UMD - rulebased rules #2 (k=3, num_rules=0, acc=0.91)

Class 0:

R1: The main feature is a long flat block in the middle at a level very different from the early and late segments, either as a deep central valley or as a high plateau whose rise already begins in the early region.

R2: The series shows no very narrow, V-shaped deep dips; any extreme excursion is a broad valley or a broad hump.

Class 1:

R1: A single broad, almost rectangular high plateau is confined to the middle segment and sits clearly above both the early and late segments.

R2: Early and late segments stay near the same medium level with only small noise and no deep valleys, narrow spikes, or V-shaped dips.


Class 2:

R1: At least one very narrow, V-shaped dip occurs in the early or late segment, dropping far below the neighbouring values.


R2: Apart from these dips, the middle segment forms a broader block (either a high plateau or a low valley) whose level is clearly different from the early and late baselines.

UMD - rulebased support examples #3 (k=3, num_rules=0, acc=0.92)


Class 0 - idx 17




Class 0 - idx 3




Class 0 - idx 2




Class 1 - idx 1




Class 1 - idx 18




Class 1 - idx 13




Class 2 - idx 6



Class 2 - idx 12



Class 2 - idx 5



UMD - rulebased rules #3 (k=3, num_rules=0, acc=0.92)

Class 0:

R1: Early or late region contains a pronounced narrow upward peak that is much taller than the surrounding baseline.

R2: Between the baseline segments, the middle region forms a relatively flat block so the strong upward peak appears as a short isolated event.

Class 1:

R1: A single broad high plateau occupies the middle region, with values clearly higher than both early and late regions.

R2: Early and late regions stay near a common lower baseline and do not show any sharp peaks or deep troughs.

Class 2:

R1: Early or late region contains at least one pronounced narrow downward spike that drops well below the surrounding baseline.

R2: Apart from these spikes, the series is comparatively flat, typically with a broad plateau in the middle, so the downward spike appears as an isolated deep trough.