



The Hitchhiker's Guide to Dynamic Ion-Solvent Clustering

(...in Differential Mobility Spectrometry)


Christian Ieritano, W. Scott Hopkins

CSC 2024 (Winnipeg)



What is differential mobility spectrometry (DMS)?


Differential mobility spectrometry (DMS) harnesses the non-linear dependence of an ion's mobility to separate analytes.



At high electric field strengths, an ion's mobility changes non-linearly with the applied field

$$\alpha(E) = \frac{K(E) - K(0)}{K(0)} \quad K = \frac{3}{16} \sqrt{\frac{2\pi}{\mu k_B T}} \cdot \frac{q(1 + \alpha)}{N \cdot \Omega(T)}$$


$$T_{eff} = T_{bath} + \frac{M}{3k_b} (KE)^2 (1 + \beta)$$




Analyst, 2019, **144**, 1660 - 1670.

Analyst, 2023, **148**, 3257-3273.


Example: predicting the DMS separation of prototropic isomers in a pure N₂ environment via MobCal-MPI



Simulated α -function



Measured DMS ionogram



For a specific SV, every analyte will elute from the DMS cell at a characteristic CV related to its alpha function


*from geometries and partial charges at ω B97X-D3/Def2-TZVPP



UNIVERSITY OF
WATERLOO | FACULTY
OF SCIENCE


How does DMS relate to dynamic ion solvation?

Differential mobility spectrometry (DMS) harnesses the non-linear dependence of an ion's mobility to separate analytes.




Doping the carrier gas with solvent (1.5 mol%) induces dynamic solvation/desolvation cycles

Changes to an ion's mobility upon microsolvation can be modeled



$$\Delta G_{ass} = (n \cdot G_{H_2O}) + G_{[M+H]^+} - G_{[M+H+n(H_2O)]}$$

$$P_i = P \cdot e^{\frac{\Delta G_{ass_i}(T)}{k_B T}} \quad P = \frac{1}{\sum_i P_i}$$





The nature of the SV waveform ensures **dynamic** ion-solvent interactions during DMS transit

What's so interesting about dynamic ion solvation in DMS?

1. Modifiers enhance separations of complex mixtures


Simultaneous DMS-MS measurements of 60 compounds

SV = 167 Td



2. Understanding dynamic ion-solvent interactions enables predictions of DMS behaviour


From first-principles modelling



$\Delta CV_{avg} = -0.4 \text{ V at SV} = 167 \text{ Td}$

J. Am. Soc. Mass Spectrom.
2022, **33**, 535 – 547.
Phys. Chem. Chem. Phys. 2022,
24, 20594 – 20615.
Analyst. 2023, **148**, 3257 – 3273.
J. Am. Soc. Mass Spectrom.
2023, **34**, 1417 – 1427.

Via machine learning (Random Forest Regression)




Analyst, 2021, **146**, 4737 – 4743.
Anal. Chem., 2021, **93**, 8937 – 8944.
Phys. Chem. Chem. Phys. 2022, **24**,
20594 – 20615.

$|\Delta CV_{avg}| = 2.2 \text{ V at SV} = 167 \text{ Td}$
for 300+ compounds!

What's so interesting about dynamic ion solvation in DMS?



3. Correlating DMS behaviour with solution-phase properties intrinsic to the drug-design process

Hypothesis: Dynamic microsolvation processes during an ion's transit in the DMS cell mimics its solution-phase primary solvation shell.



Linear trends observed within identical compound classes for several properties but not amongst different compound classes

Machine learning enables general predictions (5-fold cross validation)



Random forest models trained using only DMS-MS data (SV/CV pairs) acquired in a MeOH modified environment using only nanograms of material.

Phys. Chem. Chem. Phys. 2022, **24**, 20594 – 20615.
Anal. Chem. 2023, **95**, 10309 – 10321.


“Ion microsolvation” is more than a buzz word

4. Charge manipulation



<https://phys.org/news/2015-09-brazilian-wasp-venom-cancer-cells.html>

Polybia-MP1




Lys Residues & N-terminus (cationic)

Addition of solvent vapour to the DMS cell alters the charge state distribution of MP1

Dry N₂

[MP1 + 3H]³⁺




N₂ with 1.5% MeOH

[MP1 + 2H]²⁺

Shift must be mediated by microsolvation


[MP1 + 3H]³⁺




The shift in charge state distribution is likely mediated by microsolvation

Solvation/desolvation processes can induce proton transfer



[PrNH₂ + H + (MeOH)₈]⁺




[(MeOH)₈ + H]⁺



[(MeOH)₈]



Microsolvation facilitates ion-solvent proton transfer




Gas-phase basicity ordering: **ACE** > IPA > **MeCN** > EtOH > MeOH > H₂O

No charge transfer with aprotic modifiers because of inability to form H-bonded networks with high GPB!

Instead, retention of high charge states that otherwise fragment during DMS-transit


The +3 ion re-emerges at high bath gas temperatures and high SV fields



Speaking of proton transfers ...



5. Harnessing the separation power of dynamic microsolvation and HDX

Humulone: A brewer's best friend








Tautomer resolution requires detection as silver adduct (*i.e.*, $[M + Ag]^+$ **and** 1.5 mol% of IPA doped into the curtain gas.

HDX reagent (e.g., D_2O) is bubbled into the **DR gas line**, but the excess IPA would destroy any HDX reagent by back-exchange



Retro-HDX: Deuterate pre-ESI, then watch back-exchange with protic modifier




HDX events happen only at points of solvent accretion (*i.e.*, the charge site!)

Speaking of proton transfers ...



5. Harnessing the separation power of dynamic microsolvation and HDX

Humulone: A brewer's best friend




Tautomer resolution requires detection as silver adduct (*i.e.*, $[M + Ag]^+$ **and** 1.5 mol% of IPA doped into the curtain gas.


HDX reagent (e.g., D_2O) is bubbled into the **DR gas line**, but the excess IPA would destroy any HDX reagent by back-exchange




UVPD: DMS-select each tautomer, then interrogate w/ tunable output of the YAG-pumped OPO



dienol clearly distinguished




cis-keto



...but the keto tautomers are somewhat ambiguous


trans-keto



WaterFEL – A Canadian Free Electron Laser (FEL) Facility


What is an FEL?

Linear accelerator-based light sources provide **tunable, high-power, narrow bandwidth IR light from 20 – 4000 cm⁻¹**



2023 CFI award for \$10M to construct a Canadian FEL at the University of Waterloo (UW)

With matching funding from Ontario and additional contributions from UW and other institutions




$\bar{\nu} / \text{cm}^{-1}$	Energy per pulse (mJ; 10 Hz)	Bandwidth ($\Delta\bar{\nu} / \text{cm}^{-1}$)
3900*	20 – 30*	$\pm 8.3^*$
3000	50	± 6.75
2000	60	± 5.625
1000	75	± 2.25
500	50	± 1.125
100	>100	± 0.225

*not shown; hot off the presses at FHI!

WaterFEL – A Canadian Free Electron Laser (FEL) Facility

What is an FEL?

Linear accelerator-based light sources provide **tunable, high-power, narrow bandwidth IR light from 20 – 4000 cm^{-1}**




FRITZ-HABER-INSTITUT
MAX-PLANCK-GESELLSCHAFT



2023 CFI award for \$10M to construct a Canadian FEL at the University of Waterloo (UW)

With matching funding from Ontario and additional contributions from UW and other institutions

WaterFEL instrumentation




Facility open to the global community, with beam-time being awarded based on merit of proposals

First user proposals for early 2029

WaterFEL – Some of the many applications


Site-specific information about biological modifications





...combined with bottom-up approaches enable sequencing w/ unambiguous PTM assignment

Unambiguous distinction of “indistinguishable” isobars

3,3,3,2-tetrafluoropropionic acid



3,3,2,2-tetrafluoropropionic acid



Arthur Lee

Combine IRMPD and AI for molecular fingerprinting

Tuesday 4:00 – 4:20pm
(IN, MT, PTC) Modern Spectroscopic Approaches
for Deciphering Complex Phenomena

Acknowledgements

The Brain Trust



Prof. W. Scott
Hopkins



Dr. Alexander
Haack

Hopkins Group

Dr. Neville Coughlan
Dr. Nour Mashmoushi
Dr. Joshua Featherstone
Dr. Jeff Crouse
Dr. Dan Rickert
Dr. Steve Walker
Justine Bissonnette
Chris Ryan
Cailum Steinstra
Patrick Thomas
Arthur Lee


SCIEX Gurus

Dr. J. C. Yves Le Blanc
Dr. Brad Schneider
Dr. Mircea Guna

Academic friends

Dr. John Janiszewski (NIH)
Dr. Gilles Goetz (Pfizer)
Dr. Larry Campbell
(Bedrock)

For more information on our work, see:



The Hitchhiker's Guide to Dynamic Ion-Solvent Clustering

Phys. Chem. Chem. Phys., 2022, 24,
20594-20615.

Resources and Funding



Digital Research
Alliance of Canada

Alliance de recherche
numérique du Canada

