Functions

J. Alexander Branham

Fall 2015

• Exponents tell you to multiply that thing by its base x times:

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x:

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get
 x:
 - $log_3(81) = 4$

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x:
 - $log_3(81) = 4$
 - Note that logarithms with negative arguments are undefined

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get
 x:
 - $log_3(81) = 4$
 - Note that logarithms with negative arguments are undefined
 - Sometimes log(n) means $log_{10}(n)$

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 * 3 * 3 * 3 =$
- Logarithms ask how many times you must raise the base to get x:
 - $log_3(81) = 4$
 - Note that logarithms with negative arguments are undefined
 - Sometimes log(n) means $log_{10}(n)$
 - Othertimes, it means $log_e(n) = ln(n)$

$$a^m a^n = a^{m+n}$$

$$a^m a^n = a^{m+n}$$

$$a^{\frac{a}{n}} = a^{m-n}$$

- $a^m a^n = a^{m+n}$
- $\bullet \ \frac{a^{m}}{a^{n}} = a^{m-n}$
- $\bullet \ (a^m)^n = a^{mn}$

$$a^m a^n = a^{m+n}$$

$$a^m a^n = a^{m+n}$$

$$\frac{a^m}{a^n} = a^{m-n}$$

$$\bullet \ (a^m)^n = a^{mn}$$

•
$$a^0 = 1$$

$$\bullet \ a^m a^n = a^{m+n}$$

$$a^m a^n = a^{m+n}$$

$$\frac{a^m}{a^n} = a^{m-n}$$

$$\bullet \ (a^m)^n = a^{mn}$$

•
$$a^0 = 1$$

•
$$a^{1/n} = \sqrt[n]{a}$$

•
$$a^m a^n = a^{m+n}$$

•
$$a^m a^n = a^{m+n}$$

• $\frac{a^m}{a^n} = a^{m-n}$

•
$$(a^m)^n = a^{mn}$$

•
$$a^0 = 1$$

•
$$a^{1/n} = \sqrt[n]{a}$$

$$\bullet \left(\frac{a}{b}\right)^n = \frac{a^n}{b^n} = a^n b^{-n} \qquad \forall b \neq 0$$

•
$$a^m a^n = a^{m+n}$$

$$a^m a^n = a^{m+n}$$

$$\frac{a^m}{a^n} = a^{m-n}$$

$$\bullet (a^m)^n = a^{mn}$$

•
$$a^0 = 1$$

•
$$a^{1/n} = \sqrt[n]{a}$$

•
$$\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n} = a^n b^{-n} \quad \forall b \neq 0$$

• $\left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n \quad \forall a, b \neq 0$

•
$$log_x(ab) = log_x a + log_x b$$

- $log_x(ab) = log_x a + log_x b$
- $\log_{\mathsf{X}}(\frac{\mathsf{a}}{\mathsf{b}}) = \log_{\mathsf{X}}\mathsf{a} \log_{\mathsf{X}}\mathsf{b}$

- $log_x(ab) = log_x a + log_x b$
- $log_x(\frac{a}{b}) = log_x a log_x b$
- $log_X a^b = blog_X a$

- $log_x(ab) = log_x a + log_x b$
- $log_x(\frac{a}{b}) = log_x a log_x b$
- $log_x a^b = blog_x a$
- $log_{x}1 = 0$

- $log_x(ab) = log_x a + log_x b$
- $log_x(\frac{a}{b}) = log_x a log_x b$
- $log_x a^b = blog_x a$
- $log_{x}1 = 0$
- $m^{\log_m(a)} = a$

•
$$log_x(ab) = log_x a + log_x b$$

•
$$log_x(\frac{a}{b}) = log_x a - log_x b$$

•
$$log_x a^b = blog_x a$$

•
$$log_{x}1 = 0$$


•
$$m^{log_m(a)} = a$$

• Anything that takes input and gives one output

- Anything that takes input and gives one output
- In math, this usually looks something like f(x, z) = y

- Anything that takes input and gives one output
- In math, this usually looks something like f(x, z) = y
 - x and z are the arguments that the function takes

- Anything that takes input and gives one output
- In math, this usually looks something like f(x, z) = y
 - x and z are the arguments that the function takes
 - *y* is the *output* from the function



• We can make a function that describes a line pretty easily

- We can make a function that describes a line pretty easily
- \bullet y = mx + b

- We can make a function that describes a line pretty easily
- y = mx + b
 - m is the slope (for every one unit increase in x, y increases m units)

- We can make a function that describes a line pretty easily
- y = mx + b
 - m is the slope (for every one unit increase in x, y increases m units)
 - b is the x-intercept: the value of y when x = 0

Functions

```
ggplot(data.frame(x=c(-3, 3)), aes(x)) +
  stat_function(fun=function(x)-2*x+3, geom="line") +
  stat_function(fun=function(x)(1/2)*x+1)
```




Figure 1:

Quadratics


• These lines have one curve

Quadratics

- These lines have one curve
- $y = ax^2 + bx + c$

Quadratics


```
ggplot(data.frame(x=c(-3,4)), aes(x)) +
  stat_function(fun=function(x)x^2, color="green") +
  stat_function(fun=function(x)2*x^2, color="red") +
  stat_function(fun=function(x)-2*x^2 + 6*x -4) +
  ylim(c(-5, 5))
```



Exponential


• General form: $y = a * b^{kx} + k$

```
ggplot(data.frame(x=c(-1, 5)), aes(x)) +
   stat_function(fun=function(x)exp(x)) +
   stat_function(fun=function(x)2*exp(x), color="red")
```



• General form: y = a * log(bx) + k

```
ggplot(data.frame(x=c(0,5)), aes(x)) +
  stat_function(fun=function(x)log(x)) +
  stat_function(fun=function(x)log10(x), color="red") +
  stat_function(fun=function(x)2*log(x), color="green")
```



• The log and exponent charts are obviously related

- The log and exponent charts are obviously related
- In fact, these functions are each others "inverse" function

- The log and exponent charts are obviously related
- In fact, these functions are each others "inverse" function
 - Plug in y to find x

- The log and exponent charts are obviously related
- In fact, these functions are each others "inverse" function
 - Plug in y to find x
- Exponents have horizontal asymptote

- The log and exponent charts are obviously related
- In fact, these functions are each others "inverse" function
 - Plug in y to find x
- Exponents have horizontal asymptote
- Logs have vertical asymptote