

# Introduction to Probability

J. Alexander Branham

Fall 2015

# Probability

- Frequency with which an event occurs

# Probability

- Frequency with which an event occurs
- Typically:

$$Pr(A) = P(A) = \pi(A) = \frac{\text{Number of ways an event can occur}}{\text{Total number of possible outcomes}}$$

# Probability

- Frequency with which an event occurs
- Typically:

$$Pr(A) = P(A) = \pi(A) = \frac{\text{Number of ways an event can occur}}{\text{Total number of possible outcomes}}$$

- There are other ways of thinking about probability, but we'll stick with this one

# Kolmogorov's Axioms

- First:  $Pr(E) \in \mathbb{R}, Pr(E) \geq 0 \quad \forall E \in F$

# Kolmogorov's Axioms

- First:  $Pr(E) \in \mathbb{R}, Pr(E) \geq 0 \quad \forall E \in F$ 
  - Probabilities must be non-negative

# Kolmogorov's Axioms

- First:  $Pr(E) \in \mathbb{R}, Pr(E) \geq 0 \quad \forall E \in \mathcal{F}$ 
  - Probabilities must be non-negative
- Second:  $Pr(\Omega) = 1$

# Kolmogorov's Axioms

- First:  $Pr(E) \in \mathbb{R}, Pr(E) \geq 0 \quad \forall E \in \mathcal{F}$ 
  - Probabilities must be non-negative
- Second:  $Pr(\Omega) = 1$ 
  - Something has to happen



# Kolmogorov's Axioms

- First:  $Pr(E) \in \mathbb{R}, Pr(E) \geq 0 \quad \forall E \in \mathcal{F}$ 
  - Probabilities must be non-negative
- Second:  $Pr(\Omega) = 1$ 
  - Something has to happen
  - Probabilities sum/integrate to 1

# Kolmogorov's Axioms

- First:  $Pr(E) \in \mathbb{R}, Pr(E) \geq 0 \quad \forall E \in \mathcal{F}$ 
  - Probabilities must be non-negative
- Second:  $Pr(\Omega) = 1$ 
  - Something has to happen
  - Probabilities sum/integrate to 1
- Third:  $Pr(\bigcup_{i=1}^{\infty} E_i) = \sum_{i=1}^{\infty} Pr(E_i)$

# Kolmogorov's Axioms

- First:  $Pr(E) \in \mathbb{R}, Pr(E) \geq 0 \quad \forall E \in \mathcal{F}$ 
  - Probabilities must be non-negative
- Second:  $Pr(\Omega) = 1$ 
  - Something has to happen
  - Probabilities sum/integrate to 1
- Third:  $Pr(\bigcup_{i=1}^{\infty} E_i) = \sum_{i=1}^{\infty} Pr(E_i)$ 
  - The probability of disjoint (mutually exclusive) sets is equal to their sums

# Discrete probabilities

- What's the probability that we'll roll a 3 on one die roll:

$$Pr(y = 3) = \frac{1}{6}$$

# Discrete probabilities

- What's the probability that we'll roll a 3 on one die roll:

$$Pr(y = 3) = \frac{1}{6}$$

- If one roll of the die is an “experiment”

# Discrete probabilities

- What's the probability that we'll roll a 3 on one die roll:

$$Pr(y = 3) = \frac{1}{6}$$

- If one roll of the die is an “experiment”
- We can think of a 3 as a “success”

# Discrete probabilities

- What's the probability that we'll roll a 3 on one die roll:

$$Pr(y = 3) = \frac{1}{6}$$

- If one roll of the die is an “experiment”
- We can think of a 3 as a “success”
- And  $Y \sim \text{Bern}(\frac{1}{6})$

# Discrete probabilities

- What's the probability that we'll roll a 3 on one die roll:

$$Pr(y = 3) = \frac{1}{6}$$

- If one roll of the die is an “experiment”
- We can think of a 3 as a “success”
- And  $Y \sim \text{Bern}(\frac{1}{6})$
- Fair coins are  $\sim \text{Bern}(.5)$  for example



## Discrete probabilities

- What's the probability that we'll roll a 3 on one die roll:

$$Pr(y = 3) = \frac{1}{6}$$

- If one roll of the die is an “experiment”
- We can think of a 3 as a “success”
- And  $Y \sim \text{Bern}(\frac{1}{6})$
- Fair coins are  $\sim \text{Bern}(.5)$  for example
- More generally  $\text{Bern}(\pi)$

## Discrete probabilities

- What's the probability that we'll roll a 3 on one die roll:

$$Pr(y = 3) = \frac{1}{6}$$

- If one roll of the die is an “experiment”
- We can think of a 3 as a “success”
- And  $Y \sim \text{Bern}(\frac{1}{6})$
- Fair coins are  $\sim \text{Bern}(.5)$  for example
- More generally  $\text{Bern}(\pi)$ 
  - $\pi$  represents the probability of success

# Binomial distributions

- Before we ran the experiment just once

# Binomial distributions

- Before we ran the experiment just once
- What happens if we run it multiple times?

$$Pr(y_1 = 3, y_2 = 3) = \frac{1}{6} * \frac{1}{6}$$

# Binomial distributions

- Before we ran the experiment just once
- What happens if we run it multiple times?

$$Pr(y_1 = 3, y_2 = 3) = \frac{1}{6} * \frac{1}{6}$$

- Now  $Y \sim \text{Binom}(2, \frac{1}{6})$

# Binomial distributions

- Before we ran the experiment just once
- What happens if we run it multiple times?

$$Pr(y_1 = 3, y_2 = 3) = \frac{1}{6} * \frac{1}{6}$$

- Now  $Y \sim \text{Binom}(2, \frac{1}{6})$ 
  - Generally,  $Y \sim \text{Binom}(n, p)$

# Binomial distributions

- Before we ran the experiment just once
- What happens if we run it multiple times?

$$Pr(y_1 = 3, y_2 = 3) = \frac{1}{6} * \frac{1}{6}$$

- Now  $Y \sim \text{Binom}(2, \frac{1}{6})$ 
  - Generally,  $Y \sim \text{Binom}(n, p)$
  - $n$  = number of trials,  $p$  = probability of success

# Binomial distributions

- Before we ran the experiment just once
- What happens if we run it multiple times?

$$Pr(y_1 = 3, y_2 = 3) = \frac{1}{6} * \frac{1}{6}$$

- Now  $Y \sim \text{Binom}(2, \frac{1}{6})$ 
  - Generally,  $Y \sim \text{Binom}(n, p)$
  - $n$  = number of trials,  $p$  = probability of success
- PMF:

$$\binom{n}{k} p^k (1 - p)^{n-k}$$



# The basics

- What happens when our outcome is continuous

# The basics

- What happens when our outcome is continuous
- Much harder to think about. . .

# The basics

- What happens when our outcome is continuous
- Much harder to think about. . .
- There are infinity possible outcomes

# The basics

- What happens when our outcome is continuous
- Much harder to think about. . .
- There are infinity possible outcomes
- Makes the denominator of our fraction difficult to work with

# The basics

- What happens when our outcome is continuous
- Much harder to think about. . .
- There are infinity possible outcomes
- Makes the denominator of our fraction difficult to work with
- What's the probability that you make \$15,293.19/year?

# The basics

- What happens when our outcome is continuous
- Much harder to think about. . .
- There are infinity possible outcomes
- Makes the denominator of our fraction difficult to work with
- What's the probability that you make \$15,293.19/year?
- Probability of the whole space must equal 1

# The basics

- What happens when our outcome is continuous
- Much harder to think about. . .
- There are infinity possible outcomes
- Makes the denominator of our fraction difficult to work with
- What's the probability that you make \$15,293.19/year?
- Probability of the whole space must equal 1
- Even if all events equally likely,  $\frac{1}{\infty} = 0$


# The basics

- What happens when our outcome is continuous
- Much harder to think about. . .
- There are infinity possible outcomes
- Makes the denominator of our fraction difficult to work with
- What's the probability that you make \$15,293.19/year?
- Probability of the whole space must equal 1
- Even if all events equally likely,  $\frac{1}{\infty} = 0$ 
  - Kinda. . .




# Continuous distributions - Normal

```
ggplot(data.frame(x = c(-5, 5)), aes(x)) +  
  stat_function(fun = dnorm)
```



## Continuous distributions - Uniform

```
ggplot(data.frame(x = c(-2, 2)), aes(x)) +  
  stat_function(fun = dunif)
```



# Probabilities and continuous distributions

- $Pr(y = c) = 0$

# Probabilities and continuous distributions

- $Pr(y = c) = 0$
- $Pr(0 < y < .5) = Pr(0 \geq y \leq .5)$

$$= \int_0^{.5} f(y) dy$$

# Probabilities and continuous distributions

- $Pr(y = c) = 0$
- $Pr(0 < y < .5) = Pr(0 \geq y \leq .5)$

$$= \int_0^{.5} f(y) dy$$

- For uniform from previous slide,  $Pr(0 < y < .5) = 0.5$

# CDF

- CDF = Cumulative Distribution Function

# CDF

- CDF = Cumulative Distribution Function
- $F_X(x) = Pr(X \leq x)$

# Discrete

- $Y \sim \text{Binom}(10, .5)$



# Discrete

- $Y \sim \text{Binom}(10, .5)$
- What's the probability that  $y < 5$  ?

# Discrete

- $Y \sim \text{Binom}(10, .5)$
- What's the probability that  $y < 5$  ?
- 

$$\sum_{y_i < y} \text{Pr}(Y = y_i)$$

$$Pr(y \leq 5)$$

$$\begin{aligned} & \binom{10}{1} .5^1 (1 - .5)^{10-1} + \\ & \binom{10}{2} .5^2 (1 - .5)^{10-2} + \\ & \binom{10}{3} .5^3 (1 - .5)^{10-3} + \\ & \binom{10}{4} .5^4 (1 - .5)^{10-4} + \\ & \binom{10}{5} .5^5 (1 - .5)^{10-5} \end{aligned}$$

# Continuous

- We can't sum probabilities for continuous distributions (remember the 0 problem)

# Continuous

- We can't sum probabilities for continuous distributions (remember the 0 problem)
- Solution: integration

# Continuous

- We can't sum probabilities for continuous distributions (remember the 0 problem)
- Solution: integration
- $F_Y(y) = \int_{-\infty}^y f(y)dy$