

Introduction to Math for Political Scientists

J. Alexander Branham & Megan Moeller

Fall 2015

Let's start real slow...

- So we all know that

$$2 + 2 = 4$$

Let's start real slow...

- So we all know that

$$2 + 2 = 4$$

- But addition and subtraction have some cool (grool?) rules

Properties of Addition and subtraction

- Communicative:

Properties of Addition and subtraction

- Communicative:
 - $a \pm b = b \pm a$

Properties of Addition and subtraction

- Communicative:
 - $a \pm b = b \pm a$
- Associative

Properties of Addition and subtraction

- Communicative:
 - $a \pm b = b \pm a$
- Associative
 - $(a \pm b) \pm c = a \pm (b \pm c)$

Multiplication

- Multiplication - I have these 4 things 10 times.

$$4+4+4+4+4+4+4+4+4+4$$

[1] 40

Multiplication

- Multiplication - I have these 4 things 10 times.

$$4+4+4+4+4+4+4+4+4+4$$

[1] 40

Multiplication

- Multiplication - I have these 4 things 10 times.

$$4+4+4+4+4+4+4+4+4+4$$

[1] 40

Or I could just do

$$4*10$$

[1] 40

Division

- Just fancy multiplication.

Division

- Just fancy multiplication.

Division

- Just fancy multiplication.
- I have these four things one of ten times.

$$4 * (1/10)$$

```
## [1] 0.4
```

Properties of Multiplication and Division

- Communicative

Properties of Multiplication and Division

- Communicative
 - $a * b = b * a$

Properties of Multiplication and Division

- Communicative
 - $a * b = b * a$
- Associative

Properties of Multiplication and Division

- Communicative
 - $a * b = b * a$
- Associative
 - $(ab)c = a(bc)$

Properties of Multiplication and Division

- Communicative
 - $a * b = b * a$
- Associative
 - $(ab)c = a(bc)$
- Distributive

Properties of Multiplication and Division

- Communicative
 - $a * b = b * a$
- Associative
 - $(ab)c = a(bc)$
- Distributive
 - $a(b + c) = ab + ac$

Properties of Multiplication and Division

- Communicative

- $a * b = b * a$

- Associative

- $(ab)c = a(bc)$

- Distributive

- $a(b + c) = ab + ac$

- Note that this works for division: $\frac{a + b}{c} = \frac{a}{c} + \frac{b}{c}$

Relationships that hold with (real) numbers

- $a = b \iff b = a$ (Symmetric relationships)

Relationships that hold with (real) numbers

- $a = b \iff b = a$ (Symmetric relationships)
- $a = b$ and $b = c \Rightarrow a = c$ (Transitive relationships)

Relationships that hold with (real) numbers

- $a = b \iff b = a$ (Symmetric relationships)
- $a = b$ and $b = c \Rightarrow a = c$ (Transitive relationships)
 - $a > b$ and $b > c \Rightarrow a > c$

PEMDAS

- Parentheses

PEMDAS

- Parentheses
- Exponents

PEMDAS

- Parentheses
- Exponents
- Multiplication and division (left to right)

PEMDAS

- Parentheses
- Exponents
- Multiplication and division (left to right)
- Addition and subtraction (left to right)

$$(10 - 48 \div 12 * 2)^2 + 3^2 * (8 - 6)$$

Exponents and Logarithms

- Exponents tell you to multiply that thing by its base x times:

Exponents and Logarithms

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 333*3 = 81$

Exponents and Logarithms

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 333*3 = 81$
- Logarithms ask how many times you must raise the base to get x :

Exponents and Logarithms

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 333*3 = 81$
- Logarithms ask how many times you must raise the base to get x :
 - $\log_3(81) = 4$

Exponents and Logarithms

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 333 \cdot 3 = 81$
- Logarithms ask how many times you must raise the base to get x :
 - $\log_3(81) = 4$
 - Note that logarithms with negative arguments are undefined

Exponents and Logarithms

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 \cdot 3 \cdot 3 \cdot 3 = 81$
- Logarithms ask how many times you must raise the base to get x :
 - $\log_3(81) = 4$
 - Note that logarithms with negative arguments are undefined
 - Sometimes $\log(n)$ means $\log_{10}(n)$

Exponents and Logarithms

- Exponents tell you to multiply that thing by its base x times:
 - $3^4 = 3 \cdot 3 \cdot 3 \cdot 3 = 81$
- Logarithms ask how many times you must raise the base to get x :
 - $\log_3(81) = 4$
 - Note that logarithms with negative arguments are undefined
 - Sometimes $\log(n)$ means $\log_{10}(n)$
 - Othertimes, it means $\log_e(n) = \ln(n)$

Properties of exponents

- $a^m a^n = a^{m+n}$

Properties of exponents

- $a^m a^n = a^{m+n}$
- $\frac{a^m}{a^n} = a^{m-n}$

Properties of exponents

- $a^m a^n = a^{m+n}$
- $\frac{a^m}{a^n} = a^{m-n}$
- $(a^m)^n = a^{mn}$

Properties of exponents

- $a^m a^n = a^{m+n}$
- $\frac{a^m}{a^n} = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $a^0 = 1$

Properties of exponents

- $a^m a^n = a^{m+n}$
- $\frac{a^m}{a^n} = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $a^0 = 1$
- $a^{1/n} = \sqrt[n]{a}$

Properties of exponents

- $a^m a^n = a^{m+n}$
- $\frac{a^m}{a^n} = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $a^0 = 1$
- $a^{1/n} = \sqrt[n]{a}$
- $\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n} = a^n b^{-n} \quad \forall b \neq 0$

Properties of exponents

- $a^m a^n = a^{m+n}$
- $\frac{a^m}{a^n} = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $a^0 = 1$
- $a^{1/n} = \sqrt[n]{a}$
- $\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n} = a^n b^{-n} \quad \forall b \neq 0$
- $\left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n \quad \forall a, b \neq 0$

Properties of logs

- $\log_x(ab) = \log_x a + \log_x b$

Properties of logs

- $\log_x(ab) = \log_x a + \log_x b$
- $\log_x\left(\frac{a}{b}\right) = \log_x a - \log_x b$

Properties of logs

- $\log_x(ab) = \log_x a + \log_x b$
- $\log_x\left(\frac{a}{b}\right) = \log_x a - \log_x b$
- $\log_x a^b = b \log_x a$

Properties of logs

- $\log_x(ab) = \log_x a + \log_x b$
- $\log_x\left(\frac{a}{b}\right) = \log_x a - \log_x b$
- $\log_x a^b = b \log_x a$
- $\log_x 1 = 0$

Properties of logs

- $\log_x(ab) = \log_x a + \log_x b$
- $\log_x\left(\frac{a}{b}\right) = \log_x a - \log_x b$
- $\log_x a^b = b \log_x a$
- $\log_x 1 = 0$
- $m^{\log_m(a)} = a$

Properties of logs

- $\log_x(ab) = \log_x a + \log_x b$
- $\log_x\left(\frac{a}{b}\right) = \log_x a - \log_x b$
- $\log_x a^b = b \log_x a$
- $\log_x 1 = 0$
- $m^{\log_m(a)} = a$
- $\frac{\log_x n}{\log_x m} = \log_m n$