

Doctoral training school

Methods in Malaria Modeling

Simulation algorithms & numerics for epidemiological models

Christian Selinger, christian.selinger@aims.ac.rw

Nov 13 - Dec 8, 2023



AIMS Center Senegal, Mbour

Outline

1 Introduction

2 Numerics

3 Stochastics

- Randomness from the computer
- Biochemical reaction systems
- Stochastic simulation algorithms

Question block

Discuss in class!



Important block

This is important!



Search block




Search and think outside the box (and the classroom)!

sites.google.com/aims.ac.rw/mamodafrica-trainingschool/week-3/ **modsimul**

Influenza outbreak in a boarding school: the data

date	in bed	convalescent	total
1978-01-22	3	0	763
1978-01-23	8	0	763
1978-01-24	26	0	763
1978-01-25	76	0	763
1978-01-26	225	9	763
1978-01-27	298	17	763
1978-01-28	258	105	763
1978-01-29	233	162	763

The R programming language logo, consisting of a blue letter 'R' inside a grey hexagon.

- **time series** of symptomatic (i.e. infectious) cases

- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now

- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population

- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population
compartments for susceptible S , infectious I , recovered R individuals

- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population
compartments for susceptible S , infectious I , recovered R individuals
- **closed** population, no deaths

- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population
compartments for susceptible S , infectious I , recovered R individuals
- **closed** population, no deaths
constant population size, no births nor migration

- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population
compartments for susceptible S , infectious I , recovered R individuals
- **closed** population, no deaths
constant population size, no births nor migration
- **well-mixed** population

- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population
compartments for susceptible S , infectious I , recovered R individuals
- **closed** population, no deaths
constant population size, no births nor migration
- **well-mixed** population
mass action principle, force $\lambda(I)$ acting on mass S


- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population
compartments for susceptible S , infectious I , recovered R individuals
- **closed** population, no deaths
constant population size, no births nor migration
- **well-mixed** population
mass action principle, force $\lambda(I)$ acting on mass S
- **denser** population in dormitory \Rightarrow more infections

- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population
compartments for susceptible S , infectious I , recovered R individuals
- **closed** population, no deaths
constant population size, no births nor migration
- **well-mixed** population
mass action principle, force $\lambda(I)$ acting on mass S
- **denser** population in dormitory \Rightarrow more infections
density-dependent (vs. frequency-dependent) force


- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population
compartments for susceptible S , infectious I , recovered R individuals
- **closed** population, no deaths
constant population size, no births nor migration
- **well-mixed** population
mass action principle, force $\lambda(I)$ acting on mass S
- **denser** population in dormitory \Rightarrow more infections
density-dependent (vs. frequency-dependent) force
- **full immunity** upon recovery

- **time series** of symptomatic (i.e. infectious) cases
ordinary differential equations/ Markov process, what happens next depends only on now
- **homogeneous** population
compartments for susceptible S , infectious I , recovered R individuals
- **closed** population, no deaths
constant population size, no births nor migration
- **well-mixed** population
mass action principle, force $\lambda(I)$ acting on mass S
- **denser** population in dormitory \Rightarrow more infections
density-dependent (vs. frequency-dependent) force
- **full immunity** upon recovery
recovered individuals cannot become susceptible again


Influenza outbreak in a boarding school: bathtubs



Influenza outbreak in a boarding school: bathtubs



Influenza outbreak in a boarding school: bathtubs



Influenza outbreak in a boarding school: compartments



Influenza outbreak in a boarding school: compartments



- **rate** = number of events happening within time step Δ

Influenza outbreak in a boarding school: compartments



- **rate** = number of events happening within time step Δ
- γ **recovery rate** from infection



- **rate** = number of events happening within time step Δ
- γ **recovery rate** from infection
- $\lambda \equiv \lambda(I)$ **force of infection rate**

Influenza outbreak in a boarding school: compartments

Recurrence equation with update linear in time increment Δ

$$\begin{aligned} S(t + \Delta) &= S(t) + \Delta \{-\lambda(I(t))S(t)\} \\ I(t + \Delta) &= I(t) + \Delta \{\lambda(I(t))S(t) - \gamma I\} \\ R(t + \Delta) &= R(t) + \Delta \{\gamma I(t)\} \end{aligned}$$

Initial condition

$$\begin{aligned} S(0) &= S_0 < N \\ I(0) &= I_0 > 0 \\ R(0) &= 0 \end{aligned}$$

Force of infection



$\lambda(I)$ rate at which new infectious created from susceptible

Force of infection



$\lambda(I)$ rate at which new infectious created from susceptible

Density-dependent transmission



Per capita contact rate between susceptible and infected depends on the **population density**. Transmission rates increase with density.

Force of infection



$\lambda(I)$ rate at which new infectious created from susceptible

Density-dependent transmission







Per capita contact rate between susceptible and infected depends on the **population density**. Transmission rates increase with density.

Frequency-dependent transmission



Per capita contact rate between susceptible and infected **does not depend** on the population density. Transmission rates do not change with density.



Density- vs frequency-dependent transmission



Density- vs frequency-dependent transmission



Density-dependent transmission

more individuals per area increases transmission



Frequency-dependent transmission



more individuals, no impact transmission



Density- vs frequency-dependent transmission

Density-dependent transmission



more individuals per area increases transmission



Influenza, Coronavirus, Malaria?, Polio

Frequency-dependent transmission

more individuals, no impact transmission



HIV, Malaria?

Force of infection formula



$\lambda(I) = c \frac{I}{N} v$ with **contact rate**, probability of contact with infected individual, **probability that contact $S \longleftrightarrow I$ leads to transmission**

Two choices for **contact rate**:

1 **$c = k \frac{N}{A}$** : slope k of **density-dependent** contact rate per area A :

$$\lambda(I) = k \frac{N}{A} v \frac{I}{N} = \underbrace{\frac{k}{A} v}_{\beta} I = \beta I$$

2 **$c = k'$ constant, frequency-dependent contact rate:**

$$\lambda(I) = \underbrace{k' v}_{\beta'} \frac{I}{N} = \beta' \frac{I}{N}$$

Force of infection formula



$\lambda(I) = c \frac{I}{N} v$ with contact rate, probability of contact with infected individual, probability that contact $S \longleftrightarrow I$ leads to transmission

1 **density-dependent** $\lambda(I) = \beta I$


2 **frequency-dependent** $\lambda(I) = \beta' \frac{I}{N}$

If N constant: mathematically equivalent but $\beta, \frac{\beta'}{N}$ different **biological meaning**

Begon et al. 


Non-linear force of infection (foi)

- **linear** $\lambda(I) \sim I$: mass action




Non-linear force of infection (foi)

- **linear** $\lambda(I) \sim I$: mass action
- **quadratic** $\lambda(I) \sim I^2$: panic behavior




Non-linear force of infection (foi)

- **linear** $\lambda(I) \sim I$: mass action
- **quadratic** $\lambda(I) \sim I^2$: panic behavior
- **Michaelis-Menten** $\lambda(I) \sim \frac{aI}{b+I}$: a maximum rate, b level of I by which half of λ reached, saturation




Non-linear force of infection (foi)

- **linear** $\lambda(I) \sim I$: mass action
- **quadratic** $\lambda(I) \sim I^2$: panic behavior
- **Michaelis-Menten** $\lambda(I) \sim \frac{aI}{b+I}$: a maximum rate, b level of I by which half of λ reached, saturation
- **crowding** $\lambda(I) \sim \frac{aI^2}{b+I^2}$: saturation



Non-linear force of infection (foi)

- **linear** $\lambda(I) \sim I$: mass action
- **quadratic** $\lambda(I) \sim I^2$: panic behavior
- **Michaelis-Menten** $\lambda(I) \sim \frac{aI}{b+I}$: a maximum rate, b level of I by which half of λ reached, saturation
- **crowding** $\lambda(I) \sim \frac{aI^2}{b+I^2}$: saturation
- **intervention** $\lambda \sim \frac{I}{f(I)}$, $f > 0$, $f' \geq 0$



- Recurrence equation with time increment Δ and $t_0 = 0$:

$$S(t + \Delta) = S(t) + \Delta \{-\lambda(I(t))S(t)\} \quad (1)$$

$$I(t + \Delta) = I(t) + \Delta \{\lambda(I(t))S(t) - \gamma I(t)\} \quad (2)$$

$$R(t + \Delta) = R(t) + \Delta \gamma I(t) \quad (3)$$

(4)

- Recurrence equation with time increment Δ and $t_0 = 0$:

$$S(t + \Delta) = S(t) + \Delta \{-\lambda(I(t))S(t)\} \quad (1)$$

$$I(t + \Delta) = I(t) + \Delta \{\lambda(I(t))S(t) - \gamma I(t)\} \quad (2)$$

$$R(t + \Delta) = R(t) + \Delta \gamma I(t) \quad (3)$$

(4)

Influenza outbreak in a boarding school: differential equation

- Recurrence equation with time increment Δ and $t_0 = 0$:

$$S(t + \Delta) = S(t) + \Delta \{-\lambda(I(t))S(t)\} \quad (1)$$

$$I(t + \Delta) = I(t) + \Delta \{\lambda(I(t))S(t) - \gamma I(t)\} \quad (2)$$

$$R(t + \Delta) = R(t) + \Delta \gamma I(t) \quad (3)$$

(4)

- First order differential equation, for all $t \geq 0$:

$$\lim_{\Delta \rightarrow 0} \frac{S(t + \Delta) - S(t)}{\Delta} = \frac{dS}{dt} = -\lambda(I)S$$

$$\lim_{\Delta \rightarrow 0} \frac{I(t + \Delta) - I(t)}{\Delta} = \frac{dI}{dt} = \lambda(I)S - \gamma I$$

$$\lim_{\Delta \rightarrow 0} \frac{R(t + \Delta) - R(t)}{\Delta} = \frac{dR}{dt} = \gamma I$$

For equidistant time points $0 = t_0 < t_1 < \dots < t_n$ write $\Delta \equiv \Delta t = t_{i+1} - t_i$, and $t_k = k\Delta t$:

$$S(t_{i+1}) = S(t_i) + \Delta \left\{ -\frac{\beta}{N} I(t_i) S(t_i) \right\} \quad (5)$$

$$I(t_{i+1}) = I(t_i) + \Delta \left\{ \frac{\beta}{N} I(t_i) S(t_i) - \gamma I \right\} \quad (6)$$

$$R(t_{i+1}) = R(t_i) + \Delta \gamma I(t_i) \quad (7)$$

For equidistant time points $0 = t_0 < t_1 < \dots < t_n$ write $\Delta \equiv \Delta t = t_{i+1} - t_i$, and $t_k = k\Delta t$:

$$S(t_{i+1}) = S(t_i) + \Delta \left\{ -\frac{\beta}{N} I(t_i) S(t_i) \right\} \quad (5)$$

$$I(t_{i+1}) = I(t_i) + \Delta \left\{ \frac{\beta}{N} I(t_i) S(t_i) - \gamma I \right\} \quad (6)$$

$$R(t_{i+1}) = R(t_i) + \Delta \gamma I(t_i) \quad (7)$$

Group work: Solve the influenza SIR model numerically in R!



- Create a sequence of time steps t_i up to 15 days with step size $\Delta = 0.5$
- Create data frame, first row is initial condition $S = 762, I = 1, R = 0$
- $\beta = 1.1, \gamma = 0.5, N = 763$, try different β such that $\frac{\beta}{\gamma} < 1$ or $\frac{\beta}{\gamma} > 1$
- Write a loop over i and plot the graph $t_i \mapsto I(t_i)$  01_ForwardEulerSIR.R

Numerical scheme for ordinary differential equation



Given an ODE $\frac{dx}{dt} = f(t, x)$ an (explicit one-step) scheme is given by continuous function $\Phi(t, x, h)$ with mesh $0 = t_0 < t_1 < \dots t_n = T$ and $\Delta t = t_{i+1} - t_i$ s.th.

$$x^{k+1} = x^k + \Delta t \Phi(t_k, x^k, \Delta t)$$

Truncation error



The truncation error is $T_k(\Delta t) = \frac{x^{k+1} - x^k}{\Delta t} - \Phi(t_k, x(t_k), \Delta t)$

$$\lim_{\Delta t \rightarrow 0} T_k(\Delta t) = \frac{dx}{dt} - \Phi(t_k, x, 0)$$

Consistency

The scheme is **consistent** with the ODE if $\Phi(t, x, 0) = f(t, x)$



Stability

The scheme is **stable** if $x \mapsto \Phi(t, x, h)$ is globally Lipschitz (i.e. almost differentiable)



Convergence

The scheme is converging if the global error $|x^k - x(t_k)| \rightarrow 0$ as $\Delta t \rightarrow 0$



Dahlquist-Lax Theorem

Convergence \Leftrightarrow Consistency + Stability



Explicit Euler is convergent



Set $\Phi(t_k, x^k, h) = f(t_k, x^k)$, for $h \in [0, H]$, $t \in [0, T]$. Discuss why this scheme is convergent!

Remember from highschool: **Taylor** expansion

Any smooth function φ can be written locally around a point a :

$$\varphi(x) = \varphi(a) + \frac{(x-a)}{1!} \frac{d}{dx} \varphi(a) + \frac{(x-a)^2}{2!} \frac{d^2}{dx^2} \varphi(a) + \dots$$

- Apply Taylor to solution curve $t \mapsto x(t)$ at discretization points t_k :

$$x(t_{k+1}) = x(t_k + \Delta t) = x(t_k) + \frac{\Delta t}{1!} \frac{d}{dt} x(t_k) + \frac{(\Delta t)^2}{2!} \frac{d^2}{dt^2} x(t_k) + \dots$$

- since $\frac{d}{dt} x(t_k) = f(t, x(t_k))$, and $\frac{d^2}{dt^2} x(t_k) = \frac{\partial f}{\partial t}(t, x_k) + \frac{\partial f}{\partial x}(t, x_k) \frac{d}{dt} x(t, x_k)$
- numeric scheme

$$x(t_{k+1}) = x(t_k) + (\Delta t) f(t, x(t_k)) + \frac{1}{2} (\Delta t)^2 \left\{ \frac{\partial f}{\partial t}(t, x_k) + \frac{\partial f}{\partial x}(t, x_k) f(t, x(t_k)) \right\}$$

Second order for SIR model



Calculate the second order term of the scheme for each component of the SIR model and add it to the R code! Idem for the SIR model with quadratic force of infection function! Compare!

- Second order scheme: accurate, but f needs to be differentiable \Rightarrow integral equation:

$$x(t_{k+1}) = x(t_k) + \int_{t_k}^{t_{k+1}} f(s)ds$$


- **Left endpoint** rule: $\int_{t_k}^{t_{k+1}} f(s)ds \approx (\Delta t)f(t_k)$ with (forward Euler) scheme:

$$x(t_{k+1}) = x(t_k) + (\Delta t)f(t_k, x^k)$$

Implicit numerical schemes: Trapezoidal rule

- **Trapezoidal rule:** $\int_{t_k}^{t_{k+1}} f(s)ds \approx (\Delta t) \frac{1}{2} (f(t_k, x(t_k)) + f(t_{k+1}, x(t_{k+1})))$ with **implicit** scheme

$$x(t_{k+1}) = x(t_k) + (\Delta t) \frac{1}{2} \left(f(t_k, x^k) + f(t_{k+1}, x^{k+1}) \right)$$



Group A work: Solve influenza SIR model numerically in R!



- Solve the SIR model numerically using the function `ode` in the package `deSolve` (e.g. find syntax on stackoverflow or ChatGPT)
- look up in the help menu `?ode` different methods and their required parameters 
`02_deSolveSIR.R`


Group B work: Do-it-yourself trapezoidal scheme!



- Solve the SIR model numerically by implementing the trapezoidal scheme in R!
~~Don't use ChatGPT~~ you can use ChatGPT, but explain the result. 
`03_SIR_trapezoidal.R`

Serial interval, generation time, incubation time

The figure below from Lehtinen et al. shows between infector i and infectee j :



- G , **generation time**: time between infection of i and j
- S , **serial interval**: time between symptom onset of i and j
- I , **incubation time**: time between infection of i and symptom onset of j

Generation time and serial interval

Which of the three quantities G, S, I can be negative? What is the epidemiological meaning in this case? Can you give an example?



- time increments $t_i = i \in \mathbb{N}$

- time increments $t_i = i \in \mathbb{N}$
- **generation time** distribution $g : \mathbb{N} \rightarrow [0, 1]$, i.e. $g(k)$ is probability of a primary infection causing a secondary infection after k time steps

- time increments $t_i = i \in \mathbb{N}$
- **generation time** distribution $g : \mathbb{N} \rightarrow [0, 1]$, i.e. $g(k)$ is probability of a primary infection causing a secondary infection after k time steps
- force of infection $\lambda(I)(i) = \beta \sum_k \frac{I(i-k)}{N(i-k)} g(k)$, **non-Markovian**

- time increments $t_i = i \in \mathbb{N}$
- **generation time** distribution $g : \mathbb{N} \rightarrow [0, 1]$, i.e. $g(k)$ is probability of a primary infection causing a secondary infection after k time steps
- force of infection $\lambda(I)(i) = \beta \sum_k \frac{I(i-k)}{N(i-k)} g(k)$, **non-Markovian**
- **Difference equation:** β, γ are probabilities

$$\begin{aligned} S(i+1) &= S(i) - \lambda(I)(i)S(i) \\ I(i+1) &= I(i) + \lambda(I)(i)S(i) - \gamma I(i) \\ R(i+1) &= \gamma I(i) \end{aligned}$$

- time increments $t_i = i \in \mathbb{N}$
- **generation time** distribution $g : \mathbb{N} \rightarrow [0, 1]$, i.e. $g(k)$ is probability of a primary infection causing a secondary infection after k time steps
- force of infection $\lambda(I)(i) = \beta \sum_k \frac{I(i-k)}{N(i-k)} g(k)$, **non-Markovian**
- **Difference equation:** β, γ are probabilities


$$S(i+1) = S(i) - \lambda(I)(i)S(i)$$


$$I(i+1) = I(i) + \lambda(I)(i)S(i) - \gamma I(i)$$

$$R(i+1) = \gamma I(i)$$

Update for next time step depends not only on now, but also past events!


event: infectious individual arrives






event: infectious individuals transmits, two infected

dormitory




event: the first infected individual recovers

dormitory




Why stochastic dynamics?


Many phenomena in biology are **intrinsically random** and **multi-scale**!




microbiology



biophysics




individual to population



- stochastic algorithms need **rules**, not explicit functions, **flexible**!
- stochastic algorithms explore **probabilistic** questions: extinction, criticality

"mean-field approximation" of deterministic equations by stochastic algorithm

Law of Large Numbers (LLN)



Mean of iid samples converges to expected value!

X_i iid r.v., then

$$\lim_{n \rightarrow \infty} \frac{1}{n}(X_1 + \dots + X_n) = \mathbb{E}(X_1)$$

strong LLN: a.s. convergence

weak LLN: convergence in probability

"mean-field approximation" of deterministic equations by stochastic algorithm

Central Limit Theorem (CLT)



Rescaled mean of iid samples with equal variance has Gaussian law as limit distribution!

X_i iid r.v. with $\text{var}(X_i) = \sigma^2$, and Y r.v. with law $\mathcal{N}(0, \sigma^2)$, then

$$\lim_{n \rightarrow \infty} \sqrt{n} \frac{1}{n} (X_1 + \dots + X_n) = Y$$

CLT: convergence in probability

stochastic=random=aleatory=chance=?

stochastic=random=aleatory=chance=?

axioms of probability: universe+events+probability

stochastic=random=aleatory=chance=?

axioms of probability: universe+events+probability

universe Ω : things, e.g. head, tail or infection, recovery

stochastic=random=aleatory=chance=?

axioms of probability: universe+events+probability

universe Ω : things, e.g. head, tail or infection, recovery

events \mathcal{F} , what can happen with things, e.g. head/tail in coin toss

stochastic=random=aleatory=chance=?

axioms of probability: universe+events+probability

universe Ω : things, e.g. head, tail or infection, recovery

events \mathcal{F} , what can happen with things, e.g. head/tail in coin toss

probability $\mathbb{P} : (\Omega, \mathcal{F}) \rightarrow [0, 1]$

stochastic=random=aleatory=chance=?

axioms of probability: universe+events+probability

universe Ω : things, e.g. head, tail or infection, recovery

events \mathcal{F} , what can happen with things, e.g. head/tail in coin toss

probability $\mathbb{P} : (\Omega, \mathcal{F}) \rightarrow [0, 1]$

Axiom 1 $\mathbb{P}(\Omega) = 1$

stochastic=random=aleatory=chance=?

axioms of probability: universe+events+probability

universe Ω : things, e.g. head, tail or infection, recovery

events \mathcal{F} , what can happen with things, e.g. head/tail in coin toss

probability $\mathbb{P} : (\Omega, \mathcal{F}) \rightarrow [0, 1]$

Axiom 1 $\mathbb{P}(\Omega) = 1$

Axiom 2 For any event E : $\mathbb{P}(E) = 1 - \mathbb{P}(\Omega \setminus E)$

stochastic=random=aleatory=chance=?

axioms of probability: universe+events+probability

universe Ω : things, e.g. head, tail or infection, recovery

events \mathcal{F} , what can happen with things, e.g. head/tail in coin toss

probability $\mathbb{P} : (\Omega, \mathcal{F}) \rightarrow [0, 1]$

Axiom 1 $\mathbb{P}(\Omega) = 1$

Axiom 2 For any event E : $\mathbb{P}(E) = 1 - \mathbb{P}(\Omega \setminus E)$

Axiom 3 if E_i disjoint, then $\mathbb{P}(\bigcup_i E_i) = \sum_i \mathbb{P}(E_i)$

stochastic=random=aleatory=chance=?

axioms of probability: universe+events+probability

universe Ω : things, e.g. head, tail or infection, recovery

events \mathcal{F} , what can happen with things, e.g. head/tail in coin toss

probability $\mathbb{P} : (\Omega, \mathcal{F}) \rightarrow [0, 1]$

Axiom 1 $\mathbb{P}(\Omega) = 1$

Axiom 2 For any event E : $\mathbb{P}(E) = 1 - \mathbb{P}(\Omega \setminus E)$

Axiom 3 if E_i disjoint, then $\mathbb{P}(\bigcup_i E_i) = \sum_i \mathbb{P}(E_i)$

random variable (r.v.) $X : (\Omega, \mathcal{F}, \mathbb{P}) \rightarrow (A, \mathcal{A})$ measurable

stochastic=random=aleatory=chance=?

axioms of probability: universe+events+probability

universe Ω : things, e.g. head, tail or infection, recovery

events \mathcal{F} , what can happen with things, e.g. head/tail in coin toss

probability $\mathbb{P} : (\Omega, \mathcal{F}) \rightarrow [0, 1]$

Axiom 1 $\mathbb{P}(\Omega) = 1$

Axiom 2 For any event E : $\mathbb{P}(E) = 1 - \mathbb{P}(\Omega \setminus E)$

Axiom 3 if E_i disjoint, then $\mathbb{P}(\bigcup_i E_i) = \sum_i \mathbb{P}(E_i)$

random variable (r.v.) $X : (\Omega, \mathcal{F}, \mathbb{P}) \rightarrow (A, \mathcal{A})$ measurable

probability law of r.v. $f_X : (A, \mathcal{A}) \rightarrow [0, 1]$ with

$f(A) = \mathbb{P}(X^{-1}(A)) = \mathbb{P}(E \in \mathcal{F} : X(E) = A)$ for $A \in \mathcal{A}$

Coin toss



Write down a **fair** coin toss as a r.v. using the definitions from above!

Discrete & continuous r.v.



discrete r.v. universe is countable or finite

Coin toss



Write down a **fair** coin toss as a r.v. using the definitions from above!

Discrete & continuous r.v.



discrete r.v. universe is countable or finite

$$\Omega = \{0, 1\} \text{ and } f_X(\{1\}) = p \in [0, 1] \text{ "Bernoulli"}$$

Coin toss



Write down a **fair** coin toss as a r.v. using the definitions from above!

Discrete & continuous r.v.



discrete r.v. universe is countable or finite

$\Omega = \{0, 1\}$ and $f_X(\{1\}) = p \in [0, 1]$ "Bernoulli"

$\Omega = \{0, 1, \dots\}$ and $f_X(\{k\}) = e^{-\lambda} \frac{\lambda^k}{k!}$ "Poisson"

Coin toss



Write down a **fair** coin toss as a r.v. using the definitions from above!

Discrete & continuous r.v.



discrete r.v. universe is countable or finite

$\Omega = \{0, 1\}$ and $f_X(\{1\}) = p \in [0, 1]$ "Bernoulli"

$\Omega = \{0, 1, \dots\}$ and $f_X(\{k\}) = e^{-\lambda} \frac{\lambda^k}{k!}$ "Poisson"

continuous r.v. universe is uncountable

Coin toss



Write down a **fair** coin toss as a r.v. using the definitions from above!

Discrete & continuous r.v.



discrete r.v. universe is countable or finite

$\Omega = \{0, 1\}$ and $f_X(\{1\}) = p \in [0, 1]$ "Bernoulli"

$\Omega = \{0, 1, \dots\}$ and $f_X(\{k\}) = e^{-\lambda} \frac{\lambda^k}{k!}$ "Poisson"

continuous r.v. universe is uncountable

$\Omega = \mathbb{R}_+$ and $f_X([0, a]) = \lambda \int_0^a e^{-\lambda y} dy$, but $f_X(\{b\}) = 0!$
"exponential distribution"

From random events to stochastic dynamics

observable $\varphi : A \rightarrow \mathbb{R}$, then $\varphi(X)$ observable

From random events to stochastic dynamics

observable $\varphi : A \rightarrow \mathbb{R}$, then $\varphi(X)$ observable

expectation : $\mathbb{E}(\varphi(X)) := \sum_z \varphi(z) f_X(z)$ resp. $\int_A \varphi(y) f_X(y) dy$

observable $\varphi : A \rightarrow \mathbb{R}$, then $\varphi(X)$ observable

expectation : $\mathbb{E}(\varphi(X)) := \sum_z \varphi(z) f_X(z)$ resp. $\int_A \varphi(y) f_X(y) dy$

moments $f(x) = x^n$, then $\mathbb{E}(X^n)$ is nth-moment, moments fully determine probability law of r.v.!

observable $\varphi : A \rightarrow \mathbb{R}$, then $\varphi(X)$ observable

expectation : $\mathbb{E}(\varphi(X)) := \sum_z \varphi(z) f_X(z)$ resp. $\int_A \varphi(y) f_X(y) dy$

moments $f(x) = x^n$, then $\mathbb{E}(X^n)$ is nth-moment, moments fully determine probability law of r.v.!

independence X, Y r.v. are independent $X \perp Y$, iff

$$\mathbb{P}(\{X \in A\} \cap \{Y \in B\}) = \mathbb{P}(X \in A)\mathbb{P}(Y \in B)$$

observable $\varphi : A \rightarrow \mathbb{R}$, then $\varphi(X)$ observable

expectation : $\mathbb{E}(\varphi(X)) := \sum_z \varphi(z) f_X(z)$ resp. $\int_A \varphi(y) f_X(y) dy$

moments $f(x) = x^n$, then $\mathbb{E}(X^n)$ is nth-moment, moments fully determine probability law of r.v.!

independence X, Y r.v. are independent $X \perp Y$, iff

$$\mathbb{P}(\{X \in A\} \cap \{Y \in B\}) = \mathbb{P}(X \in A)\mathbb{P}(Y \in B)$$

iid X, Y iid if independent, identically distributed

observable $\varphi : A \rightarrow \mathbb{R}$, then $\varphi(X)$ observable

expectation : $\mathbb{E}(\varphi(X)) := \sum_z \varphi(y) f_X(y)$ resp. $\int_A \varphi(y) f_X(y) dy$

moments $f(x) = x^n$, then $\mathbb{E}(X^n)$ is nth-moment, moments fully determine probability law of r.v.!

independence X, Y r.v. are independent $X \perp Y$, iff

$$\mathbb{P}(\{X \in A\} \cap \{Y \in B\}) = \mathbb{P}(X \in A)\mathbb{P}(Y \in B)$$

iid X, Y iid if independent, identically distributed

Warm-up



Calculate the expectation of Bernoulli and exponential r.v.!

From random events to stochastic dynamics

stochastic process collection of r.v. indexed by time: $t \mapsto X(t)$

stochastic process collection of r.v. indexed by time: $t \mapsto X(t)$

filtration \mathcal{F}_t collection of sets of events indexed by time, information about $X(t)$ that is available up to time t

convergence almost sure \rightarrow in probability \rightarrow in expectation

stochastic process collection of r.v. indexed by time: $t \mapsto X(t)$

filtration \mathcal{F}_t collection of sets of events indexed by time, information about $X(t)$ that is available up to time t

convergence almost sure \rightarrow in probability \rightarrow in expectation

Markov property $\mathbb{P}(X_{t+a} \in A | \mathcal{F}_t) = \mathbb{P}(X_{t+a} \in A | \sigma(X_t))$
"what happens in the future depends only on the present state"

Markov chain $\mathbb{P}(X_{n+1} = x_{n+1} | X_n = x_n, \dots, X_1 = x_1) = \mathbb{P}(X_{n+1} = x_{n+1} | X_n = x_n)$
"what happens in the future depends only on the present state"

From memory-less processes to differential equations

The r.v. X in \mathbb{R}_+ is **without memory** if:

$$\mathbb{P}(X > t + s | X > s) = \mathbb{P}(X > t)$$

The r.v. X in \mathbb{R}_+ is **without memory** if:

$$\mathbb{P}(X > t + s | X > s) = \mathbb{P}(X > t)$$

with

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

The r.v. X in \mathbb{R}_+ is **without memory** if:

$$\mathbb{P}(X > t + s | X > s) = \mathbb{P}(X > t)$$

with

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

\iff

$$\mathbb{P}(X > t + s) = \mathbb{P}(X > t)\mathbb{P}(X > s)$$

The r.v. X in \mathbb{R}_+ is **without memory** if:

$$\mathbb{P}(X > t + s | X > s) = \mathbb{P}(X > t)$$

with

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

\iff

$$\mathbb{P}(X > t + s) = \mathbb{P}(X > t)\mathbb{P}(X > s)$$

\iff functional equation:

$$\mathbb{P}(X > a) = \mathbb{P}(X > 1)^a = e^{\log(\mathbb{P}(X > 1))a} = e^{-\lambda a}$$

The r.v. X in \mathbb{R}_+ is **without memory** if:

$$\mathbb{P}(X > t + s | X > s) = \mathbb{P}(X > t)$$

with

$$\mathbb{P}(A|B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}$$

\iff

$$\mathbb{P}(X > t + s) = \mathbb{P}(X > t)\mathbb{P}(X > s)$$

\iff functional equation:

$$\mathbb{P}(X > a) = \mathbb{P}(X > 1)^a = e^{\log(\mathbb{P}(X > 1))a} = e^{-\lambda a}$$

Memory-less r.v.



X r.v. **without memory** $\iff X$ exponentially distr. with $\lambda = -\log(\mathbb{P}(X > 1))$


A MILLION Random Digits

WITH
100,000 Normal Deviates

buy the book... RAND

Random number generation: the physical way

...or move your mouse: hardware random number generator

A screenshot of a terminal window titled "bestuser@workstation:~". The window contains the following text:

```
bestuser@workstation:~  
File Edit View Search Terminal Help  
"Best User (Best Company) <bestuser@example.com>"  
  
Change (N)ame, (C)omment, (E)mail or (O)kay/(Q)uit? O  
We need to generate a lot of random bytes. It is a good idea to perform  
some other action (type on the keyboard, move the mouse, utilize the  
disks) during the prime generation; this gives the random number  
generator a better chance to gain enough entropy.  
We need to generate a lot of random bytes. It is a good idea to perform  
some other action (type on the keyboard, move the mouse, utilize the  
disks) during the prime generation; this gives the random number  
generator a better chance to gain enough entropy.  
gpg: /home/bestuser/.gnupg/trustdb.gpg: trustdb created  
gpg: key 94F45C144CD3559D marked as ultimately trusted  
gpg: directory '/home/bestuser/.gnupg/openpgp-revocs.d' created  
gpg: revocation certificate stored as '/home/bestuser/.gnupg/openpgp-revocs.d/CC  
1795E6F83B091A7B813A6D94F45C144CD3559D.rev'  
public and secret key created and signed.  
  
pub    rsa2048 2020-04-23 [SC] [expires: 2021-04-23]  
      CC1795E6F83B091A7B813A6D94F45C144CD3559D  
uid    Best User (Best Company) <bestuser@example.com>  
sub    rsa2048 2020-04-23 [E] [expires: 2021-04-23]  
  
[bestuser@workstation ~]$
```

The terminal window has a dark background with light-colored text. The user's command history is visible at the bottom of the window.

- **pseudo-random** number generator: linear congruence, Mersenne Twister

Random number generation: the algorithmic way

- **pseudo-random** number generator: linear congruence, Mersenne Twister
- **quasi-random** number generator: low-discrepancy sequence

- **pseudo-random** number generator: linear congruence, Mersenne Twister
- **quasi-random** number generator: low-discrepancy sequence
- statistical tests of randomness

- **pseudo-random** number generator: linear congruence, Mersenne Twister
- **quasi-random** number generator: low-discrepancy sequence
- statistical tests of randomness
- **seed** of random number generation, index used to "replicate" simulation

Random number generation: the algorithmic way

- **pseudo-random** number generator: linear congruence, Mersenne Twister
- **quasi-random** number generator: low-discrepancy sequence
- statistical tests of randomness
- **seed** of random number generation, index used to "replicate" simulation
- cumulative distribution function (cdf) $F_X(t) = \mathbb{P}(X \leq t)$, inverse cdf (icdf) F_X^{-1}

- **pseudo-random** number generator: linear congruence, Mersenne Twister
- **quasi-random** number generator: low-discrepancy sequence
- statistical tests of randomness
- **seed** of random number generation, index used to "replicate" simulation
- cumulative distribution function (cdf) $F_X(t) = \mathbb{P}(X \leq t)$, inverse cdf (icdf) F_X^{-1}

inverse transform sampling



X real-valued r.v. and U uniformly distributed r.v. on $[0, 1]$, then r.v. $F_X^{-1}(U) \sim X$

Sample from exponential distribution



random draws for r.v. $X \sim \text{Exp}(\lambda)$ by using uniformly distributed random numbers in the interval $[0, 1]$

cdf $F_X(t) = 1 - e^{-\lambda t}$ \Rightarrow **icdf** $F_X^{-1}(t) = -\log(-(t - 1))/\lambda = \frac{\log(1-t)}{-\lambda}$ 

04_inversesampling.R

Sample from standard normal distribution



random draws for r.v. $X \sim \mathcal{N}(0, 1)$ by using uniformly distributed random numbers in the interval $[0, 1]$

Sample from discrete set




X is r.v. with values in discrete set $K = \{k_1, k_2, \dots\}$ with $\mathbb{P}(X = k_i) = p_i$ such that $\sum_i p_i = 1$.

Define $g : [0, 1] \rightarrow K$ with

$$g(x) = k_j \Leftrightarrow \sum_{i=1}^{j-1} p_i < x \leq \sum_{i=1}^j p_i$$

If $U \sim \text{uniform}$, then $X \sim g(U)$:


$$\mathbb{P}(g(U) = k_j) = \mathbb{P}\left(\sum_{i=1}^{j-1} p_i < U \leq \sum_{i=1}^j p_i\right) = p_j$$



How to measure computational complexity

- Landau notation: $f(n) = \mathcal{O}(g(n))$ for $n \rightarrow \infty$ if there are $M, n_0 > 0$ such that for all $n \geq n_0$:

$$|f(n)| \leq Mg(n)$$




How to measure computational complexity

- Landau notation: $f(n) = \mathcal{O}(g(n))$ for $n \rightarrow \infty$ if there are $M, n_0 > 0$ such that for all $n \geq n_0$:

$$|f(n)| \leq Mg(n)$$

- e.g. $f(n) = 5n^3 + n + 5 \Rightarrow f(n) = \mathcal{O}(n^3)$




How to measure computational complexity

- Landau notation: $f(n) = \mathcal{O}(g(n))$ for $n \rightarrow \infty$ if there are $M, n_0 > 0$ such that for all $n \geq n_0$:

$$|f(n)| \leq Mg(n)$$

- e.g. $f(n) = 5n^3 + n + 5 \Rightarrow f(n) = \mathcal{O}(n^3)$
- **runtime** of algorithm: input size n , algorithm needs $\mathcal{O}(g(n))$ computation time for solution




How to measure computational complexity

- Landau notation: $f(n) = \mathcal{O}(g(n))$ for $n \rightarrow \infty$ if there are $M, n_0 > 0$ such that for all $n \geq n_0$:

$$|f(n)| \leq Mg(n)$$

- e.g. $f(n) = 5n^3 + n + 5 \Rightarrow f(n) = \mathcal{O}(n^3)$
- **runtime** of algorithm: input size n , algorithm needs $\mathcal{O}(g(n))$ computation time for solution
- e.g. **binary search** in list of size n has **logarithmic** run time, i.e. algorithm needs $\mathcal{O}(\log n)$ computation steps for solution



Benchmarking: compare the computing time of programs with same input/output

Benchmarking: compare the computing time of programs with same input/output

Recursive vs dynamic programming



The Fibonacci numbers are defined by the recursion:

$$F_1 = 1, F_2 = 1, F_n = F_{n-1} + F_{n-2}$$

for $n > 2$. Calculate F_n by both recursive and dynamic programming (i.e. using already stored numbers). Use the R package `microbenchmark` to benchmark both functions and `system.time` to calculate the runtime as a function of n . What do you observe?  `05_benchmarking.R`

Profiling: diagnosing required memory, frequency and duration of functional calls for each line of your computer code

Profiling: diagnosing required memory, frequency and duration of functional calls for each line of your computer code

Profiling



Use the R function `Rprof` to profile both implementations of the Fibonacci number calculations. What do you observe?  [06_profiling.R](#)

Profiling: diagnosing required memory, frequency and duration of functional calls for each line of your computer code

Profiling



Use the R function `Rprof` to profile both implementations of the Fibonacci number calculations. What do you observe?  [06_profiling.R](#)

Cyclomatic complexity: number of linearly independent paths through code

Profiling: diagnosing required memory, frequency and duration of functional calls for each line of your computer code

Profiling



Use the R function `Rprof` to profile both implementations of the Fibonacci number calculations. What do you observe?  `06_profiling.R`

Cyclomatic complexity: number of linearly independent paths through code

Cyclomatic complexity



Use the R package `cyclocomp` to profile both implementations of the Fibonacci number calculations. What do you observe? What are your conclusion?  `07_cyclocomp.R`

- species \mathcal{S} : chemical compounds whose dynamics we model
- reactions \mathcal{R} : how to convert one complex into another

Example

$$\mathcal{S} = \{A, B, C\}, \mathcal{C} = \{A + B, 2B, C, \emptyset\}, \mathcal{R} = \{A + B \rightarrow 2B, B \rightarrow C, C \rightarrow \emptyset\}$$

- species \mathcal{S} : chemical compounds whose dynamics we model
- complexes \mathcal{C} : nonnegative **linear** combinations of species (i.e. interactions)
- reactions \mathcal{R} : how to convert one complex into another

Example

$$\mathcal{S} = \{A, B, C\}, \mathcal{C} = \{A + B, 2B, C, \emptyset\}, \mathcal{R} = \{A + B \rightarrow 2B, B \rightarrow C, C \rightarrow \emptyset\}$$

- species \mathcal{S} : chemical compounds whose dynamics we model
- complexes \mathcal{C} : nonnegative **linear** combinations of species (i.e. interactions)
- reactions \mathcal{R} : how to convert one complex into another

Example

$$\mathcal{S} = \{A, B, C\}, \mathcal{C} = \{A + B, 2B, C, \emptyset\}, \mathcal{R} = \{A + B \rightarrow 2B, B \rightarrow C, C \rightarrow \emptyset\}$$

- $A + B \rightarrow 2B$: A active, B inactive form of protein, B catalyzes A

- species \mathcal{S} : chemical compounds whose dynamics we model
- complexes \mathcal{C} : nonnegative **linear** combinations of species (i.e. interactions)
- reactions \mathcal{R} : how to convert one complex into another

Example

$$\mathcal{S} = \{A, B, C\}, \mathcal{C} = \{A + B, 2B, C, \emptyset\}, \mathcal{R} = \{A + B \rightarrow 2B, B \rightarrow C, C \rightarrow \emptyset\}$$

- $A + B \rightarrow 2B$: A active, B inactive form of protein, B catalyzes A
- $B \rightarrow C$: B undergoes conformational change to become C

- species \mathcal{S} : chemical compounds whose dynamics we model
- complexes \mathcal{C} : nonnegative **linear** combinations of species (i.e. interactions)
- reactions \mathcal{R} : how to convert one complex into another

Example

$$\mathcal{S} = \{A, B, C\}, \mathcal{C} = \{A + B, 2B, C, \emptyset\}, \mathcal{R} = \{A + B \rightarrow 2B, B \rightarrow C, C \rightarrow \emptyset\}$$

- $A + B \rightarrow 2B$: A active, B inactive form of protein, B catalyzes A
- $B \rightarrow C$: B undergoes conformational change to become C
- $C \rightarrow \emptyset$: C is degraded

- $\mathcal{R} = \{y_k \rightarrow y'_k; y_k, y'_k \in \mathcal{C}\}$ with $y_k \equiv \sum_i y_{k,i} S_i$
- **stoichiometric** vectors of network: $\zeta_k := y'_k - y_k \in \mathbb{Z}^n$

Example

$$\mathcal{S} = \{A, B, C\}, \mathcal{C} = \{A + B, 2B, C, \emptyset\}, \mathcal{R} = \{A + B \rightarrow 2B, B \rightarrow C, C \rightarrow \emptyset\}$$

$$\zeta_1 = [0, 2, 0] - [1, 1, 0] = [-1, 1, 0]$$

$$\zeta_2 = [0, 0, 1] - [0, 1, 0] = [0, -1, 1]$$

$$\zeta_3 = [0, 0, 0] - [0, 0, 1] = [0, 0, -1]$$

Counting processes via Poisson

Counting process:

$t \mapsto N(t) \in \mathbb{N}$ such that $N(0) = 0$, N constant except jumps of size +1.


Poisson (point) process



Counting process such that

- $N(0) = 0$
- independent increments, i.e. $N(t_{k+1}) - N(t_k)$ are independent r.v. for $0 < t_1 < \dots < t_{k+1}$
- distribution of $N(t + \Delta) - N(t)$ does not depend on t
it follows: $\mathbb{P}(N(t) = k) = \frac{(\lambda t)^k}{k!} e^{-\lambda t}$ "number of arrivals until time t "
- jump time $S_k = \min\{t : N(t) \geq k\}$, then $S_k - S_{k-1} \sim \text{Exp}(\lambda)$

Poisson process



Counting processes and biochemical reactions

- $R_k(t)$ counting process for occurrences of reaction k by time t

Counting processes and biochemical reactions

- $R_k(t)$ counting process for occurrences of reaction k by time t
- dynamical system of molecules

$$X(t) = X(0) + \sum_k R_k(t) \zeta_k$$

Counting processes and biochemical reactions

- $R_k(t)$ counting process for occurrences of reaction k by time t
- dynamical system of molecules

$$X(t) = X(0) + \sum_k R_k(t) \zeta_k$$

Reaction dynamics as Markov jump processes



Let $\lambda_k : \mathbb{N}^S \rightarrow \mathbb{R}_+$ be intensity function of reaction k for given molecular state.

The counting processes R_k can be represented by iid Poisson processes Y_k with intensity 1 such that for **intensity function** $\lambda_k : \mathbb{N}^S \rightarrow \mathbb{R}_+$:

$$R^k(t) = Y_k \left(\int_0^t \lambda_k(X(s)) ds \right)$$



Anderson & Kurtz, chapter 1, pp5

Mass-action kinetics, seen by the chemist



At constant temperature, the rate of chemical reaction is directly proportional to the product of molar concentrations of reacting species.

Mass-action kinetics, seen by the chemist



At constant temperature, the rate of chemical reaction is directly proportional to the product of molar concentrations of reacting species.

Mass-action kinetics, seen by the mathematician



$$\lambda_k(x) = \kappa_k \prod_i \frac{x_i!}{(x_i - y_{ki})!}$$

x_i = #species i , y_{ki} = #species i needed for reaction k , "falling factorial"

Mass-action kinetics, seen by the mathematician



$$\lambda_k(x) = \kappa_k \prod_i \frac{x_i!}{(x_i - y_{ki})!}$$

x_i = #species i , y_{ki} = #species i needed for reaction k , "falling factorial"

Mass-action kinetics, seen by the mathematician



$$\lambda_k(x) = \kappa_k \prod_i \frac{x_i!}{(x_i - y_{ki})!}$$

x_i = #species i , y_{ki} = #species i needed for reaction k , "falling factorial"

Mass-action kinetics, seen by the mathematician



λ_k is proportional to the number of distinct subsets of the molecules present that can form the inputs for the reaction. E.g. for reaction $A + B \rightarrow 2B$, $\lambda_1(x) = \kappa_1 x_1 x_2$.

Beyond mass-action kinetics



Think of physical or chemical reasons that could prevent the validity of the principle of mass-action!

Beyond mass-action kinetics



Think of physical or chemical reasons that could prevent the validity of the principle of mass-action!

Beyond mass-action kinetics



Explain the concept of **cooperative binding** and how it would change the assumptions on biochemical reaction dynamics!

Beyond mass-action kinetics



Think of physical or chemical reasons that could prevent the validity of the principle of mass-action!

Beyond mass-action kinetics



Explain the concept of **cooperative binding** and how it would change the assumptions on biochemical reaction dynamics!

Beyond mass-action kinetics



Can you give an example for **non**-mass-action kinetics in epidemic processes?

stoichiometrically admissible: $\lambda_k(x) = 0$ if $x_i < y_{k,i}$ for all i (e.g. for $A + B \rightarrow 2B$ we need at least one A and one B for the reaction to happen)

stoichiometrically admissible: $\lambda_k(x) = 0$ if $x_i < y_{k,i}$ for all i (e.g. for $A + B \rightarrow 2B$ we need at least one A and one B for the reaction to happen)

Reaction	Intensity function	Rate	molecular
$A + B \rightarrow 2B$	$\lambda_1(x) = \kappa_1 x_1 x_2$	κ_1	catalysis of protein inactivation
$B \rightarrow C$	$\lambda_2(x) = \kappa_2 x_2$	κ_2	conformational change
$C \rightarrow \emptyset$	$\lambda_3(x) = \kappa_3 x_3$	κ_3	degradation

Epidemics as biochemical reaction systems

stoichiometrically admissible: $\lambda_k(x) = 0$ if $x_i < y_{k,i}$ for all i (e.g. for $A + B \rightarrow 2B$ we need at least one A and one B for the reaction to happen)

Reaction	Intensity function	Rate	molecular
$A + B \rightarrow 2B$	$\lambda_1(x) = \kappa_1 x_1 x_2$	κ_1	catalysis of protein inactivation
$B \rightarrow C$	$\lambda_2(x) = \kappa_2 x_2$	κ_2	conformational change
$C \rightarrow \emptyset$	$\lambda_3(x) = \kappa_3 x_3$	κ_3	degradation

Reaction	Intensity function	Rate	molecular	epi
$S + I \rightarrow 2I$	$\lambda_1(S, I, R) = \beta SI$	β	catalysis of protein inactivation	new infections
$I \rightarrow R$	$\lambda_2(S, I, R) = \gamma I$	γ	conformational change	recovery
$R \rightarrow \emptyset$	$\lambda_3(S, I, R) = \delta R$	δ	degradation	death

Reaction	Intensity function	Rate	Stoichiometry ζ	epi
$S + I \rightarrow 2I$	$\lambda_1(S, I, R) = \beta SI$	β	$[-1, 1, 0]$	new infections
$I \rightarrow R$	$\lambda_2(S, I, R) = \gamma I$	γ	$[0, -1, 1]$	recovery

Reaction	Intensity function	Rate	Stoichiometry ζ	epi
$S + I \rightarrow 2I$	$\lambda_1(S, I, R) = \beta SI$	β	$[-1, 1, 0]$	new infections
$I \rightarrow R$	$\lambda_2(S, I, R) = \gamma I$	γ	$[0, -1, 1]$	recovery

- time evolution of molecules per species is given by solution of the equation

$$X(t) = X(0) + Y^1 \left(\int_0^t \lambda_1(X_s) ds \right) \zeta_1 + Y^2 \left(\int_0^t \lambda_2(Y_s) ds \right) \zeta_2$$

Reaction	Intensity function	Rate	Stoichiometry ζ	epi
$S + I \rightarrow 2I$	$\lambda_1(S, I, R) = \beta SI$	β	$[-1, 1, 0]$	new infections
$I \rightarrow R$	$\lambda_2(S, I, R) = \gamma I$	γ	$[0, -1, 1]$	recovery

- time evolution of molecules per species is given by solution of the equation

$$X(t) = X(0) + Y^1 \left(\int_0^t \lambda_1(X_s) ds \right) \zeta_1 + Y^2 \left(\int_0^t \lambda_2(Y_s) ds \right) \zeta_2$$

$$\begin{bmatrix} S(t) \\ I(t) \\ R(t) \end{bmatrix} = \begin{bmatrix} S(0) \\ I(0) \\ R(0) \end{bmatrix} + Y^1 \left(\int_0^t \beta S(s) I(s) ds \right) \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} + Y^2 \left(\int_0^t \gamma I(s) ds \right) \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

Group work: simulate variations of the SIR models

For each of the examples draw the flow diagram for the disease dynamics, write the biochemical reaction network and stoichiometric vectors!

Incubation period



From exposure to infectiousness, 5 days pass on average. Add a compartment for exposed but not yet infectious hosts!

Ebola-like dynamics



In addition to the basic model used for influenza, we consider also a fraction p of individuals to die from the disease. Contact of susceptibles with dead bodies before burial will lead to additional infections.

For each of the examples draw the flow diagram for the disease dynamics, write the biochemical reaction network and stoichiometric vectors!

Two rooms in the dorm




In the boarding school there is a respiratory disease outbreak among the students. All of them live together in the same building, but there are two dormitories A and B . Students living in B prefer to visit those living in A , but not so much the other way around. We have seen the following contact rates:



Group work: solutions

For infectivity $\beta_A > \beta_B$ and recovery rate γ , the flow diagram reads:



The **species** are $\{S_A, I_A, R_A, S_B, I_B, R_B\}$, the **complexes** are $\{S_A + I_A, 2I_A, R_A, S_B + I_B, 2I_B, R_B, S_A + I_B, I_A + I_B, S_B + I_A\}$, the **reactions** are $R_1 : S_A + I_A \rightarrow 2I_A$, $R_2 : I_A \rightarrow R_A$, $R_3 : S_B + I_B \rightarrow 2I_B$, $R_4 : I_B \rightarrow R_B$, $R_5 : S_A + I_B \rightarrow I_A + I_B$, $R_6 : S_B + I_A \rightarrow I_A + I_B$ and the stoichiometric matrix is

$$\begin{bmatrix} -1 & 0 & 0 & 0 & -1 & 0 \\ -1 & -1 & 0 & 0 & +1 & 0 \\ 0 & +1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & -1 \\ 0 & 0 & +1 & -1 & 0 & +1 \\ 0 & 0 & 0 & +1 & 0 & 0 \end{bmatrix}$$

From Markov jump processes to differential equations

- replace molecule numbers $X(t)$ by **concentration** $C(t) = \frac{1}{N}X(t)$

- replace molecule numbers $X(t)$ by **concentration** $C(t) = \frac{1}{N}X(t)$
- N total number of molecules at given volume (e.g. Avogadro's number \times volume v , or total population)

- replace molecule numbers $X(t)$ by **concentration** $C(t) = \frac{1}{N}X(t)$
- N total number of molecules at given volume (e.g. Avogadro's number \times volume v , or total population)
- reaction rate inversely proportional to volume

- replace molecule numbers $X(t)$ by **concentration** $C(t) = \frac{1}{N}X(t)$
- N total number of molecules at given volume (e.g. Avogadro's number \times volume v , or total population)
- reaction rate inversely proportional to volume

Concentration dynamics



$$C^N(t) = C^N(0) + \sum_{k=1} N^{-1} Y_k \left(N \int_0^t \lambda_k(C^N(s)) ds \right) \zeta_k$$

From Markov jump processes to differential equations

- replace molecule numbers $X(t)$ by **concentration** $C(t) = \frac{1}{N_v} X(t)$

From Markov jump processes to differential equations

- replace molecule numbers $X(t)$ by **concentration** $C(t) = \frac{1}{N_v} X(t)$
- N_v total number of molecules at given volume (e.g. Avogadro's number \times volume v , or total population)

- replace molecule numbers $X(t)$ by **concentration** $C(t) = \frac{1}{N_v} X(t)$
- N_v total number of molecules at given volume (e.g. Avogadro's number \times volume v , or total population)
- reaction rates vary inversely with volume

- replace molecule numbers $X(t)$ by **concentration** $C(t) = \frac{1}{N_v} X(t)$
- N_v total number of molecules at given volume (e.g. Avogadro's number \times volume v , or total population)
- reaction rates vary inversely with volume

Concentration dynamics



$$C^{N_v}(t) := C^{N_v}(0) + \sum_{i=1}^m N_v^{-1} Y_k(N_v \int_0^t \lambda_k(C^{N_v}(s)) ds) \zeta_k$$

From Markov jump processes to differential equations

- what happens if $N \rightarrow \infty$?

From Markov jump processes to differential equations

- what happens if $N \rightarrow \infty$?
- $F(x) := \sum_k \lambda_k(x) \zeta_k$ globally Lipschitz

From Markov jump processes to differential equations

- what happens if $N \rightarrow \infty$?
- $F(x) := \sum_k \lambda_k(x) \zeta_k$ globally Lipschitz
- deterministic integral equation

$$x(t) = x(0) + \int_0^t F(x(s)) ds \quad (8)$$

- what happens if $N \rightarrow \infty$?
- $F(x) := \sum_k \lambda_k(x) \zeta_k$ globally Lipschitz
- deterministic integral equation

$$x(t) = x(0) + \int_0^t F(x(s)) ds \quad (8)$$

Convergence theorem



$$\lim_{N \rightarrow \infty} \mathbb{P}\left(\sup_{s \leq t} |C^N(s) - x(s)| \geq \epsilon\right) = 0$$

for each $\epsilon, t > 0$, weak law of large numbers

- what happens if $N \rightarrow \infty$?
- $F(x) := \sum_k \lambda_k(x) \zeta_k$ globally Lipschitz
- deterministic integral equation

$$x(t) = x(0) + \int_0^t F(x(s)) ds \tag{8}$$

Convergence theorem



$$\lim_{N \rightarrow \infty} \mathbb{P}(\sup_{s \leq t} |C^N(s) - x(s)| \geq \epsilon) = 0$$

for each $\epsilon, t > 0$, weak law of large numbers

- Proof built on Gronwall & Doob inequalities and martingale theory:  Anderson & Kurtz, page 44f

- **species** of molecules can form complexes, and changes between complexes define reactions

- **species** of molecules can form complexes, and changes between complexes define reactions
- **reactions** can be described by stoichiometric vectors
- under the **mass action assumption**, the rate of reaction is proportional to the number of molecules involved

- **species** of molecules can form complexes, and changes between complexes define reactions
- **reactions** can be described by stoichiometric vectors
- under the **mass action assumption**, the rate of reaction is proportional to the number of molecules involved
- the evolution of molecules over time can be described mathematically as a Poisson process

- **species** of molecules can form complexes, and changes between complexes define reactions
- **reactions** can be described by stoichiometric vectors
- under the **mass action assumption**, the rate of reaction is proportional to the number of molecules involved
- the evolution of molecules over time can be described mathematically as a Poisson process
- molecular **concentrations converge** towards deterministic limit as number of molecules goes to infinity

Back to epidemic processes



For the simple SIR model, show that assumptions of the convergence theorem are satisfied.

Write explicitly the integral equation (8) and show how it relates to the ODE system.

Gillespie's algorithm: direct method

Simulate systems of biochemical **reactions** (e.g. susceptible meets infectious), assuming no more than two individuals at a time are involved in the reaction or events:

- ① reminder:

$$X(t) = X(0) + Y^1 \left(\int_0^t \lambda_1(X_s) ds \right) \zeta_1 + Y^2 \left(\int_0^t \lambda_2(Y_s) ds \right) \zeta_2$$

- ② suppose transition times between states t_i , define $X(t) = X(t_k)$ for $t \in [t_k, t_{k+1})$

- ③ initial condition $X(0) = x_0$

- 1 for $t = t_k$ calculate $\lambda_k(X_t)$ for all k

- 2 τ time to next event follows $\text{Exp}(\sum_k \lambda_k(X_t))$

- 3 next event K sampled with $\frac{\lambda_k(X_t)}{\sum_k \lambda_k(X_t)}$

- 4 update time: $t_{k+1} = t_k + \tau$


- 5 update state: $X(t_{k+1}) = X_{t_k} + \zeta_K$

Gillespie's algorithm: direct method for SIR

```
1: Initial trajectory  $\mathcal{T} = (t, S, I, R) = (0, 762, 1, 0)$ 
2: while  $I > 0$  do
3:   Current state  $S$  last row of  $\mathcal{T}$ ,  $S = \mathcal{S}[2]$ ,  $I = \mathcal{S}[3]$ ,  $R = \mathcal{S}[4]$ 
4:   possible events vector:  $\mathcal{E} = (\underbrace{\text{new infection}, \dots, \text{new infection}}_{S \text{ times}}, \underbrace{\text{clearance}, \dots, \text{clearance}}_{I \text{ times}})$ 
5:   rates vector:  $\lambda = (\underbrace{\beta I / N, \dots, \beta I / N}_{S \text{ times}}, \underbrace{\gamma, \dots, \gamma}_{I \text{ times}})$ 
6:   time to next event: draw sample  $\tau$  from  $\text{Exp}(\sum_i \lambda_i)$ 
7:   choose next event: sample from  $\mathcal{E}$  with probability  $\frac{\lambda_i}{\sum_i \lambda_i}$ 
8:   if next event is "new infection" then
9:      $S \leftarrow S + (\tau, -1, 1, 0)$ 
10:    else if next event is "clearance" then
11:       $S \leftarrow S + (\tau, 0, -1, 1)$ 
12:    end if
13:     $\mathcal{T} \leftarrow [\mathcal{T}, S]$ 
14:  end while
15: return  $\mathcal{T}$ 
```

- write the Gillespie direct method in R for an SIR model
- use the optimal parameters obtained for the ODE system: $\beta = 1.6692258, \gamma = 0.4434502$
- perform 100 realizations of the stochastic process and compare to the ODE solution
- does the law of large numbers hold?
-  08_GillespieDirect.R

Gillespie's direct method for influenza SIR model



What happens if you choose $\beta = 0.7$?

Gillespie's algorithm: first reaction method for SIR

```
1: Initial trajectory  $\mathcal{T} = (t, S, I, R) = (0, 762, 1, 0)$ 
2: while  $I > 0$  do
3:   Current state  $S$  last row of  $\mathcal{T}$ ,  $S = \mathcal{S}[2]$ ,  $I = \mathcal{S}[3]$ ,  $R = \mathcal{S}[4]$ 
4:   possible events vector:  $\mathcal{E} = (\underbrace{\text{new infection}, \dots, \text{new infection}}_{S \text{ times}}, \underbrace{\text{clearance}, \dots, \text{clearance}}_{I \text{ times}})$ 
5:   rates vector:  $\lambda = (\underbrace{\beta I / N, \dots, \beta I / N}_{S \text{ times}}, \underbrace{\gamma, \dots, \gamma}_{I \text{ times}})$ 
6:   time to event  $i$ : draw sample  $\tau_i$  from  $\text{Exp}(\lambda_i)$ 
7:   choose next event  $E_\mu \in \mathcal{E}$ : for  $\mu = \arg \min_i \tau_i$ 
8:   if next event is "new infection" then
9:      $S \leftarrow S + (\tau_\mu, -1, 1, 0)$ 
10:    else if next event is "clearance" then
11:       $S \leftarrow S + (\tau_\mu, 0, -1, 1)$ 
12:    end if
13:     $\mathcal{T} \leftarrow [\mathcal{T}, S]$ 
14:  end while
15:  return  $\mathcal{T}$ 
```

Gillespie's algorithm: tau-leap

- direct method: sample time to next reaction, **only one reaction** per time step
- **tau-leap:** fix time to next reaction $\tau > 0$, sample **several reactions**
- **tau-leap assumption:** rates of reactions do not change within $[t, t + \tau)$
- tau-leap: $X(t + \tau) = X(t) + \sum_j P_j(\lambda_j \tau)$
- tau-leap: $P_j(x)$ are independent Poisson random variables with intensity x
 - 1 for each event j , sample $K_j \sim \text{Poisson}(\lambda_j \tau)$ "number of times of event"
 - 2 update: $S[t + \tau] = S[t] + \sum_j K_j v_{ij}$ for v_{ij} stoichiometric vector, state i , event j
- really fast, τ can be optimized, check assumptions!
-  09_GillespieTau.R

Simulate SIR variations with Gillespie




For the three models from the group work (incubation period, Ebola-like and two rooms in dormitory) use the R package GillespieSSA to simulate trajectories! Play around with parameters!

- **Problem:** model with **structure** for age, location, immunity, network, etc. has many different species and possible reactions ⇒ Gillespie slow: two random number draw per iteration, event/rate updates
- **Solution:** Gibson-Bruck algorithm with **data structure**
- **dependency graph** between events ⇒ event/rate update
- **indexed priority queue** of event times ⇒ single random number draw needed

Gibson-Bruck method: dependency graph

node	reaction	propensity	affects	depends	event
1	$S + I \rightarrow I + I$	$\beta S(t)I(t)^a$	I, S	I, S	new infection
2	$I \rightarrow R$	$\gamma I(t)$	I, R	I	clearance
3	$I \rightarrow \emptyset$	$\nu I(t)$	I	I	virulence
4	$\emptyset \rightarrow S$	π	S	\emptyset	birth
5	$R \rightarrow S$	$\rho R(t)$	R, S	R	immunity loss


^a $I(t)$ denotes sum of all I at time t etc.



Gibson-Bruck method: dependency graph

node	reaction	propensity	affects	depends	event
1	$S + I \rightarrow I + I$	$\beta S(t)I(t)^a$	I, S	I, S	new infection
2	$I \rightarrow R$	$\gamma I(t)$	I, R	I	clearance
3	$I \rightarrow \emptyset$	$\nu I(t)$	I	I	virulence
4	$\emptyset \rightarrow S$	π	S	\emptyset	birth
5	$R \rightarrow S$	$\rho R(t)$	R, S	R	immunity loss

^a $I(t)$ denotes sum of all I at time t etc.




dependency graph: draw edge E_{ij} iff $\text{affects}(i) \cap \text{depends}(j) \neq \emptyset$

Gibson-Bruck method: dependency graph

node	reaction	propensity	affects	depends	event
1	$S + I \rightarrow I + I$	$\beta S(t)I(t)^a$	I, S	I, S	new infection
2	$I \rightarrow R$	$\gamma I(t)$	I, R	I	clearance
3	$I \rightarrow \emptyset$	$\nu I(t)$	I	I	virulence
4	$\emptyset \rightarrow S$	π	S	\emptyset	birth
5	$R \rightarrow S$	$\rho R(t)$	R, S	R	immunity loss

^a $I(t)$ denotes sum of all I at time t etc.





dependency graph: draw edge E_{ij} iff $\text{affects}(i) \cap \text{depends}(j) \neq \emptyset$

update: reaction i happens \rightarrow propensity update for $U_i = \{j : E_{ij} \neq 0\}$

Gibson-Bruck method: indexed priority queue


- priority queue: data structure such that elements with highest priority are served first
- binary heap: complete binary tree, key stored in each node is either less than or equal to the keys in the node's children
- $\mathcal{O}(\log n)$ performance for inserts and removals, and $\mathcal{O}(n \log n)$ to build heap from n elements



Gibson-Bruck method: algorithm


-
- 1: set $t = 0$; generate dependency graph \mathcal{D} of reactions; calculate propensity function α_i for each reaction $i = 1, \dots, M$; draw $\tau_i \sim \text{Exp}(\alpha_i)$; write absolute time $t_i = t + \tau_i$ in an indexed priority queue given by heap \mathcal{Q} .
 - 2: **while** $t < t_{\max}$ **do**
 - 3: choose next reaction R_μ with μ root in \mathcal{Q}
 - 4: update stoichiometry, i.e. copy number of molecules after reaction R_μ , set $t = t_\mu$
 - 5: update reaction rates α_i for $i \in U_\mu$ using \mathcal{D}
 - 6: update next reaction times in \mathcal{Q} for updated α_i **without** new random number draw:
- $$t_{i,\text{new}} = \underbrace{\frac{\alpha_{i,\text{old}}}{\alpha_{i,\text{new}}}}_{\tau_{i,\text{new}}} (t_{i,\text{old}} - t) + t$$
- 7: **end while**
 - 8: **return** trajectory for each species and reaction times
-

Malaria toy model: Gillespie-type simulation



- α host seeking and biting rate by (female) mosquito
- γ recovery rate from human infection
- β acquisition rate from infectious host to susceptible mosquito
- μ mosquito birth rate (from both susceptible and infected mosquitoes)
- δ death rate of adult mosquito

Malaria toy model: Gillespie-type simulation



$$\begin{aligned}
 \frac{dS}{dt} &= -\alpha S \frac{V}{H} + \gamma I \\
 \frac{dI}{dt} &= \alpha S \frac{V}{H} - \gamma I \\
 \frac{dU}{dt} &= -\beta U \frac{I}{H} + \mu(U + V) - \delta U \\
 \frac{dV}{dt} &= \beta U \frac{I}{H} - \delta V
 \end{aligned}$$

Where does the term $\frac{V}{H}$ come from?

- α host seeking and biting rate
- then, $\alpha(U + V)$ is expected number of bites
- $\frac{\alpha(U+V)}{H}$ are expected number of bites per human
- multiply with infectious mosquito density $\frac{V}{U+V}$ gives
- $\frac{V}{U+V} \frac{\alpha(U+V)}{H} = \frac{\alpha V}{H}$

$$\begin{aligned}\frac{dS}{dt} &= -\alpha S \frac{V}{H} + \gamma I \\ \frac{dI}{dt} &= \alpha S \frac{V}{H} - \gamma I \\ \frac{dU}{dt} &= -\beta U \frac{I}{H} + \mu(U + V) - \delta U \\ \frac{dV}{dt} &= \beta U \frac{I}{H} - \delta V\end{aligned}$$

Malaria toy model: Gillespie-type simulation

$$\frac{dS}{dt} = -\alpha V \frac{S}{H} + \gamma I$$

$$\frac{dI}{dt} = \alpha V \frac{S}{H} - \gamma I$$

$$\frac{dU}{dt} = -\beta U \frac{I}{H} + \mu(U + V) - \delta U$$

$$\frac{dV}{dt} = \beta U \frac{I}{H} - \delta V$$

- 4 species: S, I, U, V
- 7 reactions: $S + V \rightarrow I + V$, $I \rightarrow S$, $U + I \rightarrow V + I$, $U \rightarrow \emptyset$, $V \rightarrow \emptyset$, $U \rightarrow 2U$, $V \rightarrow U + V$
- 7 intensities: $\alpha \frac{S}{H}$, γI , $\beta \frac{U}{H}$, δU , δV , μU , μV ,
- stoichiometry 4×7 matrix:

$$\begin{bmatrix} -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 & 1 & 1 \\ 0 & 0 & +1 & 0 & -1 & 0 & 0 \end{bmatrix}$$

For this model, the basic reproduction number is

$$\mathcal{R}_0 = \sqrt{\frac{\alpha\beta(U + V)/H}{\mu\gamma}}$$

Malaria toy model



Simulate with GillespieSSA and obtain an endemic equilibrium!


Choose $H = 1000$, $U + V = 5000$, $\mu = \delta = 1/10$, $\beta = \alpha = 0.03$ and search some values for γ in literature! Change the ratio of mosquito $M = U + V$ to human $H = S + I$!

Plot the curve of infected humans over two year!

 10_RossMcDonaldGillespie.R

Malaria toy model: Gillespie-type simulation

$$\begin{aligned}\frac{dS}{dt} &= -\alpha V \frac{S}{H} + \gamma I \\ \frac{dI}{dt} &= \alpha V \frac{S}{H} - \gamma I \\ \frac{dU}{dt} &= -\beta U \frac{I}{H} + \mu(U + V) - \delta U \\ \frac{dV}{dt} &= \beta U \frac{I}{H} - \delta V\end{aligned}$$




Asymptomatic infections



Based on our Malaria toy model, we consider three classes of infected populations: confirmed cases I_c who are treated before gametocytemia, asymptomatic with high gametocytemia I_h and asymptomatic with low gametocytemia I_l . We assume that hosts with high/low gametocytemia have a transmission rate β_h, β_l . The duration of infection with positive gametocytemia for I_c, I_h, I_l is 10, 45, 15 days resp.

Asymptomatic Malaria toy model: Gillespie simulation



Andolina et al. 2021: The bar heights indicate the proportion of mosquitoes that became infected when feeding on this population. The bar widths indicate the proportion of the infected population.

Asymptomatic infections



Use the figure to discuss parameters for proportions of I_c , I_h , I_e and the ratio of β_h over β_e . Draw the flow diagram, use parameters from the toy model, write the reactions, rates and stoichiometric vectors. Simulate the dynamics of infection compartments with the Gillespie algorithm!


Asymptomatic Malaria toy model: Gillespie simulation

Proposed solution:

- New infections: Exposure to infectious mosquitoes V creates new infections in I_c , I_h , I_ℓ at rate $\alpha_c = p_c\alpha$, $\alpha_h = p_h\alpha$ and $\alpha_\ell = p_\ell\alpha$ where EIR $\alpha = 2$ and $[p_c, p_h, p_\ell] = [0.05, 0.17, 0.78]$ the relative proportion of confirmed/treated, high and low parasitemia infections after a mosquito bite.
- clearance rates for human infections are $\gamma_c = 1/10$, $\gamma_h = 1/45$, $\gamma_\ell = 1/15$.
- For transmission from humans to vectors, we assume that $\beta_c = 0.03$ and $\beta_h = 0.08$ and $\beta_\ell = K\beta_h$ for $K = 0.84/0.16$, i.e. the transmission ratio into I_h vs I_c infections.
- The life-cycle for the mosquitoes populations remains as in the toy-model before.
-  `11_RossMcDonaldGillespieAsymptomatic.R`


Asymptomatic Malaria toy model: biochemical reaction system

- ① $S + V \rightarrow I_c + V$ at rate $\alpha p_c \frac{S}{H}$
- ② $S + V \rightarrow I_h + V$ at rate $\alpha p_h \frac{S}{H}$
- ③ $S + V \rightarrow I_\ell + V$ at rate $\alpha p_\ell \frac{S}{H}$
- ④ $I_c \rightarrow \emptyset$ at rate γ_c
- ⑤ $I_h \rightarrow \emptyset$ at rate γ_h
- ⑥ $I_\ell \rightarrow \emptyset$ at rate γ_ℓ
- ⑦ $U + I_c \rightarrow V + I_c$ at rate $\beta_c \frac{U}{H}$
- ⑧ $U + I_h \rightarrow V + I_h$ at rate $\beta_h \frac{U}{H}$
- ⑨ $U + I_\ell \rightarrow V + I_\ell$ at rate $\beta_\ell \frac{U}{H}$
- ⑩ $U \rightarrow \emptyset$ at rate δ
- ⑪ $V \rightarrow \emptyset$ at rate δ
- ⑫ $U \rightarrow U + U$ at rate μ
- ⑬ $V \rightarrow U + V$ at rate μ



Asymptomatic Malaria toy model: Gillespie simulation

With initial conditions $H = 10000$, $I_c(t) = 20$, $I_h(0) = 5$, $I_\ell(0) = 10$ and $V(0) = 8$ and $U(0) = 49992$, we obtain an endemic equilibrium of confirmed cases prevalence at roughly 3%, while a large part of the population is infected without symptoms at low-level parasitemia:



Test and treat vs. mass drug administration



For the asymptomatic model, we want to evaluate two different intervention strategies:

- test and treat: the antigen-based diagnostics has a sensitivity to detect 95% of asymptomatic cases with high gametocytemia and 15% with low gametocytemia, all positively tested are treated.
- mass drug administration: 95% of the entire population gets drug treatment, regardless of infection status

Simulate trajectories for the two strategies and the counterfactual starting from the endemic equilibrium obtained from the preceding exercise!

What is your metric of evaluation and which intervention would you recommend?


Asymptomatic Malaria intervention toy model: reaction system

For test and treat we assume that both I_h and I_ℓ move to treated compartment T . For mass drug administration, we also assume that S move into T .

- ⑭ $I_h \rightarrow T$ at rate T_h
- ⑮ $I_\ell \rightarrow T$ at rate T_ℓ
- ⑯ $S \rightarrow T$ at rate M
- ⑰ $T \rightarrow S$ at rate $r = 1/30$


R

12_RossMcDonaldGillespieAsymptomatic_TTvsmDA.R



Asymptomatic Malaria intervention toy model: Gillespie simulation

- test and treat: $T_h = 0.95$, $T_\ell = 0.15$ and $M = 0$, $r = 1/30$
- MDA: $T_h = T_\ell = M = 0.98$, $r = 1/30$



Prevalence is close to 0 within 3 months, test sensitivity for I_ℓ is crucial to achieve elimination.

Stochastic differential equation heuristics

- consider SIR stochastic process $X(t) = [S(t), I(t)]$ s.th. for $\bar{X}(t) = \mathbb{E}(X(t))$

$$\frac{d\bar{S}}{dt} = -\beta \bar{S} \frac{\bar{I}}{\bar{N}}$$
$$\frac{d\bar{I}}{dt} = \beta \bar{S} \frac{\bar{I}}{\bar{N}} - \gamma \bar{I}$$

Stochastic differential equation heuristics

- consider SIR stochastic process $X(t) = [S(t), I(t)]$ s.th. for $\bar{X}(t) = \mathbb{E}(X(t))$

$$\frac{d\bar{S}}{dt} = -\beta \bar{S} \frac{\bar{I}}{\bar{N}} \quad \frac{d\bar{I}}{dt} = \beta \bar{S} \frac{\bar{I}}{\bar{N}} - \gamma \bar{I}$$

- divide time interval $[0, t]$ into subintervals of length Δt , with

$$\Delta X(t) = [\Delta S(t), \Delta I(t)] = [S(t + \Delta t) - S(t), I(t + \Delta t) - I(t)]$$

Stochastic differential equation heuristics

- consider SIR stochastic process $X(t) = [S(t), I(t)]$ s.th. for $\bar{X}(t) = \mathbb{E}(X(t))$

$$\frac{d\bar{S}}{dt} = -\beta \bar{S} \frac{\bar{I}}{\bar{N}} \quad \frac{d\bar{I}}{dt} = \beta \bar{S} \frac{\bar{I}}{\bar{N}} - \gamma \bar{I}$$

- divide time interval $[0, t]$ into subintervals of length Δt , with

$$\Delta X(t) = [\Delta S(t), \Delta I(t)] = [S(t + \Delta t) - S(t), I(t + \Delta t) - I(t)]$$

- further divide Δt s.th. for $\Delta t_i = t_i - t_{i-1}$: $\sum_i^n \Delta t_i = \Delta t$ and

$$\Delta X(t) = \sum_i \Delta X(t_i)$$

Stochastic differential equation heuristics

- consider SIR stochastic process $X(t) = [S(t), I(t)]$ s.th. for $\bar{X}(t) = \mathbb{E}(X(t))$

$$\frac{d\bar{S}}{dt} = -\beta \bar{S} \frac{\bar{I}}{\bar{N}} \quad \frac{d\bar{I}}{dt} = \beta \bar{S} \frac{\bar{I}}{\bar{N}} - \gamma \bar{I}$$

- divide time interval $[0, t]$ into subintervals of length Δt , with

$$\Delta X(t) = [\Delta S(t), \Delta I(t)] = [S(t + \Delta t) - S(t), I(t + \Delta t) - I(t)]$$

- further divide Δt s.th. for $\Delta t_i = t_i - t_{i-1}$: $\sum_i^n \Delta t_i = \Delta t$ and

$$\Delta X(t) = \sum_i \Delta X(t_i)$$

- if Δt_i small, assume $\Delta X(t_i)$ are iid on Δt

Stochastic differential equation heuristics

- consider SIR stochastic process $X(t) = [S(t), I(t)]$ s.th. for $\bar{X}(t) = \mathbb{E}(X(t))$

$$\frac{d\bar{S}}{dt} = -\beta \bar{S} \frac{\bar{I}}{\bar{N}} \quad \frac{d\bar{I}}{dt} = \beta \bar{S} \frac{\bar{I}}{\bar{N}} - \gamma \bar{I}$$

- divide time interval $[0, t]$ into subintervals of length Δt , with

$$\Delta X(t) = [\Delta S(t), \Delta I(t)] = [S(t + \Delta t) - S(t), I(t + \Delta t) - I(t)]$$

- further divide Δt s.th. for $\Delta t_i = t_i - t_{i-1}$: $\sum_i^n \Delta t_i = \Delta t$ and

$$\Delta X(t) = \sum_i \Delta X(t_i)$$

- if Δt_i small, assume $\Delta X(t_i)$ are iid on Δt
- for n large, CTL: $\frac{1}{\sqrt{n}} (\Delta X(t) - \mathbb{E}(\Delta X(t))) \sim \mathcal{N}(0, \text{cov}(\Delta X(t)))$

At the order of Δt :

$$\mathbb{E}(\Delta X(t)) \approx \left[-\beta \bar{S} \frac{\bar{I}}{N}, \beta \bar{S} \frac{\bar{I}}{N} - \gamma \bar{I} \right] \Delta t = f \Delta t$$

At the order of Δt :

$$\mathbb{E}(\Delta X(t)) \approx \left[-\beta \bar{S} \frac{\bar{I}}{N}, \beta \bar{S} \frac{\bar{I}}{N} - \gamma \bar{I} \right] \Delta t = f \Delta t$$

At the order of Δt :

$$\text{cov}(\Delta X) \approx \mathbb{E}((\Delta X)(\Delta X)^T) = \begin{pmatrix} \text{cov}(\Delta S, \Delta S) & \text{cov}(\Delta S, \Delta I) \\ \text{cov}(\Delta S, \Delta I) & \text{cov}(\Delta I, \Delta I) \end{pmatrix}$$

At the order of Δt :

$$\mathbb{E}(\Delta X(t)) \approx \left[-\beta \bar{S} \frac{\bar{I}}{N}, \beta \bar{S} \frac{\bar{I}}{N} - \gamma \bar{I} \right] \Delta t = f \Delta t$$

At the order of Δt :

$$\text{cov}(\Delta X) \approx \mathbb{E}((\Delta X)(\Delta X)^T) = \begin{pmatrix} \text{cov}(\Delta S, \Delta S) & \text{cov}(\Delta S, \Delta I) \\ \text{cov}(\Delta S, \Delta I) & \text{cov}(\Delta I, \Delta I) \end{pmatrix}$$

$$\text{cov}(\Delta X) \approx \begin{pmatrix} \beta S \frac{I}{N} & -\beta S \frac{I}{N} \\ -\beta S \frac{I}{N} & \beta S \frac{I}{N} + \gamma I \end{pmatrix} \Delta t = C \Delta t$$

By assumption $\Delta X(t_i)$ are iid on Δt , and with $\Delta t = n\Delta t_i$ s.th.

$$\begin{aligned}\mathbb{E}(\Delta X_1^2) &= \mathbb{E}(\Delta S^2) = \mathbb{E}\left(\sum_i \Delta S(t_i)^2\right) \\ &= \sum_i \mathbb{E}(\Delta S(t_i)^2) + 2 \sum_{i < j} \mathbb{E}(\Delta S(t_i))\mathbb{E}(\Delta S(t_j)) \\ &= n\mathbb{E}(\Delta S(t_0)^2) + n(n-1)\mathbb{E}(\Delta S(t_0))^2 \\ &= n(-1)^2 \Delta t_1 \beta I(t_0) \frac{S(t_0)}{N} + n 0^2 (1 - \Delta t_1 \beta I(t_1) \frac{S(t_1)}{N}) + (\Delta t)^2 (1 - \frac{1}{n})(\beta I(t_1) \frac{S(t_1)}{N})^2 \\ &\approx \Delta t \beta I(t_0) \frac{S(t_0)}{N}\end{aligned}$$

at the order of Δt with $\mathbb{P}(\Delta S(t_i) = -1) = \Delta t_i \beta I(t_{i-1}) \frac{S(t_{i-1})}{N}$

Stochastic differential equation



$$\Delta X(t) \approx f(X(t))\Delta t + G(X(t))\Delta W(t)$$

Stochastic differential equation



$$\Delta X(t) \approx f(X(t))\Delta t + G(X(t))\Delta W(t)$$

- Here, the matrix G is such that $GG^T = C$ and

Stochastic differential equation



$$\Delta X(t) \approx f(X(t))\Delta t + G(X(t))\Delta W(t)$$

- Here, the matrix G is such that $GG^T = C$ and
- $\Delta W = [\Delta W_1, \Delta W_2]$ with $\Delta W_i \sim \mathcal{N}(0, \Delta t)$

Euler-Maruyama algorithm



We implement

$$\Delta X(t) \approx f(X(t))\Delta t + G(X(t))\Delta W(t)$$

by first order scheme

$$X[i+1] = X[i] + f(X[i])\Delta t + G(X[i])\eta\sqrt{\Delta t}$$

where $\eta \in \mathbb{R}^d$ with $\eta_k \sim \mathcal{N}(0, 1)$ and d is the number of reactions.

Euler-Maruyama algorithm for SIR

In our SIR example:

$$f(S, I) = \begin{pmatrix} -\beta S \frac{I}{N} \\ \beta S \frac{I}{N} - \gamma I \end{pmatrix}$$
$$G(S, I) = \begin{pmatrix} -\sqrt{\beta S \frac{I}{N}} & 0 \\ \sqrt{\beta S \frac{I}{N}} & -\sqrt{\gamma I} \end{pmatrix}$$


Just take square roots of the rates from the ODE!

Implementation of Euler-Maruyama




Code the Euler-Maruyama scheme in R for the influenza boarding school SIR model! Simulate several trajectories! When choosing $\beta > \gamma$, do you have simulations where I get extinct early on?  13_ForwardEulerMaruyamaSIR.R

Euler-Maruyama algorithm for SIR



Malaria toy model with stochastic differential equation



$$\begin{aligned} dS &= f_1 dt - \sqrt{\alpha V \frac{S}{H}} dW^1 + \sqrt{\gamma I} dW^2 \\ dI &= f_2 dt + \sqrt{\alpha V \frac{S}{H}} dW^1 - \sqrt{\gamma I} dW^2 \\ dU &= f_3 dt - \sqrt{\beta U \frac{I}{H}} dW^3 + \sqrt{\mu U} dW^4 + \sqrt{\mu V} dW^5 - \sqrt{\delta U} dW^6 \\ dV &= f_4 dt + \sqrt{\beta U \frac{I}{H}} dW^3 - \sqrt{\delta V} dW^7 \end{aligned}$$


Euler-Maruyama for Malaria toy model



Implement the stochastic differential equation version of the Malaria toy model in R.


R 14_RossMcDonaldForwardEulerMaruyama.R

Euler-Maruyama for Malaria toy model



Sellke's method: infectious pressure

- **probabilistically equivalent** to SIR process
- iid $Q_1, \dots, Q_n \sim \text{Exp}(1)$ for n susceptibles
- iid $T_{-(m-1)}, \dots, T_n$ infection durations, **any distribution** on \mathbb{R}_+ (e.g. gamma, Weibull)
- m initially infected $T_{-(m-1)}, \dots, T_0$
- $I(t) =$ number of infected at time t
- infectious "pressure" $\Lambda(t) = \int_0^t I(s)ds$



Sellke's method: infectious pressure

- susceptible i accumulates exposure to infection at rate equal to number of infected individuals
- i th susceptible becomes infected by time t_i if infectious pressure reached: $\Lambda(t_i) = Q_i$
- individual who was j th infected remains infected for time T_j and then clears
- infections happen at the right time:

$$\begin{aligned}\mathbb{P}(\text{susceptible } i \text{ infected by } t + dt | \text{not infected by } t) &= \\ &= \mathbb{P}(Q_i < \Lambda(t + dt) | Q_i > \Lambda(t)) = \frac{\mathbb{P}(\Lambda(t) < Q_i < \Lambda(t + dt))}{\mathbb{P}(Q_i > \Lambda(t))} \\ &\approx \frac{(1 - e^{-\Lambda(t+dt)}) - (1 - e^{-\Lambda(t)})}{e^{-\Lambda(t)}} = 1 - e^{-[\Lambda(t+dt) - \Lambda(t)]} = 1 - e^{-\Lambda'(t)dt} \\ &= \Lambda'(t)dt = \beta I(t)dt\end{aligned}$$

- **advantage:** generalize straight-forward to infection duration **with memory**

Stochastic simulation algorithms: summary table

algorithm	time	space	conv	non-Mark.	in practice
exact Gillespie	C	D	✓	✗	only for simple systems, slow
first reaction Gillespie	C	D	✗	✗	no need to sample next reaction
tau-leap Gillespie	D	D	✗	✗	fast for simple systems, step size tuning
Gillespie-Boguña	C	D	✗	✓	only for simple systems, slow
Gillespie-Gibson-Bruck	C	D	✗	✓	fast for system with many reactions
Sellke	C	D	✓	✓	only for simple systems
Euler-Maruyama	C	C	✓	✗	faster to simulate for large populations

Thanks for financial support to

**BILL & MELINDA
GATES foundation**



Swiss Tropical and Public Health Institute

internet mathoverflow, stackoverflow, chatGPT, google colab

lecture notes Anderson, Kurtz: Stochastic Analysis of Biochemical Systems 

lecture notes Allen: Stochastic Population and Epidemic Models 

lecture notes Ammari, Wu, Yu: Numerical Methods for ODEs 

history Ross: An Application of the Theory of Probabilities to the Study of a priori
pathometry. -Part I 

- article** Lehtinen et al. 
- article** Begon et al. 