

# Bairstow's Method

MATH 4701 Numerical Analysis

# Synthetic Division

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x - a$  we follow these steps,

# Synthetic Division

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x - a$  we follow these steps,

- 1 We form a table with **three** rows and  $n + 1$  columns.

# Synthetic Division

To divide a polynomial  $P(x) = x^n + a_nx^{n-1} + \dots + a_1x + a_0$  by a factor of  $x - a$  we follow these steps,

- ① We form a table with **three** rows and  $n + 1$  columns.
- ② We write the coefficients of  $P(x)$  sorted by degree in the first row.

# Synthetic Division

To divide a polynomial  $P(x) = x^n + a_nx^{n-1} + \dots + a_1x + a_0$  by a factor of  $x - a$  we follow these steps,

- ① We form a table with **three** rows and  $n + 1$  columns.
- ② We write the coefficients of  $P(x)$  sorted by degree in the first row.
- ③ We start by writing 0 in the second row of the first column.

# Synthetic Division

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x - a$  we follow these steps,

- ① We form a table with **three** rows and  $n + 1$  columns.
- ② We write the coefficients of  $P(x)$  sorted by degree in the first row.
- ③ We start by writing 0 in the second row of the first column.
- ④ In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

# Synthetic Division

To divide a polynomial  $P(x) = x^n + a_nx^{n-1} + \dots + a_1x + a_0$  by a factor of  $x - a$  we follow these steps,

- ① We form a table with **three** rows and  $n + 1$  columns.
- ② We write the coefficients of  $P(x)$  sorted by degree in the first row.
- ③ We start by writing 0 in the second row of the first column.
- ④ In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.
- ⑤ Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

# Synthetic Division

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x - a$  we follow these steps,

- 1 We form a table with **three** rows and  $n + 1$  columns.
- 2 We write the coefficients of  $P(x)$  sorted by degree in the first row.
- 3 We start by writing 0 in the second row of the first column.
- 4 In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.
- 5 Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.
- 6 When we fill the entry in the third row of the last column, this entry is the remainder of division of  $P(x)$  by  $x - a$ .



# Synthetic Division

To divide a polynomial  $P(x) = x^n + a_nx^{n-1} + \dots + a_1x + a_0$  by a factor of  $x - a$  we follow these steps,

- 1 We form a table with **three** rows and  $n + 1$  columns.
- 2 We write the coefficients of  $P(x)$  sorted by degree in the first row.
- 3 We start by writing 0 in the second row of the first column.
- 4 In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.
- 5 Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.
- 6 When we fill the entry in the third row of the last column, this entry is the remainder of division of  $P(x)$  by  $x - a$ .
- 7 The rest of the numbers in the third row in the order they appear are the coefficients of the quotient of division of  $P(x)$  by  $x - a$ .

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

We form a table with three rows and  $n + 1 = 6$  columns.

$$a = 2$$


# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

We write the coefficients of  $P(x)$  sorted by degree in the first row.

$$a = 2$$

1	-4	3	2	-5	-3

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

We start by writing 0 in the second row of the first column.

$$a = 2$$

1	-4	3	2	-5	-3
0					

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0					

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0					
1					

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2				
1					



# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2				
1	-2				

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4			
1	-2				

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4			
1	-2	-1			

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2		
1	-2	-1			

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2		
1	-2	-1	0		

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2	0	
1	-2	-1	0		

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2	0	
1	-2	-1	0	-5	

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2	0	-10
1	-2	-1	0	-5	



# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

In each step we calculate the sum of the entries in the first and second rows of the current column and write the sum in the third row under the same column.

Then we multiply the number in the third row we just found by  $a$  and we write the product in the second row of the next column.

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2	0	-10
1	-2	-1	0	-5	-13

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

When we fill the entry in the third row of the last column, this entry is the remainder of division of  $P(x)$  by  $x - a$ .

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2	0	-10
1	-2	-1	0	-5	-13

Remainder =  $-13$

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

The rest of the numbers in the third row in the order they appear are the coefficients of the quotient of division of  $P(x)$  by  $x - a$ .

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2	0	-10
1	-2	-1	0	-5	-13

$$\text{Remainder} = -13$$

$$\text{Quotient} = x^4 - 2x^3 - x^2 - 5$$

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

The rest of the numbers in the third row in the order they appear are the coefficients of the quotient of division of  $P(x)$  by  $x - a$ .

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2	0	-10
1	-2	-1	0	-5	-13

$$\text{Remainder} = -13$$

$$\text{Quotient} = x^4 - 2x^3 - x^2 - 5$$

Notice that since  $P(x) = Q(x)(x - a) + R$ ,  $P(a) = Q(a)(a - a) + R = R$ .

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

The rest of the numbers in the third row in the order they appear are the coefficients of the quotient of division of  $P(x)$  by  $x - a$ .

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2	0	-10
1	-2	-1	0	-5	-13

$$\text{Remainder} = -13$$

$$\text{Quotient} = x^4 - 2x^3 - x^2 - 5$$

Notice that since  $P(x) = Q(x)(x - a) + R$ ,  $P(a) = Q(a)(a - a) + R = R$ . Therefore, the remainder of division of  $P(x)$  by  $x - a$  calculates  $P(a)$ .

# Synthetic Division

**Example:** Find the quotient and remainder of division of  $P(x) = x^5 - 4x^4 + 3x^3 + 2x^2 - 5x - 3$  by  $x - 2$ .

The rest of the numbers in the third row in the order they appear are the coefficients of the quotient of division of  $P(x)$  by  $x - a$ .

$$a = 2$$

1	-4	3	2	-5	-3
0	2	-4	-2	0	-10
1	-2	-1	0	-5	-13

$$\text{Remainder} = -13$$

$$\text{Quotient} = x^4 - 2x^3 - x^2 - 5$$

Notice that since  $P(x) = Q(x)(x - a) + R$ ,  $P(a) = Q(a)(a - a) + R = R$ . Therefore, the remainder of division of  $P(x)$  by  $x - a$  calculates  $P(a)$ . In this example  $P(2) = -13$ .

# Synthetic Division by a Quadratic Polynomial

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x^2 - px - q$  we follow these steps,

# Synthetic Division by a Quadratic Polynomial

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x^2 - px - q$  we follow these steps,

- 1 We form a table with **four** rows and  $n + 1$  columns.



# Synthetic Division by a Quadratic Polynomial

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x^2 - px - q$  we follow these steps,

- 1 We form a table with **four** rows and  $n + 1$  columns.
- 2 We write the coefficients of  $P(x)$  sorted by degree in the first row.

# Synthetic Division by a Quadratic Polynomial

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x^2 - px - q$  we follow these steps,

- ① We form a table with **four** rows and  $n + 1$  columns.
- ② We write the coefficients of  $P(x)$  sorted by degree in the first row.
- ③ We start by writing 0 in the second and third rows of the first column, the second row of the second column, and the third row of the last column.

# Synthetic Division by a Quadratic Polynomial

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x^2 - px - q$  we follow these steps,

- ① We form a table with **four** rows and  $n + 1$  columns.
- ② We write the coefficients of  $P(x)$  sorted by degree in the first row.
- ③ We start by writing 0 in the second and third rows of the first column, the second row of the second column, and the third row of the last column.
- ④ In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

# Synthetic Division by a Quadratic Polynomial

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x^2 - px - q$  we follow these steps,

- ① We form a table with **four** rows and  $n + 1$  columns.
- ② We write the coefficients of  $P(x)$  sorted by degree in the first row.
- ③ We start by writing 0 in the second and third rows of the first column, the second row of the second column, and the third row of the last column.
- ④ In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.
- ⑤ Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

# Synthetic Division by a Quadratic Polynomial

To divide a polynomial  $P(x) = x^n + a_n x^{n-1} + \dots + a_1 x + a_0$  by a factor of  $x^2 - px - q$  we follow these steps,

- 1 We form a table with **four** rows and  $n + 1$  columns.
- 2 We write the coefficients of  $P(x)$  sorted by degree in the first row.
- 3 We start by writing 0 in the second and third rows of the first column, the second row of the second column, and the third row of the last column.
- 4 In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.
- 5 Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.
- 6 We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

# Synthetic Division by a Quadratic Polynomial

- 1 We form a table with **four** rows and  $n + 1$  columns.
- 2 We write the coefficients of  $P(x)$  sorted by degree in the first row.
- 3 We start by writing 0 in the second and third rows of the first column, the second row of the second column, and the third row of the last column.
- 4 In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.
- 5 Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.
- 6 We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.
- 7 When we fill the entry in the fourth row of the last column, the last two entries  $a$  and  $b$  in the fourth row determine the remainder of division of  $P(x)$  by  $x^2 - px - q$  as  $ax + b$ .

# Synthetic Division by a Quadratic Polynomial

- ① We form a table with **four** rows and  $n + 1$  columns.
- ② We write the coefficients of  $P(x)$  sorted by degree in the first row.
- ③
- ④
- ⑤ Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.
- ⑥ We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.
- ⑦ When we fill the entry in the fourth row of the last column, the last two entries  $a$  and  $b$  in the fourth row determine the remainder of division of  $P(x)$  by  $x^2 - px - q$  as  $ax + b$ .
- ⑧ The rest of the numbers in the fourth row in the order they appear are the coefficients of the quotient of division of  $P(x)$  by  $x^2 - px - q$ .

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .



# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

We form a table with **four** rows and  $n + 1 = 5$  columns.

$$p = 2 \quad q = -5$$


# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

We write the coefficients of  $P(x)$  sorted by degree in the first row.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

We start by writing 0 in the second and third rows of the first column, the second row of the second column, and the third row of the last column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0			
0				0

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0			
0				0

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0			
0				0
1				

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0			
0	2			0
1				

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0			
0	2			0
1	0			

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0			
0	2	0		0
1	0			



# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0	-5		
0	2	0		0
1	0			

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0	-5		
0	2	0		0
1	0	-1		

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0	-5		
0	2	0	-2	0
1	0	-1		

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0	-5	0	
0	2	0	-2	0
1	0	-1		

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0	-5	0	
0	2	0	-2	0
1	0	-1	-3	

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0	-5	0	5
0	2	0	-2	0
1	0	-1	-3	

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

In each step we calculate the sum of the entries in the first three rows of the current column and write the sum in the fourth row under the same column.

Then we multiply the number in the fourth row of the current column by  $p$  and we write the product in the third row of the next column.

We multiply the number in the fourth row of the previous column by  $q$  and we write the product in the second row of the next column.

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0	-5	0	5
0	2	0	-2	0
1	0	-1	-3	12

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

When we fill the entry in the fourth row of the last column, the last two entries  $a$  and  $b$  in the fourth row determine the remainder of division of  $P(x)$  by  $x^2 - px - q$  as  $ax + b$ .

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0	-5	0	5
0	2	0	-2	0
1	0	-1	-3	12

Remainder =  $-3x + 12$



# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

The rest of the numbers in the fourth row in the order they appear are the coefficients of the quotient of division of  $P(x)$  by  $x^2 - px - q$ .

$$p = 2 \quad q = -5$$

1	-2	4	-1	7
0	0	-5	0	5
0	2	0	-2	0
1	0	-1	-3	12

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

For every complex number  $\omega = a + bi$ , its conjugate  $\bar{\omega} = a - bi$  satisfies  $\omega\bar{\omega} = |\omega|^2 = a^2 + b^2$  and  $\omega + \bar{\omega} = 2a$  which are both real numbers.

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

For every complex number  $\omega = a + b\mathbf{i}$ , its conjugate  $\bar{\omega} = a - b\mathbf{i}$  satisfies  $\omega\bar{\omega} = |\omega|^2 = a^2 + b^2$  and  $\omega + \bar{\omega} = 2a$  which are both real numbers.

Therefore,  $q(x) = (x - \omega)(x - \bar{\omega}) = x^2 - 2ax + a^2 + b^2$  is a quadratic polynomial with real coefficients.

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

For every complex number  $\omega = a + b\mathbf{i}$ , its conjugate  $\bar{\omega} = a - b\mathbf{i}$  satisfies  $\omega\bar{\omega} = |\omega|^2 = a^2 + b^2$  and  $\omega + \bar{\omega} = 2a$  which are both real numbers.

Therefore,  $q(x) = (x - \omega)(x - \bar{\omega}) = x^2 - 2ax + a^2 + b^2$  is a quadratic polynomial with real coefficients.

Now if  $Q(x)$  and  $R(x)$  are quotient and remainder of division of  $P(x)$  by  $q(x)$ , then  $P(\omega) = Q(\omega)q(\omega) + R(\omega)$

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

For every complex number  $\omega = a + bi$ , its conjugate  $\bar{\omega} = a - bi$  satisfies  $\omega\bar{\omega} = |\omega|^2 = a^2 + b^2$  and  $\omega + \bar{\omega} = 2a$  which are both real numbers.

Therefore,  $q(x) = (x - \omega)(x - \bar{\omega}) = x^2 - 2ax + a^2 + b^2$  is a quadratic polynomial with real coefficients.

Now if  $Q(x)$  and  $R(x)$  are quotient and remainder of division of  $P(x)$  by  $q(x)$ , then  $P(\omega) = Q(\omega)q(\omega) + R(\omega) = R(\omega)$  because  $q(\omega) = (\omega - \omega)(\omega - \bar{\omega}) = 0$ .

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

Now if  $Q(x)$  and  $R(x)$  are quotient and remainder of division of  $P(x)$  by  $q(x)$ , then  $P(\omega) = Q(\omega)q(\omega) + R(\omega) = R(\omega)$  because  $q(\omega) = (\omega - \omega)(\omega - \bar{\omega}) = 0$ .

In the above example

$$x^2 - 2x + 5 = 0 \Leftrightarrow x^2 - 2x + 1 + 4 = 0$$

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

Now if  $Q(x)$  and  $R(x)$  are quotient and remainder of division of  $P(x)$  by  $q(x)$ , then  $P(\omega) = Q(\omega)q(\omega) + R(\omega) = R(\omega)$  because  $q(\omega) = (\omega - \omega)(\omega - \bar{\omega}) = 0$ .

In the above example

$$x^2 - 2x + 5 = 0 \Leftrightarrow x^2 - 2x + 1 + 4 = 0 \Leftrightarrow x^2 - 2x + 1 = -4$$



# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

Now if  $Q(x)$  and  $R(x)$  are quotient and remainder of division of  $P(x)$  by  $q(x)$ , then  $P(\omega) = Q(\omega)q(\omega) + R(\omega) = R(\omega)$  because  $q(\omega) = (\omega - \omega)(\omega - \bar{\omega}) = 0$ .

In the above example

$$\begin{aligned} x^2 - 2x + 5 = 0 &\Leftrightarrow x^2 - 2x + 1 + 4 = 0 \Leftrightarrow x^2 - 2x + 1 = -4 \\ &\Leftrightarrow (x - 1)^2 = -4 \end{aligned}$$

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

Now if  $Q(x)$  and  $R(x)$  are quotient and remainder of division of  $P(x)$  by  $q(x)$ , then  $P(\omega) = Q(\omega)q(\omega) + R(\omega) = R(\omega)$  because  $q(\omega) = (\omega - \omega)(\omega - \bar{\omega}) = 0$ .

In the above example

$$\begin{aligned} x^2 - 2x + 5 = 0 &\Leftrightarrow x^2 - 2x + 1 + 4 = 0 \Leftrightarrow x^2 - 2x + 1 = -4 \\ &\Leftrightarrow (x - 1)^2 = -4 \Leftrightarrow x - 1 = \pm 2i \end{aligned}$$

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Synthetic division by quadratic polynomials can be used to evaluate a polynomial at a complex number.

Now if  $Q(x)$  and  $R(x)$  are quotient and remainder of division of  $P(x)$  by  $q(x)$ , then  $P(\omega) = Q(\omega)q(\omega) + R(\omega) = R(\omega)$  because  $q(\omega) = (\omega - \omega)(\omega - \bar{\omega}) = 0$ .

In the above example

$$\begin{aligned} x^2 - 2x + 5 = 0 &\Leftrightarrow x^2 - 2x + 1 + 4 = 0 \Leftrightarrow x^2 - 2x + 1 = -4 \\ &\Leftrightarrow (x - 1)^2 = -4 \Leftrightarrow x - 1 = \pm 2i \Leftrightarrow x = 1 \pm 2i \end{aligned}$$

# Synthetic Division by a Quadratic Polynomial

**Example:** Find the quotient and remainder of division of  $P(x) = x^4 - 2x^3 + 4x^2 - x + 7$  by  $x^2 - 2x + 5$ .

$$\text{Remainder} = -3x + 12$$

$$\text{Quotient} = x^2 - 1$$

Now if  $Q(x)$  and  $R(x)$  are quotient and remainder of division of  $P(x)$  by  $q(x)$ , then  $P(\omega) = Q(\omega)q(\omega) + R(\omega) = R(\omega)$  because  $q(\omega) = (\omega - \omega)(\omega - \bar{\omega}) = 0$ .

In the above example

$$\begin{aligned} x^2 - 2x + 5 = 0 &\Leftrightarrow x^2 - 2x + 1 + 4 = 0 \Leftrightarrow x^2 - 2x + 1 = -4 \\ &\Leftrightarrow (x - 1)^2 = -4 \Leftrightarrow x - 1 = \pm 2i \Leftrightarrow x = 1 \pm 2i \end{aligned}$$

Therefore, to calculate  $P(1 + 2i)$  we can evaluate

$$R(1 + 2i) = -3(1 + 2i) + 12 = 9 - 6i.$$

# Recursive Relations for Division by a Quadratic Polynomial

When we divide polynomial  $P(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$  by a quadratic polynomial  $x^2 - px - q$  to get the quotient  $Q(x) = x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0$  and remainder  $R(x) = ux + v$  we have

$$P(x) = (x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0)(x^2 - px - q) + ux + v$$

# Recursive Relations for Division by a Quadratic Polynomial

When we divide polynomial  $P(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$  by a quadratic polynomial  $x^2 - px - q$  to get the quotient

$Q(x) = x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0$  and remainder  $R(x) = ux + v$  we have

$$P(x) = (x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0)(x^2 - px - q) + ux + v$$

Matching the coefficients on both sides we get the following relations

# Recursive Relations for Division by a Quadratic Polynomial

When we divide polynomial  $P(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$  by a quadratic polynomial  $x^2 - px - q$  to get the quotient

$Q(x) = x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0$  and remainder  $R(x) = ux + v$  we have

$$P(x) = (x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0)(x^2 - px - q) + ux + v$$

Matching the coefficients on both sides we get the following relations

- $b_{n-2} = 1, b_{n-1} = b_n = 0$

# Recursive Relations for Division by a Quadratic Polynomial

When we divide polynomial  $P(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$  by a quadratic polynomial  $x^2 - px - q$  to get the quotient  $Q(x) = x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0$  and remainder  $R(x) = ux + v$  we have

$$P(x) = (x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0)(x^2 - px - q) + ux + v$$

Matching the coefficients on both sides we get the following relations

- $b_{n-2} = 1, b_{n-1} = b_n = 0$
- $a_i = b_{i-2} - pb_{i-1} - qb_i$  for  $2 \leq i \leq n$



# Recursive Relations for Division by a Quadratic Polynomial

When we divide polynomial  $P(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$  by a quadratic polynomial  $x^2 - px - q$  to get the quotient  $Q(x) = x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0$  and remainder  $R(x) = ux + v$  we have

$$P(x) = (x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0)(x^2 - px - q) + ux + v$$

Matching the coefficients on both sides we get the following relations

- $b_{n-2} = 1, b_{n-1} = b_n = 0$
- $a_i = b_{i-2} - pb_{i-1} - qb_i$  for  $2 \leq i \leq n$
- $a_1 = -pb_0 - qb_1 + u$

# Recursive Relations for Division by a Quadratic Polynomial

When we divide polynomial  $P(x) = x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$  by a quadratic polynomial  $x^2 - px - q$  to get the quotient  $Q(x) = x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0$  and remainder  $R(x) = ux + v$  we have

$$P(x) = (x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0)(x^2 - px - q) + ux + v$$

Matching the coefficients on both sides we get the following relations

- $b_{n-2} = 1, b_{n-1} = b_n = 0$
- $a_i = b_{i-2} - pb_{i-1} - qb_i$  for  $2 \leq i \leq n$
- $a_1 = -pb_0 - qb_1 + u$
- $a_0 = -qb_0 + v$

# Recursive Relations for Division by a Quadratic Polynomial

$$P(x) = (x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0)(x^2 - px - q) + ux + v$$

Matching the coefficients on both sides we get the following relations

- $b_{n-2} = 1, b_{n-1} = b_n = 0$
- $a_i = b_{i-2} - pb_{i-1} - qb_i$  for  $2 \leq i \leq n$
- $a_1 = -pb_0 - qb_1 + u$
- $a_0 = -qb_0 + v$

from these relations we can define the following recursive (reverse time) system

$$b_i = a_{i+2} + pb_{i+1} + qb_{i+2}$$

starting with initial conditions  $b_{n-2} = 1, b_{n-1} = b_n = 0$ .

# Recursive Relations for Division by a Quadratic Polynomial

$$P(x) = (x^{n-2} + b_{n-3}x^{n-3} + \dots + b_1x + b_0)(x^2 - px - q) + ux + v$$

Matching the coefficients on both sides we get the following relations

- $b_{n-2} = 1, b_{n-1} = b_n = 0$
- $a_i = b_{i-2} - pb_{i-1} - qb_i$  for  $2 \leq i \leq n$
- $a_1 = -pb_0 - qb_1 + u$
- $a_0 = -qb_0 + v$

from these relations we can define the following recursive (reverse time) system

$$b_i = a_{i+2} + pb_{i+1} + qb_{i+2}$$

starting with initial conditions  $b_{n-2} = 1, b_{n-1} = b_n = 0$ . Once  $b_0$  is calculated we proceed to find

$$u = a_1 + pb_0 + qb_1$$

$$v = a_0 + qb_0$$

# Recursive Relations for Division by a Quadratic Polynomial

Matching the coefficients on both sides we get the following relations

- $b_{n-2} = 1, b_{n-1} = b_n = 0$
- $a_i = b_{i-2} - pb_{i-1} - qb_i$  for  $2 \leq i \leq n$
- $a_1 = -pb_0 - qb_1 + u$
- $a_0 = -qb_0 + v$

from these relations we can define the following recursive (reverse time) system

$$b_i = a_{i+2} + pb_{i+1} + qb_{i+2}$$

starting with initial conditions  $b_{n-2} = 1, b_{n-1} = b_n = 0$ . Once  $b_0$  is calculated we proceed to find

$$u = a_1 + pb_0 + qb_1$$

$$v = a_0 + qb_0$$

Synthetic division is a way of implementing this recursive system using a table.

If  $P(x)$  is a polynomial with real coefficients and  $\omega = a + b\mathbf{i}$  is a complex root of  $P(x)$ , then  $\bar{\omega} = a - b\mathbf{i}$  is also a root of  $P(x)$ .

If  $P(x)$  is a polynomial with real coefficients and  $\omega = a + b\mathbf{i}$  is a complex root of  $P(x)$ , then  $\bar{\omega} = a - b\mathbf{i}$  is also a root of  $P(x)$ . In particular,  $(x - \omega)(x - \bar{\omega})$  is a factor of  $P(x)$ .

If  $P(x)$  is a polynomial with real coefficients and  $\omega = a + bi$  is a complex root of  $P(x)$ , then  $\bar{\omega} = a - bi$  is also a root of  $P(x)$ . In particular,  $(x - \omega)(x - \bar{\omega})$  is a factor of  $P(x)$ .

$$(x - \omega)(x - \bar{\omega}) = x^2 - (\omega + \bar{\omega})x + \omega\bar{\omega}$$



If  $P(x)$  is a polynomial with real coefficients and  $\omega = a + b\mathbf{i}$  is a complex root of  $P(x)$ , then  $\bar{\omega} = a - b\mathbf{i}$  is also a root of  $P(x)$ . In particular,  $(x - \omega)(x - \bar{\omega})$  is a factor of  $P(x)$ .

$$(x - \omega)(x - \bar{\omega}) = x^2 - (\omega + \bar{\omega})x + \omega\bar{\omega} = x^2 - 2ax + a^2 + b^2$$

# Bairstow's Method

If  $P(x)$  is a polynomial with real coefficients and  $\omega = a + b\mathbf{i}$  is a complex root of  $P(x)$ , then  $\bar{\omega} = a - b\mathbf{i}$  is also a root of  $P(x)$ . In particular,  $(x - \omega)(x - \bar{\omega})$  is a factor of  $P(x)$ .

$$(x - \omega)(x - \bar{\omega}) = x^2 - (\omega + \bar{\omega})x + \omega\bar{\omega} = x^2 - 2ax + a^2 + b^2$$

Letting  $p = 2a$  and  $q = -a^2 - b^2$ , we can search for the root  $\omega = a + b\mathbf{i}$  by trying to find a quadratic factor  $x^2 - px - q$  of  $P(x)$ .

# Bairstow's Method

If  $P(x)$  is a polynomial with real coefficients and  $\omega = a + bi$  is a complex root of  $P(x)$ , then  $\bar{\omega} = a - bi$  is also a root of  $P(x)$ . In particular,  $(x - \omega)(x - \bar{\omega})$  is a factor of  $P(x)$ .

$$(x - \omega)(x - \bar{\omega}) = x^2 - (\omega + \bar{\omega})x + \omega\bar{\omega} = x^2 - 2ax + a^2 + b^2$$

Letting  $p = 2a$  and  $q = -a^2 - b^2$ , we can search for the root  $\omega = a + bi$  by trying to find a quadratic factor  $x^2 - px - q$  of  $P(x)$ .

In **Bairstow's method** we write the remainder  $u(p, q)x + v(p, q)$  of division of  $P(x)$  by  $x^2 - px - q$  as a function of  $p$  and  $q$ .

# Bairstow's Method

If  $P(x)$  is a polynomial with real coefficients and  $\omega = a + b\mathbf{i}$  is a complex root of  $P(x)$ , then  $\bar{\omega} = a - b\mathbf{i}$  is also a root of  $P(x)$ . In particular,  $(x - \omega)(x - \bar{\omega})$  is a factor of  $P(x)$ .

$$(x - \omega)(x - \bar{\omega}) = x^2 - (\omega + \bar{\omega})x + \omega\bar{\omega} = x^2 - 2ax + a^2 + b^2$$

Letting  $p = 2a$  and  $q = -a^2 - b^2$ , we can search for the root  $\omega = a + b\mathbf{i}$  by trying to find a quadratic factor  $x^2 - px - q$  of  $P(x)$ .

In **Bairstow's method** we write the remainder  $u(p, q)x + v(p, q)$  of division of  $P(x)$  by  $x^2 - px - q$  as a function of  $p$  and  $q$ . The goal is to find values of  $p$  and  $q$  for which both  $u(p, q) = 0$  and  $v(p, q) = 0$ .

# Bairstow's Method

If  $P(x)$  is a polynomial with real coefficients and  $\omega = a + bi$  is a complex root of  $P(x)$ , then  $\bar{\omega} = a - bi$  is also a root of  $P(x)$ . In particular,  $(x - \omega)(x - \bar{\omega})$  is a factor of  $P(x)$ .

$$(x - \omega)(x - \bar{\omega}) = x^2 - (\omega + \bar{\omega})x + \omega\bar{\omega} = x^2 - 2ax + a^2 + b^2$$

Letting  $p = 2a$  and  $q = -a^2 - b^2$ , we can search for the root  $\omega = a + bi$  by trying to find a quadratic factor  $x^2 - px - q$  of  $P(x)$ .

In **Bairstow's method** we write the remainder  $u(p, q)x + v(p, q)$  of division of  $P(x)$  by  $x^2 - px - q$  as a function of  $p$  and  $q$ . The goal is to find values of  $p$  and  $q$  for which both  $u(p, q) = 0$  and  $v(p, q) = 0$ . Starting with initial guess  $(p_0, q_0)$  corresponding with an initial guess of a quadratic factor  $x^2 - p_0x - q_0$  of  $P(x)$  we use Newton-Raphson iterations on the system of two equations  $u(p, q) = 0$  and  $v(p, q) = 0$  in two variables  $p$  and  $q$  to find a sequence  $(p_n, q_n)$  corresponding with quadratic polynomials  $x^2 - p_nx - q_n$  which are estimates of a quadratic factor of  $P(x)$ .

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0			
$p$	0				0
	1				

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0			
$p$	0	$p$			0
	1				



# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0			
$p$	0	$p$			0
	1	$p - 5$			

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$		
$p$	0	$p$			0
	1	$p - 5$			

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$		
$p$	0	$p$	$p^2 - 5p$		0
	1	$p - 5$			

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$		
$p$	0	$p$	$p^2 - 5p$		0
	1	$p - 5$	$p^2 - 5p + q + 1$		

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$		
$p$	0	$p$	$p^2 - 5p$	$p^3 - 5p^2 + qp + p$	0
	1	$p - 5$	$p^2 - 5p + q + 1$		

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$	$qp - 5q$	
$p$	0	$p$	$p^2 - 5p$	$p^3 - 5p^2 + qp + p$	0
	1	$p - 5$	$p^2 - 5p + q + 1$		

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

$q$

$p$

1	-5	1	12	15
0	0	$q$	$qp - 5q$	
0	$p$	$p^2 - 5p$	$p^3 - 5p^2 + qp + p$	0
1	$p - 5$	$p^2 - 5p + q + 1$	$p^3 - 5p^2 + 2qp + p - 5q + 12$	

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$	$qp - 5q$	$qp^2 - 5qp + q^2 + q$
$p$	0	$p$	$p^2 - 5p$	$p^3 - 5p^2 + qp + p$	0
	1	$p - 5$	$p^2 - 5p + q + 1$	$p^3 - 5p^2 + 2qp + p - 5q + 12$	



# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$	$qp - 5q$	$qp^2 - 5qp + q^2 + q$
$p$	0	$p$	$p^2 - 5p$	$p^3 - 5p^2 + qp + p$	0
	1	$p - 5$	$p^2 - 5p + q + 1$	$p^3 - 5p^2 + 2qp + p - 5q + 12$	$qp^2 - 5qp + q^2 + q + 15$

## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$	$qp - 5q$	$qp^2 - 5qp + q^2 + q$
$p$	0	$p$	$p^2 - 5p$	$p^3 - 5p^2 + qp + p$	0
	1	$p - 5$	$p^2 - 5p + q + 1$	$p^3 - 5p^2 + 2qp + p - 5q + 12$	$qp^2 - 5qp + q^2 + q + 15$

Now we use Newton-Raphson's iteration for the system of non-linear equations

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$	$qp - 5q$	$qp^2 - 5qp + q^2 + q$
$p$	0	$p$	$p^2 - 5p$	$p^3 - 5p^2 + qp + p$	0
	1	$p - 5$	$p^2 - 5p + q + 1$	$p^3 - 5p^2 + 2qp + p - 5q + 12$	$qp^2 - 5qp + q^2 + q + 15$

Now we use Newton-Raphson's iteration for the system of non-linear equations

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$U_p(p, q) = 3p^2 - 10p + 2q + 1$$

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

	1	-5	1	12	15
$q$	0	0	$q$	$qp - 5q$	$qp^2 - 5qp + q^2 + q$
$p$	0	$p$	$p^2 - 5p$	$p^3 - 5p^2 + qp + p$	0
	1	$p - 5$	$p^2 - 5p + q + 1$	$p^3 - 5p^2 + 2qp + p - 5q + 12$	$qp^2 - 5qp + q^2 + q + 15$

Now we use Newton-Raphson's iteration for the system of non-linear equations

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$U_p(p, q) = 3p^2 - 10p + 2q + 1$$

$$U_q(p, q) = 2p - 5$$

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

Now we use Newton-Raphson's iteration for the system of non-linear equations

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$U_p(p, q) = 3p^2 - 10p + 2q + 1$$

$$U_q(p, q) = 2p - 5$$

$$V_p(p, q) = 2pq - 5q$$

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

Now we use Newton-Raphson's iteration for the system of non-linear equations

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$U_p(p, q) = 3p^2 - 10p + 2q + 1$$

$$U_q(p, q) = 2p - 5$$

$$V_p(p, q) = 2pq - 5q$$

$$V_q(p, q) = p^2 - 5p + 2q + 1$$

## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial

$P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$U_p(p, q) = 3p^2 - 10p + 2q + 1$$

$$U_q(p, q) = 2p - 5$$

$$V_p(p, q) = 2pq - 5q$$

$$V_q(p, q) = p^2 - 5p + 2q + 1$$

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - \begin{bmatrix} U_p(p_n, q_n) & U_q(p_n, q_n) \\ V_p(p_n, q_n) & V_q(p_n, q_n) \end{bmatrix}^{-1} \begin{bmatrix} U(p_n, q_n) \\ V(p_n, q_n) \end{bmatrix}$$

## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$U_p(p, q) = 3p^2 - 10p + 2q + 1$$

$$U_q(p, q) = 2p - 5$$

$$V_p(p, q) = 2pq - 5q$$

$$V_q(p, q) = p^2 - 5p + 2q + 1$$

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - \begin{bmatrix} U_p(p_n, q_n) & U_q(p_n, q_n) \\ V_p(p_n, q_n) & V_q(p_n, q_n) \end{bmatrix}^{-1} \begin{bmatrix} U(p_n, q_n) \\ V(p_n, q_n) \end{bmatrix}$$

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - \begin{bmatrix} 3p_n^2 - 10p_n + 2q_n + 1 & 2p_n - 5 \\ 2p_nq_n - 5q_n & p_n^2 - 5p_n + 2q_n + 1 \end{bmatrix}^{-1} \begin{bmatrix} p^3 - 5p^2 + 2qp + p - 5q + 12 \\ qp^2 - 5qp + q^2 + q + 15 \end{bmatrix}$$



## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial

$P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - \begin{bmatrix} U_p(p_n, q_n) & U_q(p_n, q_n) \\ V_p(p_n, q_n) & V_q(p_n, q_n) \end{bmatrix}^{-1} \begin{bmatrix} U(p_n, q_n) \\ V(p_n, q_n) \end{bmatrix}$$

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - \begin{bmatrix} 3p_n^2 - 10p_n + 2q_n + 1 & 2p_n - 5 \\ 2p_n q_n - 5q_n & p_n^2 - 5p_n + 2q_n + 1 \end{bmatrix}^{-1} \begin{bmatrix} p^3 - 5p^2 + 2qp + p - 5q + 12 \\ qp^2 - 5qp + q^2 + q + 15 \end{bmatrix}$$

$$p_0 = -2 \text{ and } q_0 = -1$$

## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial

$P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - \begin{bmatrix} U_p(p_n, q_n) & U_q(p_n, q_n) \\ V_p(p_n, q_n) & V_q(p_n, q_n) \end{bmatrix}^{-1} \begin{bmatrix} U(p_n, q_n) \\ V(p_n, q_n) \end{bmatrix}$$

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - \begin{bmatrix} 3p_n^2 - 10p_n + 2q_n + 1 & 2p_n - 5 \\ 2p_nq_n - 5q_n & p_n^2 - 5p_n + 2q_n + 1 \end{bmatrix}^{-1} \begin{bmatrix} p^3 - 5p^2 + 2qp + p - 5q + 12 \\ qp^2 - 5qp + q^2 + q + 15 \end{bmatrix}$$

$$p_0 = -2 \text{ and } q_0 = -1$$

$$p_1 = -1.7768595041322314050 \text{ and } q_1 = -1.23140495867768595040$$

## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - \begin{bmatrix} 3p_n^2 - 10p_n + 2q_n + 1 & 2p_n - 5 \\ 2p_nq_n - 5q_n & p_n^2 - 5p_n + 2q_n + 1 \end{bmatrix}^{-1} \begin{bmatrix} p^3 - 5p^2 + 2qp + p - 5q + 12 \\ qp^2 - 5qp + q^2 + q + 15 \end{bmatrix}$$

$$p_0 = -2 \text{ and } q_0 = -1$$

$$p_1 = -1.7768595041322314050 \text{ and } q_1 = -1.23140495867768595040$$

$$p_2 = -1.76897527126986 \text{ and } q_2 = -1.28245105573983$$

# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial

$P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - \begin{bmatrix} 3p_n^2 - 10p_n + 2q_n + 1 & 2p_n - 5 \\ 2p_nq_n - 5q_n & p_n^2 - 5p_n + 2q_n + 1 \end{bmatrix}^{-1} \begin{bmatrix} p^3 - 5p^2 + 2qp + p - 5q + 12 \\ qp^2 - 5qp + q^2 + q + 15 \end{bmatrix}$$

$$p_0 = -2 \text{ and } q_0 = -1$$

$$p_1 = -1.7768595041322314050 \text{ and } q_1 = -1.23140495867768595040$$

$$p_2 = -1.76897527126986 \text{ and } q_2 = -1.28245105573983$$

$$p_3 = -1.76907525986458 \text{ and } q_3 = -1.28291926483910$$

## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

$$\begin{cases} U(p, q) = p^3 - 5p^2 + 2qp + p - 5q + 12 = 0 \\ V(p, q) = qp^2 - 5qp + q^2 + q + 15 = 0 \end{cases}$$

$$p_0 = -2 \text{ and } q_0 = -1$$

$$p_1 = -1.7768595041322314050 \text{ and } q_1 = -1.23140495867768595040$$

$$p_2 = -1.76897527126986 \text{ and } q_2 = -1.28245105573983$$

After three steps of Newton-Raphson iteration, we announce  $x^2 + 1.76907525986458x + 1.28291926483910$  as a quadratic factor of  $P(x)$ .

## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

After three steps of Newton-Raphson iteration, we announce  $x^2 + 1.76907525986458x + 1.28291926483910$  as a quadratic factor of  $P(x)$ .

This gives us  $a = -\frac{1.76907525986458}{2} = -0.8845376265$  and  $b = \pm\sqrt{1.28291926483910 - a^2} = 0.7074690370$  which give us the two conjugate roots  $\omega = -0.8845376265 + 0.7074690370i$  and  $\bar{\omega} = -0.8845376265 - 0.7074690370i$ .

## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

After three steps of Newton-Raphson iteration, we announce  $x^2 + 1.76907525986458x + 1.28291926483910$  as a quadratic factor of  $P(x)$ .

This gives us  $a = -\frac{1.76907525986458}{2} = -0.8845376265$  and  $b = \pm\sqrt{1.28291926483910 - a^2} = 0.7074690370$  which give us the two conjugate roots  $\omega = -0.8845376265 + 0.7074690370i$  and  $\bar{\omega} = -0.8845376265 - 0.7074690370i$ .

In this example since  $P(x)$  has degree four and the polynomial  $q(x) = x^2 + 1.76907525986458x + 1.28291926483910$  is a factor of  $P(x)$ , the quotient of division of  $P(x)$  by  $q(x)$  is also a quadratic factor of  $P(x)$ .

## Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

This gives us  $a = -\frac{1.76907525986458}{2} = -0.8845376265$  and  $b = \pm\sqrt{1.28291926483910 - a^2} = 0.7074690370$  which give us the two conjugate roots  $\omega = -0.8845376265 + 0.7074690370i$  and  $\bar{\omega} = -0.8845376265 - 0.7074690370i$ .

In this example since  $P(x)$  has degree four and the polynomial  $q(x) = x^2 + 1.76907525986458x + 1.28291926483910$  is a factor of  $P(x)$ , the quotient of division of  $P(x)$  by  $q(x)$  is also a quadratic factor of  $P(x)$ .

In the synthetic division table earlier on we found the quotient to be  $x^2 + (p - 5)x + p^2 - 5p + q + 1$ .



# Example

Use Bairstow's method to estimate a quadratic factor and its corresponding pair of complex roots for the polynomial  $P(x) = x^4 - 5x^3 + x^2 + 12x + 15$ . Use initial guess  $(p_0, q_0) = (-2, -1)$ .

In this example since  $P(x)$  has degree four and the polynomial  $q(x) = x^2 + 1.76907525986458x + 1.28291926483910$  is a factor of  $P(x)$ , the quotient of division of  $P(x)$  by  $q(x)$  is also a quadratic factor of  $P(x)$ .

In the synthetic division table earlier on we found the quotient to be  $x^2 + (p - 5)x + p^2 - 5p + q + 1$ .

Using the values of  $p_3$  and  $q_3$  for  $p$  and  $q$  we get the quotient to be approximated as  $x^2 - 6.769075260 * x + 11.69208432$  and the other two conjugate roots of  $P(x)$  to be  $3.384537630 - 0.4868155206i, 3.384537630 + 0.4868155206i$ .