

## (2.2) Fixed-Point Iteration

MATH 4701 Numerical Analysis


# Fixed Points and Solving Equations

We say  $p$  is a **fixed point** for a given function  $g$  if  $g(p) = p$ .

# Fixed Points and Solving Equations

We say  $p$  is a **fixed point** for a given function  $g$  if  $g(p) = p$ .

Fixed points of a function  $g(x)$  correspond with the points where its graph intersects the line  $y = x$ .



# Fixed Points and Solving Equations

We say  $p$  is a **fixed point** for a given function  $g$  if  $g(p) = p$ .

Fixed points of a function  $g(x)$  correspond with the points where its graph intersects the line  $y = x$ .

Given an equation  $f(x) = 0$ , if we defined  $g(x) = f(x) + x$ , then a fixed point  $p$  for  $g(x)$  is a solution of for  $f(x) = 0$ .

# Fixed Points and Solving Equations

We say  $p$  is a **fixed point** for a given function  $g$  if  $g(p) = p$ .

Fixed points of a function  $g(x)$  correspond with the points where its graph intersects the line  $y = x$ .

Given an equation  $f(x) = 0$ , if we defined  $g(x) = f(x) + x$ , then a fixed point  $p$  for  $g(x)$  is a solution of for  $f(x) = 0$ .

There are, however, numerous ways solving an equation can be turned into a question of finding a fixed point of another function.

# Fixed Points and Solving Equations

We say  $p$  is a **fixed point** for a given function  $g$  if  $g(p) = p$ .

Fixed points of a function  $g(x)$  correspond with the points where its graph intersects the line  $y = x$ .

Given an equation  $f(x) = 0$ , if we defined  $g(x) = f(x) + x$ , then a fixed point  $p$  for  $g(x)$  is a solution of for  $f(x) = 0$ .

There are, however, numerous ways solving an equation can be turned into a question of finding a fixed point of another function.

For example, for  $x^2 - 2 = 0$ , we can define  $g(x) = x^2 + x - 2$  and find fixed points of  $g(x)$  ( $x^2 + x - 2 = x \Leftrightarrow x^2 - 2 = 0$ ).

# Fixed Points and Solving Equations

We say  $p$  is a **fixed point** for a given function  $g$  if  $g(p) = p$ .

Fixed points of a function  $g(x)$  correspond with the points where its graph intersects the line  $y = x$ .

Given an equation  $f(x) = 0$ , if we defined  $g(x) = f(x) + x$ , then a fixed point  $p$  for  $g(x)$  is a solution of for  $f(x) = 0$ .

There are, however, numerous ways solving an equation can be turned into a question of finding a fixed point of another function.

For example, for  $x^2 - 2 = 0$ , we can define  $g(x) = x^2 + x - 2$  and find fixed points of  $g(x)$  ( $x^2 + x - 2 = x \Leftrightarrow x^2 - 2 = 0$ ).

On the other hand, we can define  $g(x) = \frac{2}{x}$  and find fixed points of  $g(x)$  ( $\frac{2}{x} = x \Leftrightarrow x^2 - 2 = 0$ ).

# Fixed Points and Solving Equations

We say  $p$  is a **fixed point** for a given function  $g$  if  $g(p) = p$ .

Given an equation  $f(x) = 0$ , if we defined  $g(x) = f(x) + x$ , then a fixed point  $p$  for  $g(x)$  is a solution of for  $f(x) = 0$ .

There are, however, numerous ways solving an equation can be turned into a question of finding a fixed point of another function.

For example, for  $x^2 - 2 = 0$ , we can define  $g(x) = x^2 + x - 2$  and find fixed points of  $g(x)$  ( $x^2 + x - 2 = x \Leftrightarrow x^2 - 2 = 0$ ).

On the other hand, we can define  $g(x) = \frac{2}{x}$  and find fixed points of  $g(x)$  ( $\frac{2}{x} = x \Leftrightarrow x^2 - 2 = 0$ ).

Or we can define  $g(x) = \frac{x^2+2}{2x}$  and find fixed points of  $g(x)$  ( $\frac{x^2+2}{2x} = x \Leftrightarrow x^2 + 2 = 2x^2 \Leftrightarrow x^2 - 2 = 0$ ).



# Fixed-Point Iteration

For a function  $g : [a, b] \rightarrow \mathbb{R}$ , and an **initial guess**  $p_0$ , we can consider the sequence of iterations of  $g$  defined by  $x_0 = p_0$  and  $x_{n+1} = g(x_n)$ .

# Fixed-Point Iteration

For a function  $g : [a, b] \rightarrow \mathbb{R}$ , and an **initial guess**  $p_0$ , we can consider the sequence of iterations of  $g$  defined by  $x_0 = p_0$  and  $x_{n+1} = g(x_n)$ .

When this sequence converges, its limit is a fixed point of  $g$ .

# Fixed-Point Iteration

For a function  $g : [a, b] \rightarrow \mathbb{R}$ , and an **initial guess**  $p_0$ , we can consider the sequence of iterations of  $g$  defined by  $x_0 = p_0$  and  $x_{n+1} = g(x_n)$ .

When this sequence converges, its limit is a fixed point of  $g$ .

But it is possible that this sequence does not converge or it converges slowly.

# Fixed-Point Iteration

For a function  $g : [a, b] \rightarrow \mathbb{R}$ , and an **initial guess**  $p_0$ , we can consider the sequence of iterations of  $g$  defined by  $x_0 = p_0$  and  $x_{n+1} = g(x_n)$ .

When this sequence converges, its limit is a fixed point of  $g$ .

But it is possible that this sequence does not converge or it converges slowly.

For example, for  $g(x) = x^2 + x - 2$  with  $x_0 = 4$ , we get

$$x_0 = 4 \quad x_1 = g(x_0) = 18 \quad x_2 = g(x_1) = 340 \quad x_3 = g(x_2) = 115938$$

diverges to  $+\infty$ .

# Fixed-Point Iteration

For a function  $g : [a, b] \rightarrow \mathbb{R}$ , and an **initial guess**  $p_0$ , we can consider the sequence of iterations of  $g$  defined by  $x_0 = p_0$  and  $x_{n+1} = g(x_n)$ .

When this sequence converges, its limit is a fixed point of  $g$ .

But it is possible that this sequence does not converge or it converges slowly.

For  $g(x) = \frac{2}{x}$  with  $x_0 = 4$ , we get

$$x_0 = 4 \quad x_1 = g(x_0) = 0.5 \quad x_2 = g(x_1) = 4 \quad x_3 = g(x_2) = 0.5$$

forms a divergent periodic cycle.

# Fixed-Point Iteration

For a function  $g : [a, b] \rightarrow \mathbb{R}$ , and an **initial guess**  $p_0$ , we can consider the sequence of iterations of  $g$  defined by  $x_0 = p_0$  and  $x_{n+1} = g(x_n)$ .

When this sequence converges, its limit is a fixed point of  $g$ .

But it is possible that this sequence does not converge or it converges slowly.

But for  $g(x) = \frac{x^2+2}{2x}$  with  $x_0 = 4$ , we get

$$x_0 = 4 \quad x_1 = g(x_0) = \frac{9}{4} = 2.25 \quad x_2 = g(x_1) = \frac{113}{72} = 1.569\bar{4}$$


$$x_3 = g(x_2) = \frac{23137}{16272} \approx 1.42189036 \quad x_4 = g(x_3) \approx 1.414234286$$

Converges to  $\sqrt{2} \approx 1.41421356$  which is a fixed point of  $g(x) = \frac{x^2+2}{2x}$  and a solution of  $f(x) = x^2 - 2 = 0$ .

# Fixed-Point Iteration

For a function  $g : [a, b] \rightarrow \mathbb{R}$ , and an **initial guess**  $p_0$ , we can consider the sequence of iterations of  $g$  defined by  $x_0 = p_0$  and  $x_{n+1} = g(x_n)$ .


The sequence can be geometrically traced by starting at the value of any  $x_n$  on the  $x$ -axis, drawing a vertical line until we hit the graph of  $g$ , from that point drawing a horizontal line to hit the line  $y = x$ , and finally drawing a vertical line until we hit the  $x$ -axis at the next number  $x_{n+1}$ .



# Mean Value Theorem and Estimate on Distance

**Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , then there is  $\alpha \in (a, b)$  such that

$$f'(\alpha) = \frac{f(b) - f(a)}{b - a}$$






# Mean Value Theorem and Estimate on Distance

**Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , then there is  $\alpha \in (a, b)$  such that

$$f'(\alpha) = \frac{f(b) - f(a)}{b - a}$$

In other words, on the graph of the function, there is a point in between the end-points where the slope of the tangent line is equal to the slope of the secant line through the two end-points.




# Mean Value Theorem and Estimate on Distance

**Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , then there is  $\alpha \in (a, b)$  such that

$$f'(\alpha) = \frac{f(b) - f(a)}{b - a}$$

Notice that the value  $\alpha$  may not be unique and the theorem only claims the existence of at least one such value.



# Mean Value Theorem and Estimate on Distance

**Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , then there is  $\alpha \in (a, b)$  such that

$$f'(\alpha) = \frac{f(b) - f(a)}{b - a}$$

In addition to being one of the most important theoretical theorems that is used in proving numerous other theorems in Calculus, the Mean Value Theorem allows you to estimate how much the distance between two numbers changes after you apply the function  $f$  to both of them.

# Mean Value Theorem and Estimate on Distance

**Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , then there is  $\alpha \in (a, b)$  such that

$$f'(\alpha) = \frac{f(b) - f(a)}{b - a}$$

**An Important Consequence of the Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and if there is a positive number  $K$  such that for all  $x \in (a, b)$ ,  $|f'(x)| \leq K$ , then for any  $x$  and  $y$  in  $[a, b]$ ,

$$|f(x) - f(y)| \leq K|x - y|$$

# Mean Value Theorem and Estimate on Distance

**Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , then there is  $\alpha \in (a, b)$  such that

$$f'(\alpha) = \frac{f(b) - f(a)}{b - a}$$

**An Important Consequence of the Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and if there is a positive number  $K$  such that for all  $x \in (a, b)$ ,  $|f'(x)| \leq K$ , then for any  $x$  and  $y$  in  $[a, b]$ ,

$$|f(x) - f(y)| \leq K|x - y|$$

Notice that by the Mean Value Theorem, there is  $\alpha$  between  $x$  and  $y$  such that

$$\frac{f(x) - f(y)}{x - y} = f'(\alpha)$$

# Mean Value Theorem and Estimate on Distance

**Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , then there is  $\alpha \in (a, b)$  such that

$$f'(\alpha) = \frac{f(b) - f(a)}{b - a}$$

**An Important Consequence of the Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and if there is a positive number  $K$  such that for all  $x \in (a, b)$ ,  $|f'(x)| \leq K$ , then for any  $x$  and  $y$  in  $[a, b]$ ,

$$|f(x) - f(y)| \leq K|x - y|$$

Notice that by the Mean Value Theorem, there is  $\alpha$  between  $x$  and  $y$  such that

$$\frac{f(x) - f(y)}{x - y} = f'(\alpha) \Rightarrow \frac{|f(x) - f(y)|}{|x - y|} = |f'(\alpha)| \leq K$$

# Mean Value Theorem and Estimate on Distance

**Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , then there is  $\alpha \in (a, b)$  such that

$$f'(\alpha) = \frac{f(b) - f(a)}{b - a}$$

**An Important Consequence of the Mean Value Theorem:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$ , and if there is a positive number  $K$  such that for all  $x \in (a, b)$ ,  $|f'(x)| \leq K$ , then for any  $x$  and  $y$  in  $[a, b]$ ,

$$|f(x) - f(y)| \leq K|x - y|$$

Notice that by the Mean Value Theorem, there is  $\alpha$  between  $x$  and  $y$  such that

$$\Rightarrow \frac{|f(x) - f(y)|}{|x - y|} = |f'(\alpha)| \leq K \Rightarrow |f(x) - f(y)| \leq K|x - y|$$

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .



# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .

A function  $g$  does not need to satisfy the conditions of this theorem to have a fixed point but these conditions do imply that  $g$  has a fixed point.

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .

This part can be proven as follows.

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .

This part can be proven as follows. if  $g(a) = a$  or  $g(b) = b$  then we are done so assume  $g(a) \neq a$  and  $g(b) \neq b$ .

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .

This part can be proven as follows. if  $g(a) = a$  or  $g(b) = b$  then we are done so assume  $g(a) \neq a$  and  $g(b) \neq b$ . Define  $h(x) = g(x) - x$ .

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .

This part can be proven as follows. if  $g(a) = a$  or  $g(b) = b$  then we are done so assume  $g(a) \neq a$  and  $g(b) \neq b$ . Define  $h(x) = g(x) - x$ . Since  $g(x)$  and  $i(x) = x$  are both continuous on  $[a, b]$ ,  $h(x) = g(x) - x$  is also continuous on  $[a, b]$ .

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .

This part can be proven as follows. if  $g(a) = a$  or  $g(b) = b$  then we are done so assume  $g(a) \neq a$  and  $g(b) \neq b$ . Define  $h(x) = g(x) - x$ . Since  $g(x)$  and  $i(x) = x$  are both continuous on  $[a, b]$ ,  $h(x) = g(x) - x$  is also continuous on  $[a, b]$ . Since  $g([a, b]) \subset [a, b]$  and  $g(a) \neq a$ , we have  $g(a) > a$  which implies  $h(a) > 0$ .

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .

This part can be proven as follows. if  $g(a) = a$  or  $g(b) = b$  then we are done so assume  $g(a) \neq a$  and  $g(b) \neq b$ . Define  $h(x) = g(x) - x$ . Since  $g(x)$  and  $i(x) = x$  are both continuous on  $[a, b]$ ,  $h(x) = g(x) - x$  is also continuous on  $[a, b]$ . Since  $g([a, b]) \subset [a, b]$  and  $g(a) \neq a$ , we have  $g(a) > a$  which implies  $h(a) > 0$ . Similarly, since  $g([a, b]) \subset [a, b]$  and  $g(b) \neq b$ , we have  $g(b) < b$  which implies  $h(b) < 0$ .

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .

This part can be proven as follows. if  $g(a) = a$  or  $g(b) = b$  then we are done so assume  $g(a) \neq a$  and  $g(b) \neq b$ . Define  $h(x) = g(x) - x$ . Since  $g(x)$  and  $i(x) = x$  are both continuous on  $[a, b]$ ,  $h(x) = g(x) - x$  is also continuous on  $[a, b]$ . Since  $g([a, b]) \subset [a, b]$  and  $g(a) \neq a$ , we have  $g(a) > a$  which implies  $h(a) > 0$ . Similarly, since  $g([a, b]) \subset [a, b]$  and  $g(b) \neq b$ , we have  $g(b) < b$  which implies  $h(b) < 0$ . Therefore, by the **Intermediate Value Theorem**, there is  $p \in (a, b)$  such that  $h(p) = g(p) - p = 0$ .



# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .

This part can be proven as follows. if  $g(a) = a$  or  $g(b) = b$  then we are done so assume  $g(a) \neq a$  and  $g(b) \neq b$ . Define  $h(x) = g(x) - x$ . Since  $g(x)$  and  $i(x) = x$  are both continuous on  $[a, b]$ ,  $h(x) = g(x) - x$  is also continuous on  $[a, b]$ . Since  $g([a, b]) \subset [a, b]$  and  $g(a) \neq a$ , we have  $g(a) > a$  which implies  $h(a) > 0$ . Similarly, since  $g([a, b]) \subset [a, b]$  and  $g(b) \neq b$ , we have  $g(b) < b$  which implies  $h(b) < 0$ . Therefore, by the **Intermediate Value Theorem**, there is  $p \in (a, b)$  such that  $h(p) = g(p) - p = 0$ . Hence,  $g(p) = p$  and  $g$  has a fixed point  $p \in [a, b]$ .

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .
- (2) If in addition to conditions of part (1) we also assume  $g$  is differentiable on  $(a, b)$  and if there is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  $|g'(x)| \leq K$ , then  $g$  has **exactly one fixed point** inside  $[a, b]$ .

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .
- (2) If in addition to conditions of part (1) we also assume  $g$  is differentiable on  $(a, b)$  and if there is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  $|g'(x)| \leq K$ , then  $g$  has **exactly one fixed point** inside  $[a, b]$ .

A function  $g$  does not need to satisfy the conditions of this theorem to have a unique fixed point but these conditions do imply that  $g$  has a unique fixed point.

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .
- (2) If in addition to conditions of part (1) we also assume  $g$  is differentiable on  $(a, b)$  and if there is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  $|g'(x)| \leq K$ , then  $g$  has **exactly one fixed point** inside  $[a, b]$ .

To see why the assertion of this part is true, notice that if  $p$  and  $q$  are both fixed points of  $g$  and they are both in  $[a, b]$ , by the **Mean Value Theorem**,

$$|p - q| = |g(p) - g(q)| \leq K|p - q|$$

# Existence and Uniqueness Theorem for Fixed Points

## Existence and Uniqueness Theorem for Fixed Points:

- (1) If  $g : [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  ( $g \in C[a, b]$ ) and  $g(x) \in [a, b]$  for all  $x \in [a, b]$  ( $g([a, b]) \subset [a, b]$ ) then  $g$  has at least one fixed point in  $[a, b]$ .
- (2) If in addition to conditions of part (1) we also assume  $g$  is differentiable on  $(a, b)$  and if there is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  $|g'(x)| \leq K$ , then  $g$  has **exactly one fixed point** inside  $[a, b]$ .

To see why the assertion of this part is true, notice that if  $p$  and  $q$  are both fixed points of  $g$  and they are both in  $[a, b]$ , by the **Mean Value Theorem**,

$|p - q| = |g(p) - g(q)| \leq K|p - q| \Rightarrow 1 \leq K$  which contradicts the assumption on  $K$  being strictly less than 1.

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$



# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ .

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ . Moreover, by the theorem we just proved  $g$  has a unique fixed point  $p$  in  $[a, b]$ .

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ . Moreover, by the theorem we just proved  $g$  has a unique fixed point  $p$  in  $[a, b]$ .

$$|p_n - p| = |g(p_{n-1}) - p|$$

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ . Moreover, by the theorem we just proved  $g$  has a unique fixed point  $p$  in  $[a, b]$ .

$$|p_n - p| = |g(p_{n-1}) - p| = |g(p_{n-1}) - g(p)|$$

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ . Moreover, by the theorem we just proved  $g$  has a unique fixed point  $p$  in  $[a, b]$ .

$$|p_n - p| = |g(p_{n-1}) - p| = |g(p_{n-1}) - g(p)| \leq K |p_{n-1} - p|$$

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ .

$$|p_n - p| = |g(p_{n-1}) - p| = |g(p_{n-1}) - g(p)| \leq K|p_{n-1} - p|$$

$$|p_n - p| \leq K|p_{n-1} - p|$$



# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ .

$$|p_n - p| = |g(p_{n-1}) - p| = |g(p_{n-1}) - g(p)| \leq K|p_{n-1} - p|$$

$$|p_n - p| \leq K|p_{n-1} - p| \leq K^2|p_{n-2} - p| \leq \dots \leq K^n|p_0 - p|$$

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ .

$$|p_n - p| \leq K|p_{n-1} - p| \leq K^2|p_{n-2} - p| \leq \dots \leq K^n|p_0 - p|$$
$$\lim_{n \rightarrow \infty} |p_n - p| \leq \lim_{n \rightarrow \infty} K^n|p_0 - p|$$

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ .

$$\begin{aligned} |p_n - p| &\leq K|p_{n-1} - p| \leq K^2|p_{n-2} - p| \leq \dots \leq K^n|p_0 - p| \\ \lim_{n \rightarrow \infty} |p_n - p| &\leq \lim_{n \rightarrow \infty} K^n|p_0 - p| = 0 \quad (\text{because } 0 < K < 1 \text{ implies } \lim_{n \rightarrow \infty} K^n = 0). \end{aligned}$$

# Fixed-Point Iteration Theorem

**Fixed-Point Iteration Theorem:** Suppose

- $g \in C[a, b]$
- $g([a, b]) \subset [a, b]$
- $g$  is differentiable on  $(a, b)$
- There is a positive number  $K < 1$  such that for all  $x \in (a, b)$ ,  
 $|g'(x)| \leq K$

Then for any initial guess  $p_0 \in [a, b]$ , the sequence  $p_n$  defined by  $p_{n+1} = g(p_n)$  is convergent to the unique fixed point of  $g$  in  $[a, b]$ .

**Proof:** Since  $g([a, b]) \subset [a, b]$ , all the entries  $p_n$  in the sequence are defined and they are all in  $[a, b]$ .

$\lim_{n \rightarrow \infty} |p_n - p| \leq \lim_{n \rightarrow \infty} K^n |p_0 - p| = 0$  (because  $0 < K < 1$  implies  $\lim_{n \rightarrow \infty} K^n = 0$ ). Hence,  $\lim_{n \rightarrow \infty} p_n = p$ .

# Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

# Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

$g'(x) = \frac{1}{2} - \frac{1}{x^2} = 0$  only when  $x^2 = 2$  or  $x = \pm\sqrt{2}$ .

# Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

$g'(x) = \frac{1}{2} - \frac{1}{x^2} = 0$  only when  $x^2 = 2$  or  $x = \pm\sqrt{2}$ . This means  $g(x)$  has only one critical number inside  $[1, 8]$  at  $x = \sqrt{2}$ .

# Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

$g'(x) = \frac{1}{2} - \frac{1}{x^2} = 0$  only when  $x^2 = 2$  or  $x = \pm\sqrt{2}$ . This means  $g(x)$  has only one critical number inside  $[1, 8]$  at  $x = \sqrt{2}$ .

Checking the values of  $g$  at its only critical number within  $[1, 8]$  and at the two end points of  $[1, 8]$  we have  $g(1) = \frac{3}{2}$ ,  $g(8) = \frac{33}{8}$ , and  $g(\sqrt{2}) = \sqrt{2}$ .



# Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

$g'(x) = \frac{1}{2} - \frac{1}{x^2} = 0$  only when  $x^2 = 2$  or  $x = \pm\sqrt{2}$ . This means  $g(x)$  has only one critical number inside  $[1, 8]$  at  $x = \sqrt{2}$ .

Checking the values of  $g$  at its only critical number within  $[1, 8]$  and at the two end points of  $[1, 8]$  we have  $g(1) = \frac{3}{2}$ ,  $g(8) = \frac{33}{8}$ , and  $g(\sqrt{2}) = \sqrt{2}$ .

Since  $\sqrt{2} < \frac{3}{2} < \frac{33}{8}$ , we have  $g([1, 8]) = [\sqrt{2}, \frac{33}{8}] \subset [1, 8]$

## Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

$g'(x) = \frac{1}{2} - \frac{1}{x^2} = 0$  only when  $x^2 = 2$  or  $x = \pm\sqrt{2}$ . This means  $g(x)$  has only one critical number inside  $[1, 8]$  at  $x = \sqrt{2}$ .

Checking the values of  $g$  at its only critical number within  $[1, 8]$  and at the two end points of  $[1, 8]$  we have  $g(1) = \frac{3}{2}$ ,  $g(8) = \frac{33}{8}$ , and  $g(\sqrt{2}) = \sqrt{2}$ .

Since  $\sqrt{2} < \frac{3}{2} < \frac{33}{8}$ , we have  $g([1, 8]) = [\sqrt{2}, \frac{33}{8}] \subset [1, 8]$

$g'(x)$  is increasing on  $[1, 8]$  (by increasing the value of  $x$  the value of  $\frac{1}{2} - \frac{1}{x^2}$  increases as well).

## Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

$g'(x) = \frac{1}{2} - \frac{1}{x^2} = 0$  only when  $x^2 = 2$  or  $x = \pm\sqrt{2}$ . This means  $g(x)$  has only one critical number inside  $[1, 8]$  at  $x = \sqrt{2}$ .

Checking the values of  $g$  at its only critical number within  $[1, 8]$  and at the two end points of  $[1, 8]$  we have  $g(1) = \frac{3}{2}$ ,  $g(8) = \frac{33}{8}$ , and  $g(\sqrt{2}) = \sqrt{2}$ .

Since  $\sqrt{2} < \frac{3}{2} < \frac{33}{8}$ , we have  $g([1, 8]) = [\sqrt{2}, \frac{33}{8}] \subset [1, 8]$

$g'(x)$  is increasing on  $[1, 8]$  (by increasing the value of  $x$  the value of  $\frac{1}{2} - \frac{1}{x^2}$  increases as well).  $g'(1) = -\frac{1}{2}$  and  $g'(8) = \frac{31}{64}$ .

## Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

$g'(x) = \frac{1}{2} - \frac{1}{x^2} = 0$  only when  $x^2 = 2$  or  $x = \pm\sqrt{2}$ . This means  $g(x)$  has only one critical number inside  $[1, 8]$  at  $x = \sqrt{2}$ .

Checking the values of  $g$  at its only critical number within  $[1, 8]$  and at the two end points of  $[1, 8]$  we have  $g(1) = \frac{3}{2}$ ,  $g(8) = \frac{33}{8}$ , and  $g(\sqrt{2}) = \sqrt{2}$ .

Since  $\sqrt{2} < \frac{3}{2} < \frac{33}{8}$ , we have  $g([1, 8]) = [\sqrt{2}, \frac{33}{8}] \subset [1, 8]$

$g'(x)$  is increasing on  $[1, 8]$  (by increasing the value of  $x$  the value of  $\frac{1}{2} - \frac{1}{x^2}$  increases as well).  $g'(1) = -\frac{1}{2}$  and  $g'(8) = \frac{31}{64}$ . As a result  $-\frac{1}{2} \leq g'(x) \leq \frac{31}{64}$  for all  $x \in [1, 8]$ . In particular,  $|g'(x)| \leq \frac{1}{2}$  for all  $x \in [1, 8]$ . ( $\frac{31}{64} < \frac{1}{2}$ )

## Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

Checking the values of  $g$  at its only critical number within  $[1, 8]$  and at the two end points of  $[1, 8]$  we have  $g(1) = \frac{3}{2}$ ,  $g(8) = \frac{33}{8}$ , and  $g(\sqrt{2}) = \sqrt{2}$ .

Since  $\sqrt{2} < \frac{3}{2} < \frac{33}{8}$ , we have  $g([1, 8]) = [\sqrt{2}, \frac{33}{8}] \subset [1, 8]$

$g'(x)$  is increasing on  $[1, 8]$ .  $g'(1) = -\frac{1}{2}$  and  $g'(8) = \frac{31}{64}$ . As a result  $-\frac{1}{2} \leq g'(x) \leq \frac{31}{64}$  for all  $x \in [1, 8]$ . In particular,  $|g'(x)| \leq \frac{1}{2}$  for all  $x \in [1, 8]$ . ( $\frac{31}{64} < \frac{1}{2}$ )

We have seen that  $g(x)$  is continuous on  $[1, 8]$ , differentiable on  $(1, 8)$ ,  $g([1, 8]) \subset [1, 8]$  and  $|g'(x)| \leq \frac{1}{2}$  for all  $x \in [1, 8]$ .

## Example

For  $g(x) = \frac{x^2+2}{2x} = \frac{x}{2} + \frac{1}{x}$  we choose interval  $[1, 8]$ .

Checking the values of  $g$  at its only critical number within  $[1, 8]$  and at the two end points of  $[1, 8]$  we have  $g(1) = \frac{3}{2}$ ,  $g(8) = \frac{33}{8}$ , and  $g(\sqrt{2}) = \sqrt{2}$ .

Since  $\sqrt{2} < \frac{3}{2} < \frac{33}{8}$ , we have  $g([1, 8]) = [\sqrt{2}, \frac{33}{8}] \subset [1, 8]$

$g'(x)$  is increasing on  $[1, 8]$ .  $g'(1) = -\frac{1}{2}$  and  $g'(8) = \frac{31}{64}$ . As a result  $-\frac{1}{2} \leq g'(x) \leq \frac{31}{64}$  for all  $x \in [1, 8]$ . In particular,  $|g'(x)| \leq \frac{1}{2}$  for all  $x \in [1, 8]$ . ( $\frac{31}{64} < \frac{1}{2}$ )

We have seen that  $g(x)$  is continuous on  $[1, 8]$ , differentiable on  $(1, 8)$ ,  $g([1, 8]) \subset [1, 8]$  and  $|g'(x)| \leq \frac{1}{2}$  for all  $x \in [1, 8]$ .

Since  $K = \frac{1}{2} < 1$  by the two theorems we discussed,  $g$  has a unique fixed point  $p$  in  $[1, 8]$  and the sequence of iterations  $g(p_{n+1}) = g(p_n)$  converges to  $p$  for any choice of  $p_0$  in  $[1, 8]$ .

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$



# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$  and either of the following two inequalities can be used to estimate the error in calculation of the fixed point using the function iteration sequence.

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$  and either of the following two inequalities can be used to estimate the error in calculation of the fixed point using the function iteration sequence.

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$  and either of the following two inequalities can be used to estimate the error in calculation of the fixed point using the function iteration sequence.

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$  and either of the following two inequalities can be used to estimate the error in calculation of the fixed point using the function iteration sequence.

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

We saw in the proof of the **fixed point iteration theorem** that

$$|p_n - p| \leq K^n |p_0 - p|$$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$  and either of the following two inequalities can be used to estimate the error in calculation of the fixed point using the function iteration sequence.

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

We saw in the proof of the **fixed point iteration theorem** that

$$|p_n - p| \leq K^n |p_0 - p| \leq K^n \max\{p_0 - a, b - p_0\}$$

Distance between  $p$  and  $p_0$  which is within  $[a, b]$  is less than the larger value of the distance between  $p$  and the two end-points of  $[a, b]$ .

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$  and either of the following two inequalities can be used to estimate the error in calculation of the fixed point using the function iteration sequence.

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$  and either of the following two inequalities can be used to estimate the error in calculation of the fixed point using the function iteration sequence.

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

$$|p_{n+1} - p_n| = |g(p_n) - g(p_{n-1})| \leq K |p_n - p_{n-1}|$$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$  and either of the following two inequalities can be used to estimate the error in calculation of the fixed point using the function iteration sequence.

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

For all  $n \geq 1$ ,

$$|p_{n+1} - p_n| = |g(p_n) - g(p_{n-1})| \leq K |p_n - p_{n-1}| \leq K^2 |p_{n-1} - p_{n-2}|$$



# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ . Then by the **fixed point theorem**,  $g$  has a **unique** fixed point  $p$  in  $[a, b]$  and by the **fixed point iteration theorem** for any initial guess  $p_0$  in  $[a, b]$ , the sequence  $p_{n+1} = g(p_n)$  converges to  $p$  and either of the following two inequalities can be used to estimate the error in calculation of the fixed point using the function iteration sequence.

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

For all  $n \geq 1$ ,

$$|p_{n+1} - p_n| \leq K |p_n - p_{n-1}| \leq K^2 |p_{n-1} - p_{n-2}| \dots \leq K^n |p_1 - p_0|$$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

For all  $n \geq 1$ ,  $|p_{n+1} - p_n| \leq K^n |p_1 - p_0|$ .

For all  $m > n \geq 1$

$$|p_m - p_n| = |p_m - p_{m-1} + p_{m-1} - p_{m-2} + \dots + p_{n+1} - p_n|$$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

For all  $n \geq 1$ ,  $|p_{n+1} - p_n| \leq K^n |p_1 - p_0|$ .

For all  $m > n \geq 1$

$$\begin{aligned} |p_m - p_n| &= |p_m - p_{m-1} + p_{m-1} - p_{m-2} + \dots + p_{n+1} - p_n| \\ &\leq |p_m - p_{m-1}| + |p_{m-1} - p_{m-2}| + \dots + |p_{n+1} - p_n| \\ &\text{(Triangle Inequality)} \end{aligned}$$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

For all  $n \geq 1$ ,  $|p_{n+1} - p_n| \leq K^n |p_1 - p_0|$ .

For all  $m > n \geq 1$

$$\begin{aligned} |p_m - p_n| &= |p_m - p_{m-1} + p_{m-1} - p_{m-2} + \dots + p_{n+1} - p_n| \\ &\leq |p_m - p_{m-1}| + |p_{m-1} - p_{m-2}| + \dots + |p_{n+1} - p_n| \\ &\leq K^{m-1} |p_1 - p_0| + K^{m-2} |p_1 - p_0| + \dots + K^n |p_1 - p_0| \end{aligned}$$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

For all  $m > n \geq 1$

$$\begin{aligned} |p_m - p_n| &= |p_m - p_{m-1} + p_{m-1} - p_{m-2} + \dots + p_{n+1} - p_n| \\ &\leq |p_m - p_{m-1}| + |p_{m-1} - p_{m-2}| + \dots + |p_{n+1} - p_n| \\ &\leq K^{m-1} |p_1 - p_0| + K^{m-2} |p_1 - p_0| + \dots + K^n |p_1 - p_0| \\ &= (K^{m-1} + K^{m-2} + \dots + K^n) |p_1 - p_0| \end{aligned}$$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

For all  $m > n \geq 1$

$$\begin{aligned} |p_m - p_n| &= |p_m - p_{m-1} + p_{m-1} - p_{m-2} + \dots + p_{n+1} - p_n| \\ &\leq |p_m - p_{m-1}| + |p_{m-1} - p_{m-2}| + \dots + |p_{n+1} - p_n| \\ &\leq K^{m-1} |p_1 - p_0| + K^{m-2} |p_1 - p_0| + \dots + K^n |p_1 - p_0| \\ &= (K^{m-1} + K^{m-2} + \dots + K^n) |p_1 - p_0| \leq (\sum_{i=n}^{\infty} K^i) |p_1 - p_0| \end{aligned}$$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

For all  $m > n \geq 1$

$$\begin{aligned} |p_m - p_n| &= |p_m - p_{m-1} + p_{m-1} - p_{m-2} + \dots + p_{n+1} - p_n| \\ &\leq |p_m - p_{m-1}| + |p_{m-1} - p_{m-2}| + \dots + |p_{n+1} - p_n| \\ &\leq K^{m-1} |p_1 - p_0| + K^{m-2} |p_1 - p_0| + \dots + K^n |p_1 - p_0| \\ &= (K^{m-1} + K^{m-2} + \dots + K^n) |p_1 - p_0| \leq (\sum_{i=n}^{\infty} K^i) |p_1 - p_0| = \frac{K^n}{1-K} |p_1 - p_0| \\ \sum_{i=0}^{\infty} \alpha r^i &= \frac{\alpha}{1-r} \text{ when } |r| < 1. \end{aligned}$$

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

For all  $m > n \geq 1$   $|p_m - p_n| \leq \frac{K^n}{1-K} |p_1 - p_0|$  and  $\lim_{m \rightarrow \infty} p_m = p$  implies  $|p - p_n| = |p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$ .



# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

The second inequality is more useful when the original interval  $[a, b]$  has a large length and when we might have already started close to the actual fixed point.

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

The second inequality is more useful when the original interval  $[a, b]$  has a large length and when we might have already started close to the actual fixed point. The second inequality can also be used with replacing the initial guess with a number in the sequence that is already close to the fixed point.

# Error Estimate for Fixed-Point Iteration

Suppose a function  $g$  satisfies the conditions of the fixed point iteration theorem on the interval  $[a, b]$ .

- For all  $n \geq 1$ ,  $|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$
- For all  $n \geq 1$ ,  $|p_n - p| \leq \frac{K^n}{1-K} |p_1 - p_0|$

where  $K$  is a **positive constant strictly less than 1** such that  $|g'(x)| \leq K$  for all  $x$  in  $[a, b]$ .

The second inequality is more useful when the original interval  $[a, b]$  has a large length and when we might have already started close to the actual fixed point. The second inequality can also be used with replacing the initial guess with a number in the sequence that is already close to the fixed point. By restarting the iteration at such number, we can have a better estimate on how many more iterations are needed to improve the accuracy of our estimate of the fixed point.

# Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

# Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Previously we saw that  $g$  has a unique fixed point  $p$  in  $[1, 8]$  and the sequence of iterations  $g(p_{n+1}) = g(p_n)$  converges to  $p$  for any choice of  $p_0$  in  $[1, 8]$ .

# Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Previously we saw that  $g$  has a unique fixed point  $p$  in  $[1, 8]$  and the sequence of iterations  $g(p_{n+1}) = g(p_n)$  converges to  $p$  for any choice of  $p_0$  in  $[1, 8]$ . We did this by showing  $g(x)$  is continuous on  $[1, 8]$ , differentiable on  $(1, 8)$ ,  $g([1, 8]) \subset [1, 8]$  and  $|g'(x)| \leq K = \frac{1}{2} < 1$  for all  $x \in [1, 8]$ .

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Previously we saw that  $g$  has a unique fixed point  $p$  in  $[1, 8]$  and the sequence of iterations  $g(p_{n+1}) = g(p_n)$  converges to  $p$  for any choice of  $p_0$  in  $[1, 8]$ . We did this by showing  $g(x)$  is continuous on  $[1, 8]$ , differentiable on  $(1, 8)$ ,  $g([1, 8]) \subset [1, 8]$  and  $|g'(x)| \leq K = \frac{1}{2} < 1$  for all  $x \in [1, 8]$ .

Using the first estimate we get

$$|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\}$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Previously we saw that  $g$  has a unique fixed point  $p$  in  $[1, 8]$  and the sequence of iterations  $g(p_{n+1}) = g(p_n)$  converges to  $p$  for any choice of  $p_0$  in  $[1, 8]$ . We did this by showing  $g(x)$  is continuous on  $[1, 8]$ , differentiable on  $(1, 8)$ ,  $g([1, 8]) \subset [1, 8]$  and  $|g'(x)| \leq K = \frac{1}{2} < 1$  for all  $x \in [1, 8]$ .

Using the first estimate we get

$$|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\} = \left(\frac{1}{2}\right)^n \max\{3 - 1, 8 - 3\}$$



# Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Previously we saw that  $g$  has a unique fixed point  $p$  in  $[1, 8]$  and the sequence of iterations  $g(p_{n+1}) = g(p_n)$  converges to  $p$  for any choice of  $p_0$  in  $[1, 8]$ . We did this by showing  $g(x)$  is continuous on  $[1, 8]$ , differentiable on  $(1, 8)$ ,  $g([1, 8]) \subset [1, 8]$  and  $|g'(x)| \leq K = \frac{1}{2} < 1$  for all  $x \in [1, 8]$ .

Using the first estimate we get

$$|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\} = \left(\frac{1}{2}\right)^n \max\{3 - 1, 8 - 3\} = \frac{5}{2^n}$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Previously we saw that  $g$  has a unique fixed point  $p$  in  $[1, 8]$  and the sequence of iterations  $g(p_{n+1}) = g(p_n)$  converges to  $p$  for any choice of  $p_0$  in  $[1, 8]$ . We did this by showing  $g(x)$  is continuous on  $[1, 8]$ , differentiable on  $(1, 8)$ ,  $g([1, 8]) \subset [1, 8]$  and  $|g'(x)| \leq K = \frac{1}{2} < 1$  for all  $x \in [1, 8]$ .

Using the first estimate we get

$$|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\} = \left(\frac{1}{2}\right)^n \max\{3 - 1, 8 - 3\} = \frac{5}{2^n}$$

So to make sure  $|p_n - p| < 0.0001$ , we can choose  $n$  large enough so that

$$\frac{5}{2^n} < 0.0001$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the first estimate we get

$$|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\} = \left(\frac{1}{2}\right)^n \max\{3 - 1, 8 - 3\} = \frac{5}{2^n}$$

So to make sure  $|p_n - p| < 0.0001$ , we can choose  $n$  large enough so that

$$\frac{5}{2^n} < 0.0001 \Leftrightarrow 50000 < 2^n$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the first estimate we get

$$|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\} = \left(\frac{1}{2}\right)^n \max\{3 - 1, 8 - 3\} = \frac{5}{2^n}$$

So to make sure  $|p_n - p| < 0.0001$ , we can choose  $n$  large enough so that

$$\frac{5}{2^n} < 0.0001 \Leftrightarrow 50000 < 2^n \Leftrightarrow \ln(50000) < n \ln(2)$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the first estimate we get

$$|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\} = \left(\frac{1}{2}\right)^n \max\{3 - 1, 8 - 3\} = \frac{5}{2^n}$$

So to make sure  $|p_n - p| < 0.0001$ , we can choose  $n$  large enough so that

$$\frac{5}{2^n} < 0.0001 \Leftrightarrow \ln(50000) < n \ln(2) \Leftrightarrow n > \frac{\ln(50000)}{\ln(2)} \approx 15.610$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the first estimate we get

$$|p_n - p| \leq K^n \max\{p_0 - a, b - p_0\} = \left(\frac{1}{2}\right)^n \max\{3 - 1, 8 - 3\} = \frac{5}{2^n}$$

So to make sure  $|p_n - p| < 0.0001$ , we can choose  $n$  large enough so that

$$\frac{5}{2^n} < 0.0001 \Leftrightarrow \ln(50000) < n \ln(2) \Leftrightarrow n > \frac{\ln(50000)}{\ln(2)} \approx 15.610$$

Therefore, using the first estimate, we conclude that  $p_{16}$  will be within 0.0001 of the fixed point  $p$ .

# Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the second estimate we get

$$|p_n - p| \leq \frac{K^n}{1 - K} |p_1 - p_0|$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the second estimate we get

$$|p_n - p| \leq \frac{K^n}{1 - K} |p_1 - p_0| = \frac{(\frac{1}{2})^n}{1 - \frac{1}{2}} \left| 3 - \frac{11}{6} \right|$$

$$p_1 = g(p_0) = g(3) = \frac{11}{6}$$



## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the second estimate we get

$$|p_n - p| \leq \frac{K^n}{1 - K} |p_1 - p_0| = \frac{(\frac{1}{2})^n}{1 - \frac{1}{2}} |3 - \frac{11}{6}| = \frac{7}{2^n \times 3}$$

So to make sure  $|p_n - p| < 0.0001$ , we can choose  $n$  large enough so that

$$\frac{7}{2^n \times 3} < 0.0001$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the second estimate we get

$$|p_n - p| \leq \frac{K^n}{1 - K} |p_1 - p_0| = \frac{(\frac{1}{2})^n}{1 - \frac{1}{2}} |3 - \frac{11}{6}|$$

So to make sure  $|p_n - p| < 0.0001$ , we can choose  $n$  large enough so that

$$\frac{7}{2^n \times 3} < 0.0001 \Leftrightarrow \frac{70000}{3} < 2^n$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the second estimate we get

$$|p_n - p| \leq \frac{K^n}{1 - K} |p_1 - p_0| = \frac{(\frac{1}{2})^n}{1 - \frac{1}{2}} |3 - \frac{11}{6}|$$

So to make sure  $|p_n - p| < 0.0001$ , we can choose  $n$  large enough so that

$$\frac{7}{2^n \times 3} < 0.0001 \Leftrightarrow \frac{70000}{3} < 2^n \Leftrightarrow \ln\left(\frac{70000}{3}\right) < n \ln(2)$$

## Example

For  $g(x) = \frac{x^2+2}{2x}$  on the interval  $[1, 8]$ , if we start with  $p_0 = 3$ , how large should  $n$  be so that the  $n$ th iterate  $p_n$  is within 0.0001 of the fixed point  $p$  of  $g(x)$ ?

Using the second estimate we get

$$|p_n - p| \leq \frac{K^n}{1 - K} |p_1 - p_0| = \frac{(\frac{1}{2})^n}{1 - \frac{1}{2}} |3 - \frac{11}{6}|$$

So to make sure  $|p_n - p| < 0.0001$ , we can choose  $n$  large enough so that

$$\frac{7}{2^n \times 3} < 0.0001 \Leftrightarrow \ln\left(\frac{70000}{3}\right) < n \ln(2) \Leftrightarrow n > \frac{\ln\left(\frac{70000}{3}\right)}{\ln(2)} \approx 14.511$$

Therefore, using the second estimate, we conclude that  $p_{15}$  will be within 0.0001 of the fixed point  $p$ .