

(3.1) Interpolation and Polynomial Approximation

MATH 4701 Numerical Analysis

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication.

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated.

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated. In theory, there are various ways polynomial approximation of functions can be achieved.

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated. In theory, there are various ways polynomial approximation of functions can be achieved. These different approaches apply to more general or more restrictive types of functions and they may be easier or harder to implement computationally.

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated. In theory, there are various ways polynomial approximation of functions can be achieved. These different approaches apply to more general or more restrictive types of functions and they may be easier or harder to implement computationally.

Weierstrass Approximation Theorem: Given any function f **continuous** on the interval $[a, b]$, and any $\epsilon > 0$, there is a polynomial $P(x)$ such that $|f(x) - P(x)| < \epsilon$ for all x in $[a, b]$.

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated. In theory, there are various ways polynomial approximation of functions can be achieved. These different approaches apply to more general or more restrictive types of functions and they may be easier or harder to implement computationally.

Weierstrass Approximation Theorem: Given any function f **continuous** on the interval $[a, b]$, and any $\epsilon > 0$, there is a polynomial $P(x)$ such that $|f(x) - P(x)| < \epsilon$ for all x in $[a, b]$.

This theorem covers a wide range of functions (continuous on $[a, b]$) and has an effective error bound.

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated. In theory, there are various ways polynomial approximation of functions can be achieved. These different approaches apply to more general or more restrictive types of functions and they may be easier or harder to implement computationally.

Weierstrass Approximation Theorem: Given any function f **continuous** on the interval $[a, b]$, and any $\epsilon > 0$, there is a polynomial $P(x)$ such that $|f(x) - P(x)| < \epsilon$ for all x in $[a, b]$.

This theorem covers a wide range of functions (continuous on $[a, b]$) and has an effective error bound. But it does not provide any computational way to find the polynomial $P(x)$ and it does not have any bounds on the degree of $P(x)$.

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated. In theory, there are various ways polynomial approximation of functions can be achieved. These different approaches apply to more general or more restrictive types of functions and they may be easier or harder to implement computationally.

Taylor's Theorem: Suppose

- $f \in C^n[a, b]$

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated. In theory, there are various ways polynomial approximation of functions can be achieved. These different approaches apply to more general or more restrictive types of functions and they may be easier or harder to implement computationally.

Taylor's Theorem: Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$ (the $n + 1$ th derivative of f) exists on $[a, b]$, and

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated. In theory, there are various ways polynomial approximation of functions can be achieved. These different approaches apply to more general or more restrictive types of functions and they may be easier or harder to implement computationally.

Taylor's Theorem: Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$ (the $n + 1$ th derivative of f) exists on $[a, b]$, and
- $x_0 \in [a, b]$

Polynomial Approximation of Functions

Approximation of functions by polynomials has computational appeals since in evaluation of a polynomial one has to only perform summation and multiplication. Also derivatives and integrals of polynomials are most easily calculated. In theory, there are various ways polynomial approximation of functions can be achieved. These different approaches apply to more general or more restrictive types of functions and they may be easier or harder to implement computationally.

Taylor's Theorem: Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$ (the $n + 1$ th derivative of f) exists on $[a, b]$, and
- $x_0 \in [a, b]$

Let
$$P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$$

Polynomial Approximation of Functions

Taylor's Theorem: Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$ (the $n + 1$ th derivative of f) exists on $[a, b]$, and
- $x_0 \in [a, b]$

Let $P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$

Then for every $x \in [a, b]$, there is a number $\zeta = \zeta(x)$, between x_0 and x , such that

$$f(x) - P_n(x) = \frac{f^{(n+1)}(\zeta)(x - x_0)^{n+1}}{(n + 1)!}$$

Polynomial Approximation of Functions

Taylor's Theorem: Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$ (the $n + 1$ th derivative of f) exists on $[a, b]$, and
- $x_0 \in [a, b]$

Let $P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$

Then for every $x \in [a, b]$, there is a number $\zeta = \zeta(x)$, between x_0 and x , such that

$$f(x) - P_n(x) = \frac{f^{(n+1)}(\zeta)(x - x_0)^{n+1}}{(n + 1)!}$$

Taylor's polynomial approximation, requires the values $f(x_0)$, $f'(x_0)$, ..., $f^{(n)}(x_0)$ to exist in order to be defined.

Polynomial Approximation of Functions

Taylor's Theorem: Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$ (the $n + 1$ th derivative of f) exists on $[a, b]$, and
- $x_0 \in [a, b]$

Let $P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$
Then for every $x \in [a, b]$, there is a number $\zeta = \zeta(x)$, between x_0 and x , such that

$$f(x) - P_n(x) = \frac{f^{(n+1)}(\zeta)(x - x_0)^{n+1}}{(n + 1)!}$$

Taylor's polynomial approximation, requires the values $f(x_0)$, $f'(x_0)$, ..., $f^{(n)}(x_0)$ to exist in order to be defined. It also requires $f \in C^n[a, b]$ and $f^{(n+1)}$ to exist on $[a, b]$ to have an effective error bound.

Polynomial Approximation of Functions

Taylor's Theorem: Suppose

- $f \in C^n[a, b]$
- $f^{(n+1)}$ (the $n + 1$ th derivative of f) exists on $[a, b]$, and
- $x_0 \in [a, b]$

Let $P_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$
Then for every $x \in [a, b]$, there is a number $\zeta = \zeta(x)$, between x_0 and x , such that

$$f(x) - P_n(x) = \frac{f^{(n+1)}(\zeta)(x - x_0)^{n+1}}{(n + 1)!}$$

Taylor's polynomial approximation, requires the values $f(x_0)$, $f'(x_0)$, ..., $f^{(n)}(x_0)$ to exist in order to be defined. It also requires $f \in C^n[a, b]$ and $f^{(n+1)}$ to exist on $[a, b]$ to have an effective error bound. It gives an explicit way of constructing polynomial $P_n(x)$ which only depends on the values f and its higher derivatives at $x = x_0$.

Polynomial Approximation of Functions

Lagrange interpolation polynomial is a polynomial $P(x)$ that

- can be constructed for any function

Polynomial Approximation of Functions

Lagrange interpolation polynomial is a polynomial $P(x)$ that

- can be constructed for any function
- is only based on the values of the function at $n + 1$ different points

Polynomial Approximation of Functions

Lagrange interpolation polynomial is a polynomial $P(x)$ that

- can be constructed for any function
- is only based on the values of the function at $n + 1$ different points
- agrees with the given function at these $n + 1$ points

Polynomial Approximation of Functions

Lagrange interpolation polynomial is a polynomial $P(x)$ that

- can be constructed for any function
- is only based on the values of the function at $n + 1$ different points
- agrees with the given function at these $n + 1$ points
- has degree n

Polynomial Approximation of Functions

Lagrange interpolation polynomial is a polynomial $P(x)$ that

- can be constructed for any function
- is only based on the values of the function at $n + 1$ different points
- agrees with the given function at these $n + 1$ points
- has degree n
- if $f \in C^{n+1}[a, b]$ then there are effective error bounds controlling how far $P(x)$ can be from f on the interval $[a, b]$.

Polynomial Approximation of Functions

Lagrange interpolation polynomial is a polynomial $P(x)$ that

- can be constructed for any function
- is only based on the values of the function at $n + 1$ different points
- agrees with the given function at these $n + 1$ points
- has degree n
- if $f \in C^{n+1}[a, b]$ then there are effective error bounds controlling how far $P(x)$ can be from f on the interval $[a, b]$.

Since the construction of interpolation polynomial only depends on the values of the function at finitely many points, one does not need an explicit formula for description of the function to set up the corresponding interpolation polynomial.

Polynomial Approximation of Functions

Lagrange interpolation polynomial is a polynomial $P(x)$ that

- can be constructed for any function
- is only based on the values of the function at $n + 1$ different points
- agrees with the given function at these $n + 1$ points
- has degree n
- if $f \in C^{n+1}[a, b]$ then there are effective error bounds controlling how far $P(x)$ can be from f on the interval $[a, b]$.

Since the construction of interpolation polynomial only depends on the values of the function at finitely many points, one does not need an explicit formula for description of the function to set up the corresponding interpolation polynomial. One can determine the approximating polynomial using a table of values for the function or a list consisting of data evaluations at certain sample points.

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We first introduce $n + 1$ polynomials of degree n $L_k(x)$, $0 \leq k \leq n$ such that

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We first introduce $n + 1$ polynomials of degree n $L_k(x)$, $0 \leq k \leq n$ such that

$$(a) \quad L_k(x_k) = 1$$

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We first introduce $n + 1$ polynomials of degree n $L_k(x)$, $0 \leq k \leq n$ such that

(a) $L_k(x_k) = 1$

(b) $L_k(x_j) = 0$ for all $j \neq k$, $0 \leq j \leq n$

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We first introduce $n + 1$ polynomials of degree n $L_k(x)$, $0 \leq k \leq n$ such that

(a) $L_k(x_k) = 1$

(b) $L_k(x_j) = 0$ for all $j \neq k$, $0 \leq j \leq n$

$$(x - x_0)(x - x_1) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)$$

satisfies (b) and has degree n .

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We first introduce $n + 1$ polynomials of degree n $L_k(x)$, $0 \leq k \leq n$ such that

$$(a) \quad L_k(x_k) = 1$$

$$(b) \quad L_k(x_j) = 0 \text{ for all } j \neq k, 0 \leq j \leq n$$

$$\prod_{\substack{0 \leq j \leq n \\ j \neq k}} (x - x_j) = (x - x_0)(x - x_1) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)$$

satisfies (b) and has degree n .

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We first introduce $n + 1$ polynomials of degree n $L_k(x)$, $0 \leq k \leq n$ such that

(a) $L_k(x_k) = 1$

(b) $L_k(x_j) = 0$ for all $j \neq k$, $0 \leq j \leq n$

$$\frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$$

satisfies both (a) and (b) and has degree n .

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We first introduce $n + 1$ polynomials of degree n $L_k(x)$, $0 \leq k \leq n$ such that

(a) $L_k(x_k) = 1$

(b) $L_k(x_j) = 0$ for all $j \neq k$, $0 \leq j \leq n$

$$\prod_{\substack{0 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)} = \frac{(x - x_0) \dots (x - x_{k-1})(x - x_{k+1}) \dots (x - x_n)}{(x_k - x_0) \dots (x_k - x_{k-1})(x_k - x_{k+1}) \dots (x_k - x_n)}$$

satisfies both (a) and (b) and has degree n .

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We first introduce $n + 1$ polynomials of degree n $L_k(x)$, $0 \leq k \leq n$ such that

(a) $L_k(x_k) = 1$

(b) $L_k(x_j) = 0$ for all $j \neq k$, $0 \leq j \leq n$

$$L_k(x) = \prod_{\substack{0 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$$

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We first introduce $n + 1$ polynomials of degree n $L_k(x)$, $0 \leq k \leq n$ such that

(a) $L_k(x_k) = 1$

(b) $L_k(x_j) = 0$ for all $j \neq k$, $0 \leq j \leq n$

$$L_k(x) = \prod_{\substack{0 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$$

Then $P(x) = y_0 L_0(x) + y_1 L_1(x) + \dots + y_n L_n(x)$ satisfies $P(x_i) = y_i$ for all $0 \leq i \leq n$

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

$$L_k(x) = \prod_{\substack{0 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$$

Then $P(x) = y_0L_0(x) + y_1L_1(x) + \dots + y_nL_n(x)$ satisfies $P(x_i) = y_i$ for all $0 \leq i \leq n$ and it has degree at most n .

Construction of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values (not necessarily distinct) y_0, y_1, \dots, y_n we construct a polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

$$L_k(x) = \prod_{\substack{0 \leq j \leq n \\ j \neq k}} \frac{(x - x_j)}{(x_k - x_j)}$$

Then $P(x) = y_0L_0(x) + y_1L_1(x) + \dots + y_nL_n(x)$ satisfies $P(x_i) = y_i$ for all $0 \leq i \leq n$ and it has degree at most n .

$$P(x) = \sum_{0 \leq k \leq n} y_k L_k(x)$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)} = -\frac{(x-1)(x-2)(x+2)(x-3)}{24}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)} = -\frac{(x-1)(x-2)(x+2)(x-3)}{24}$$

$$L_2(x) = \frac{(x-1)(x+1)(x+2)(x-3)}{(2-1)(2+1)(2+2)(2-3)}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)} = -\frac{(x-1)(x-2)(x+2)(x-3)}{24}$$

$$L_2(x) = \frac{(x-1)(x+1)(x+2)(x-3)}{(2-1)(2+1)(2+2)(2-3)} = -\frac{(x-1)(x+1)(x+2)(x-3)}{12}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)} = -\frac{(x-1)(x-2)(x+2)(x-3)}{24}$$

$$L_2(x) = \frac{(x-1)(x+1)(x+2)(x-3)}{(2-1)(2+1)(2+2)(2-3)} = -\frac{(x-1)(x+1)(x+2)(x-3)}{12}$$

$$L_3(x) = \frac{(x-1)(x+1)(x-2)(x-3)}{(-2-1)(-2+1)(-2-2)(-2-3)}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)} = -\frac{(x-1)(x-2)(x+2)(x-3)}{24}$$

$$L_2(x) = \frac{(x-1)(x+1)(x+2)(x-3)}{(2-1)(2+1)(2+2)(2-3)} = -\frac{(x-1)(x+1)(x+2)(x-3)}{12}$$

$$L_3(x) = \frac{(x-1)(x+1)(x-2)(x-3)}{(-2-1)(-2+1)(-2-2)(-2-3)} = \frac{(x-1)(x+1)(x-2)(x-3)}{60}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)} = -\frac{(x-1)(x-2)(x+2)(x-3)}{24}$$

$$L_2(x) = \frac{(x-1)(x+1)(x+2)(x-3)}{(2-1)(2+1)(2+2)(2-3)} = -\frac{(x-1)(x+1)(x+2)(x-3)}{12}$$

$$L_3(x) = \frac{(x-1)(x+1)(x-2)(x-3)}{(-2-1)(-2+1)(-2-2)(-2-3)} = \frac{(x-1)(x+1)(x-2)(x-3)}{60}$$

$$L_4(x) = \frac{(x-1)(x+1)(x-2)(x+2)}{(3-1)(3+1)(3-2)(3+2)}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)} = -\frac{(x-1)(x-2)(x+2)(x-3)}{24}$$

$$L_2(x) = \frac{(x-1)(x+1)(x+2)(x-3)}{(2-1)(2+1)(2+2)(2-3)} = -\frac{(x-1)(x+1)(x+2)(x-3)}{12}$$

$$L_3(x) = \frac{(x-1)(x+1)(x-2)(x-3)}{(-2-1)(-2+1)(-2-2)(-2-3)} = \frac{(x-1)(x+1)(x-2)(x-3)}{60}$$

$$L_4(x) = \frac{(x-1)(x+1)(x-2)(x+2)}{(3-1)(3+1)(3-2)(3+2)} = \frac{(x-1)(x+1)(x-2)(x+2)}{40}$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)} = -\frac{(x-1)(x-2)(x+2)(x-3)}{24}$$

$$L_2(x) = \frac{(x-1)(x+1)(x+2)(x-3)}{(2-1)(2+1)(2+2)(2-3)} = -\frac{(x-1)(x+1)(x+2)(x-3)}{12}$$

$$L_3(x) = \frac{(x-1)(x+1)(x-2)(x-3)}{(-2-1)(-2+1)(-2-2)(-2-3)} = \frac{(x-1)(x+1)(x-2)(x-3)}{60}$$

$$L_4(x) = \frac{(x-1)(x+1)(x-2)(x+2)}{(3-1)(3+1)(3-2)(3+2)} = \frac{(x-1)(x+1)(x-2)(x+2)}{40}$$

$$P(x) = 3L_0(x) + 5L_1(x) + 0L_2(x) - L_3(x) - 6L_4(x)$$

Example

Find a polynomial $P(x)$ of degree at most four such that $P(1) = 3$, $P(-1) = 5$, $P(2) = 0$, $P(-2) = -1$, and $P(3) = -6$.

We let $x_0 = 1$, $x_1 = -1$, $x_2 = 2$, $x_3 = -2$, and $x_4 = 3$.

$$L_0(x) = \frac{(x+1)(x-2)(x+2)(x-3)}{(1+1)(1-2)(1+2)(1-3)} = \frac{(x+1)(x-2)(x+2)(x-3)}{12}$$

$$L_1(x) = \frac{(x-1)(x-2)(x+2)(x-3)}{(-1-1)(-1-2)(-1+2)(-1-3)} = -\frac{(x-1)(x-2)(x+2)(x-3)}{24}$$

$$L_2(x) = \frac{(x-1)(x+1)(x+2)(x-3)}{(2-1)(2+1)(2+2)(2-3)} = -\frac{(x-1)(x+1)(x+2)(x-3)}{12}$$

$$L_3(x) = \frac{(x-1)(x+1)(x-2)(x-3)}{(-2-1)(-2+1)(-2-2)(-2-3)} = \frac{(x-1)(x+1)(x-2)(x-3)}{60}$$

$$L_4(x) = \frac{(x-1)(x+1)(x-2)(x+2)}{(3-1)(3+1)(3-2)(3+2)} = \frac{(x-1)(x+1)(x-2)(x+2)}{40}$$

$$\begin{aligned} P(x) &= 3L_0(x) + 5L_1(x) + 0L_2(x) - L_3(x) - 6L_4(x) \\ &= -\frac{x^4}{8} + \frac{5x^3}{12} - \frac{7x^2}{8} - \frac{17x}{12} + 5 \end{aligned}$$

Uniqueness of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values y_0, y_1, \dots, y_n there is a **unique** polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

Uniqueness of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values y_0, y_1, \dots, y_n there is a **unique** polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We just found a polynomial of degree at most n satisfying $P(x_i) = y_i$ for all $0 \leq i \leq n$.

Uniqueness of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values y_0, y_1, \dots, y_n there is a **unique** polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We just found a polynomial of degree at most n satisfying $P(x_i) = y_i$ for all $0 \leq i \leq n$. If $Q(x)$ is another polynomial of degree at most n satisfying $Q(x_i) = y_i$ for all $0 \leq i \leq n$,

Uniqueness of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values y_0, y_1, \dots, y_n there is a **unique** polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We just found a polynomial of degree at most n satisfying $P(x_i) = y_i$ for all $0 \leq i \leq n$. If $Q(x)$ is another polynomial of degree at most n satisfying $Q(x_i) = y_i$ for all $0 \leq i \leq n$, then $R(x) = (P - Q)(x)$ is a polynomial of degree at most n satisfying $R(x_i) = 0$ for all $0 \leq i \leq n$.

Uniqueness of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values y_0, y_1, \dots, y_n there is a **unique** polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We just found a polynomial of degree at most n satisfying $P(x_i) = y_i$ for all $0 \leq i \leq n$. If $Q(x)$ is another polynomial of degree at most n satisfying $Q(x_i) = y_i$ for all $0 \leq i \leq n$, then $R(x) = (P - Q)(x)$ is a polynomial of degree at most n satisfying $R(x_i) = 0$ for all $0 \leq i \leq n$. A non-zero polynomial of degree at most n cannot have $n + 1$ distinct roots.

Uniqueness of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values y_0, y_1, \dots, y_n there is a **unique** polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We just found a polynomial of degree at most n satisfying $P(x_i) = y_i$ for all $0 \leq i \leq n$. If $Q(x)$ is another polynomial of degree at most n satisfying $Q(x_i) = y_i$ for all $0 \leq i \leq n$, then $R(x) = (P - Q)(x)$ is a polynomial of degree at most n satisfying $R(x_i) = 0$ for all $0 \leq i \leq n$. A non-zero polynomial of degree at most n cannot have $n + 1$ distinct roots. Therefore, $R(x) = 0$ and $P(x) = Q(x)$.

Uniqueness of Lagrange Interpolating Polynomials

For distinct values x_0, x_1, \dots, x_n , and for the corresponding values y_0, y_1, \dots, y_n there is a **unique** polynomial $P(x)$ of degree at most n such that

$$P(x_i) = y_i \quad \text{for all } 0 \leq i \leq n$$

We just found a polynomial of degree at most n satisfying $P(x_i) = y_i$ for all $0 \leq i \leq n$. If $Q(x)$ is another polynomial of degree at most n satisfying $Q(x_i) = y_i$ for all $0 \leq i \leq n$, then $R(x) = (P - Q)(x)$ is a polynomial of degree at most n satisfying $R(x_i) = 0$ for all $0 \leq i \leq n$. A non-zero polynomial of degree at most n cannot have $n + 1$ distinct roots. Therefore, $R(x) = 0$ and $P(x) = Q(x)$. Hence, $P(x)$ is the only polynomial of degree at most n satisfying $P(x_i) = y_i$ for all $0 \leq i \leq n$.

Generalized Rolle's Theorem

Rolle's Theorem: If $f \in C[a, b]$ and f is differentiable on (a, b) , and if $f(a) = f(b)$,

Generalized Rolle's Theorem

Rolle's Theorem: If $f \in C[a, b]$ and f is differentiable on (a, b) , and if $f(a) = f(b)$, then there is a number $\alpha \in (a, b)$ such that $f'(\alpha) = 0$.

Generalized Rolle's Theorem

Rolle's Theorem: If $f \in C[a, b]$ and f is differentiable on (a, b) , and if $f(a) = f(b)$, then there is a number $\alpha \in (a, b)$ such that $f'(\alpha) = 0$.

Generalized Rolle's Theorem: Suppose $f \in C[a, b]$ and f is n times differentiable on (a, b) . If $a \leq x_0 < x_1 < \dots < x_n \leq b$ and $f(x_0) = f(x_1) = \dots = f(x_n)$,

Generalized Rolle's Theorem

Rolle's Theorem: If $f \in C[a, b]$ and f is differentiable on (a, b) , and if $f(a) = f(b)$, then there is a number $\alpha \in (a, b)$ such that $f'(\alpha) = 0$.

Generalized Rolle's Theorem: Suppose $f \in C[a, b]$ and f is n times differentiable on (a, b) . If $a \leq x_0 < x_1 < \dots < x_n \leq b$ and $f(x_0) = f(x_1) = \dots = f(x_n)$, then there is a number $\alpha \in (a, b)$ such that $f^{(n)}(\alpha) = 0$.

Generalized Rolle's Theorem

Rolle's Theorem: If $f \in C[a, b]$ and f is differentiable on (a, b) , and if $f(a) = f(b)$, then there is a number $\alpha \in (a, b)$ such that $f'(\alpha) = 0$.

Generalized Rolle's Theorem: Suppose $f \in C[a, b]$ and f is n times differentiable on (a, b) . If $a \leq x_0 < x_1 < \dots < x_n \leq b$ and $f(x_0) = f(x_1) = \dots = f(x_n)$, then there is a number $\alpha \in (a, b)$ such that $f^{(n)}(\alpha) = 0$.

Proof of Generalized Rolle's Theorem: By Rolle's theorem, there are points $x_0 < x_0^1 < x_1 < x_1^1 < x_2 < \dots < x_{n-1} < x_{n-1}^1 < x_n$ such that $f'(x_0^1) = f'(x_1^1) = \dots = f'(x_{n-1}^1) = 0$.

Generalized Rolle's Theorem

Rolle's Theorem: If $f \in C[a, b]$ and f is differentiable on (a, b) , and if $f(a) = f(b)$, then there is a number $\alpha \in (a, b)$ such that $f'(\alpha) = 0$.

Generalized Rolle's Theorem: Suppose $f \in C[a, b]$ and f is n times differentiable on (a, b) . If $a \leq x_0 < x_1 < \dots < x_n \leq b$ and $f(x_0) = f(x_1) = \dots = f(x_n)$, then there is a number $\alpha \in (a, b)$ such that $f^{(n)}(\alpha) = 0$.

Proof of Generalized Rolle's Theorem: By Rolle's theorem, there are points $x_0 < x_0^1 < x_1 < x_1^1 < x_2 < \dots < x_{n-1} < x_{n-1}^1 < x_n$ such that $f'(x_0^1) = f'(x_1^1) = \dots = f'(x_{n-1}^1) = 0$.

Again by Rolle's theorem, there are points $x_0^1 < x_0^2 < x_1^1 < x_1^2 < x_2^1 < \dots < x_{n-2}^1 < x_{n-2}^2 < x_{n-1}^1$ such that $f''(x_0^2) = f''(x_1^2) = \dots = f''(x_{n-2}^2) = 0$.

Generalized Rolle's Theorem

Generalized Rolle's Theorem: Suppose $f \in C[a, b]$ and f is n times differentiable on (a, b) . If $a \leq x_0 < x_1 < \dots < x_n \leq b$ and $f(x_0) = f(x_1) = \dots = f(x_n)$, then there is a number $\alpha \in (a, b)$ such that $f^{(n)}(\alpha) = 0$.

Proof of Generalized Rolle's Theorem: By Rolle's theorem, there are points $x_0 < x_0^1 < x_1 < x_1^1 < x_2 < \dots < x_{n-1} < x_{n-1}^1 < x_n$ such that $f'(x_0^1) = f'(x_1^1) = \dots = f'(x_{n-1}^1) = 0$.

Again by Rolle's theorem, there are points

$x_0^1 < x_0^2 < x_1^1 < x_1^2 < x_2^1 < \dots < x_{n-2}^1 < x_{n-2}^2 < x_{n-1}^1$ such that $f''(x_0^2) = f''(x_1^2) = \dots = f''(x_{n-2}^2) = 0$.

Continuing this way, for each $0 < k \leq n$ we have points $x_0^k, x_1^k, \dots, x_{n-k}^k$ such that $f^{(k)}(x_0^k) = f^{(k)}(x_1^k) = \dots = f^{(k)}(x_{n-k}^k) = 0$.

Generalized Rolle's Theorem

Generalized Rolle's Theorem: Suppose $f \in C[a, b]$ and f is n times differentiable on (a, b) . If $a \leq x_0 < x_1 < \dots < x_n \leq b$ and $f(x_0) = f(x_1) = \dots = f(x_n)$, then there is a number $\alpha \in (a, b)$ such that $f^{(n)}(\alpha) = 0$.

Proof of Generalized Rolle's Theorem: By Rolle's theorem, there are points $x_0 < x_0^1 < x_1 < x_1^1 < x_2 < \dots < x_{n-1} < x_{n-1}^1 < x_n$ such that $f'(x_0^1) = f'(x_1^1) = \dots = f'(x_{n-1}^1) = 0$.

Again by Rolle's theorem, there are points $x_0^1 < x_0^2 < x_1^1 < x_1^2 < x_2^1 < \dots < x_{n-2}^1 < x_{n-2}^2 < x_{n-1}^1$ such that $f''(x_0^2) = f''(x_1^2) = \dots = f''(x_{n-2}^2) = 0$.

Continuing this way, for each $0 < k \leq n$ we have points $x_0^k, x_1^k, \dots, x_{n-k}^k$ such that $f^{(k)}(x_0^k) = f^{(k)}(x_1^k) = \dots = f^{(k)}(x_{n-k}^k) = 0$.

In particular, there is x_0^n such that $f^{(n)}(x_0^n) = 0$.

Generalized Rolle's Theorem

Generalized Rolle's Theorem: Suppose $f \in C[a, b]$ and f is n times differentiable on (a, b) . If $a \leq x_0 < x_1 < \dots < x_n \leq b$ and $f(x_0) = f(x_1) = \dots = f(x_n)$, then there is a number $\alpha \in (a, b)$ such that $f^{(n)}(\alpha) = 0$.

Proof of Generalized Rolle's Theorem: By Rolle's theorem, there are points $x_0 < x_0^1 < x_1 < x_1^1 < x_2 < \dots < x_{n-1} < x_{n-1}^1 < x_n$ such that $f'(x_0^1) = f'(x_1^1) = \dots = f'(x_{n-1}^1) = 0$.

Continuing this way, for each $0 < k \leq n$ we have points $x_0^k, x_1^k, \dots, x_{n-k}^k$ such that $f^{(k)}(x_0^k) = f^{(k)}(x_1^k) = \dots = f^{(k)}(x_{n-k}^k) = 0$.

In particular, there is x_0^n such that $f^{(n)}(x_0^n) = 0$. Additionally, through all these steps all the points x_i^k remain between x_0 and x_n .

Generalized Rolle's Theorem

Generalized Rolle's Theorem: Suppose $f \in C[a, b]$ and f is n times differentiable on (a, b) . If $a \leq x_0 < x_1 < \dots < x_n \leq b$ and $f(x_0) = f(x_1) = \dots = f(x_n)$, then there is a number $\alpha \in (a, b)$ such that $f^{(n)}(\alpha) = 0$.

Proof of Generalized Rolle's Theorem: By Rolle's theorem, there are points $x_0 < x_0^1 < x_1 < x_1^1 < x_2 < \dots < x_{n-1} < x_{n-1}^1 < x_n$ such that $f'(x_0^1) = f'(x_1^1) = \dots = f'(x_{n-1}^1) = 0$.

Continuing this way, for each $0 < k \leq n$ we have points $x_0^k, x_1^k, \dots, x_{n-k}^k$ such that $f^{(k)}(x_0^k) = f^{(k)}(x_1^k) = \dots = f^{(k)}(x_{n-k}^k) = 0$.

In particular, there is x_0^n such that $f^{(n)}(x_0^n) = 0$. Additionally, through all these steps all the points x_i^k remain between x_0 and x_n . Therefore, there is a point $\alpha = x_0^n$ such that $f^{(n)}(\alpha) = 0$.

Error Bound for Lagrange Interpolating Polynomials

Notation: Suppose $x_0, x_1, x_2, \dots, x_n$ are distinct numbers. We denote by $[x_0, x_1, \dots, x_n]$ the smallest closed interval that contains all these numbers.

Error Bound for Lagrange Interpolating Polynomials

Notation: Suppose $x_0, x_1, x_2, \dots, x_n$ are distinct numbers. We denote by $[x_0, x_1, \dots, x_n]$ the smallest closed interval that contains all these numbers. In other words, if $c = \min\{x_0, x_1, \dots, x_n\}$ and $d = \max\{x_0, x_1, \dots, x_n\}$ then $[x_0, x_1, \dots, x_n] = [c, d]$.

Error Bound for Lagrange Interpolating Polynomials

Notation: Suppose $x_0, x_1, x_2, \dots, x_n$ are distinct numbers. We denote by $[x_0, x_1, \dots, x_n]$ the smallest closed interval that contains all these numbers. In other words, if $c = \min\{x_0, x_1, \dots, x_n\}$ and $d = \max\{x_0, x_1, \dots, x_n\}$ then $[x_0, x_1, \dots, x_n] = [c, d]$.

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$.

Error Bound for Lagrange Interpolating Polynomials

Notation: Suppose $x_0, x_1, x_2, \dots, x_n$ are distinct numbers. We denote by $[x_0, x_1, \dots, x_n]$ the smallest closed interval that contains all these numbers. In other words, if $c = \min\{x_0, x_1, \dots, x_n\}$ and $d = \max\{x_0, x_1, \dots, x_n\}$ then $[x_0, x_1, \dots, x_n] = [c, d]$.

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Error Bound for Lagrange Interpolating Polynomials

Notation: Suppose $x_0, x_1, x_2, \dots, x_n$ are distinct numbers. We denote by $[x_0, x_1, \dots, x_n]$ the smallest closed interval that contains all these numbers. In other words, if $c = \min\{x_0, x_1, \dots, x_n\}$ and $d = \max\{x_0, x_1, \dots, x_n\}$ then $[x_0, x_1, \dots, x_n] = [c, d]$.

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: If $x = x_k$ for any $0 \leq k \leq n$ then $f(x_k) = P(x_k)$ and both sides of the above equation are equal to zero.

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: If $x = x_k$ for any $0 \leq k \leq n$ then $f(x_k) = P(x_k)$ and both sides of the above equation are equal to zero. Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: If $x = x_k$ for any $0 \leq k \leq n$ then $f(x_k) = P(x_k)$ and both sides of the above equation are equal to zero. Now if x is distinct from all x_k ,

$0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

Notice that in this expression x , $f(x)$, and $P(x)$ are all constants and the only variable is t .

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: If $x = x_k$ for any $0 \leq k \leq n$ then $f(x_k) = P(x_k)$ and both sides of the above equation are equal to zero. Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists and the rest of the expression for $g(t)$ is a polynomial in t .

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: If $x = x_k$ for any $0 \leq k \leq n$ then $f(x_k) = P(x_k)$ and both sides of the above equation are equal to zero. Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists and the rest of the expression for $g(t)$ is a polynomial in t . Therefore, $g \in C^n[a, b]$ and $g^{(n+1)}(x)$ exists.

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists and the rest of the expression for $g(t)$ is a polynomial in t . Therefore, $g \in C^n[a, b]$ and $g^{(n+1)}(x)$ exists. For each $0 \leq k \leq n$, $g(x_k) = f(x_k) - P(x_k) - [f(x) - P(x)]0 = 0$.

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists and the rest of the expression for $g(t)$ is a polynomial in t . Therefore, $g \in C^n[a, b]$ and $g^{(n+1)}(x)$ exists. For each $0 \leq k \leq n$, $g(x_k) = f(x_k) - P(x_k) - [f(x) - P(x)]0 = 0$. Moreover, for $t = x$ we have $g(x) = f(x) - P(x) - [f(x) - P(x)] \frac{(x - x_0)(x - x_1) \dots (x - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists and the rest of the expression for $g(t)$ is a polynomial in t . Therefore, $g \in C^n[a, b]$ and $g^{(n+1)}(x)$ exists. For each $0 \leq k \leq n$, $g(x_k) = f(x_k) - P(x_k) - [f(x) - P(x)]0 = 0$. Moreover, for $t = x$ we have $g(x) = f(x) - P(x) - [f(x) - P(x)] \frac{(x - x_0)(x - x_1) \dots (x - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)} = f(x) - P(x) - [f(x) - P(x)](1) = 0$.

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$g \in C^n[a, b]$ and $g^{(n+1)}(x)$ exists. This means for $n+2$ distinct values x, x_0, x_1, \dots, x_n , $g(t) = 0$.

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$g \in C^n[a, b]$ and $g^{(n+1)}(x)$ exists. This means for $n+2$ distinct values x, x_0, x_1, \dots, x_n , $g(t) = 0$. Therefore, by generalized Rolle's theorem, there is a number $\zeta = \zeta(x)$ in $[x_0, x_1, \dots, x_n, x]$ such that $g^{(n+1)}(\zeta) = 0$.

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

Therefore, by generalized Rolle's theorem, there is a number $\zeta = \zeta(x)$ in $[x_0, x_1, \dots, x_n, x]$ such that $g^{(n+1)}(\zeta) = 0$.

$$g^{(n+1)}(t) = f^{(n+1)}(t) - P^{(n+1)}(t) - \frac{(n+1)! [f(x) - P(x)]}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

Therefore, by generalized Rolle's theorem, there is a number $\zeta = \zeta(x)$ in $[x_0, x_1, \dots, x_n, x]$ such that $g^{(n+1)}(\zeta) = 0$.

For any polynomial $Q(x) = a_{n+1}x^{n+1} + a_nx^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$, $Q^{(n+1)}(x) = (n+1)!a_{n+1}$.

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

there is a number $\zeta = \zeta(x)$ in $[x_0, x_1, \dots, x_n, x]$ such that $g^{(n+1)}(\zeta) = 0$.

For any polynomial $Q(x) = a_{n+1}x^{n+1} + a_nx^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$, $Q^{(n+1)}(x) = (n+1)!a_{n+1}$. Therefore

$$\frac{d^{n+1}}{dt^{n+1}}[f(x) - P(x)] \frac{(t-x_0)(t-x_1)\dots(t-x_n)}{(x-x_0)(x-x_1)\dots(x-x_n)} = \frac{(n+1)! [f(x) - P(x)]}{(x-x_0)(x-x_1)\dots(x-x_n)}$$

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

there is a number $\zeta = \zeta(x)$ in $[x_0, x_1, \dots, x_n, x]$ such that $g^{(n+1)}(\zeta) = 0$.

$$0 = g^{(n+1)}(\zeta) = f^{(n+1)}(\zeta) - P^{(n+1)}(\zeta) - \frac{(n+1)! [f(x) - P(x)]}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$$0 = g^{(n+1)}(\zeta) = f^{(n+1)}(\zeta) - P^{(n+1)}(\zeta) - \frac{(n+1)! [f(x) - P(x)]}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$P(x)$ is a polynomial of degree at most n so $\frac{d^{n+1}}{dt^{n+1}} P(t) = 0$.

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof: Now if x is distinct from all x_k , $0 \leq k \leq n$ then we define $g : [a, b] \rightarrow \mathbb{R}$

$$g(t) = f(t) - P(t) - [f(x) - P(x)] \frac{(t - x_0)(t - x_1) \dots (t - x_n)}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

$$0 = f^{(n+1)}(\zeta) - \frac{(n+1)! [f(x) - P(x)]}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

Error Bound for Lagrange Interpolating Polynomials

Error estimate for the Lagrange polynomial: Suppose $f \in C^n[a, b]$ and $f^{(n+1)}(x)$ exists for all $x \in (a, b)$ and $x_0, x_1, x_2, \dots, x_n$ are distinct numbers in $[a, b]$. Then for each $x \in [a, b]$, there is a number $\zeta(x) \in [x_0, x_1, \dots, x_n, x]$ such that

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Proof:

$$0 = f^{(n+1)}(\zeta) - \frac{(n+1)! [f(x) - P(x)]}{(x - x_0)(x - x_1) \dots (x - x_n)}$$

which implies

$$f(x) - P(x) = \frac{f^{(n+1)}(\zeta(x))}{(n+1)!} (x - x_0)(x - x_1) \dots (x - x_n)$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)} = -\frac{576}{595}(x - 1)(x - \frac{16}{9})(x - 4)$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)} = -\frac{576}{595}(x - 1)(x - \frac{16}{9})(x - 4)$$

$$L_3(x) = \frac{(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})}{(4 - 1)(4 - \frac{16}{9})(4 - \frac{9}{4})}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)} = -\frac{576}{595}(x - 1)(x - \frac{16}{9})(x - 4)$$

$$L_3(x) = \frac{(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})}{(4 - 1)(4 - \frac{16}{9})(4 - \frac{9}{4})} = \frac{3}{35}(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)} = -\frac{576}{595}(x - 1)(x - \frac{16}{9})(x - 4)$$

$$L_3(x) = \frac{(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})}{(4 - 1)(4 - \frac{16}{9})(4 - \frac{9}{4})} = \frac{3}{35}(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})$$

$$f(x) = \sqrt{x} \Rightarrow y_0 = \sqrt{x_0} = 1,$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)} = -\frac{576}{595}(x - 1)(x - \frac{16}{9})(x - 4)$$

$$L_3(x) = \frac{(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})}{(4 - 1)(4 - \frac{16}{9})(4 - \frac{9}{4})} = \frac{3}{35}(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})$$

$$f(x) = \sqrt{x} \Rightarrow y_0 = \sqrt{x_0} = 1, y_1 = \sqrt{x_1} = \frac{4}{3},$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)} = -\frac{576}{595}(x - 1)(x - \frac{16}{9})(x - 4)$$

$$L_3(x) = \frac{(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})}{(4 - 1)(4 - \frac{16}{9})(4 - \frac{9}{4})} = \frac{3}{35}(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})$$

$$f(x) = \sqrt{x} \Rightarrow y_0 = \sqrt{x_0} = 1, y_1 = \sqrt{x_1} = \frac{4}{3}, y_2 = \sqrt{x_2} = \frac{3}{2},$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)} = -\frac{576}{595}(x - 1)(x - \frac{16}{9})(x - 4)$$

$$L_3(x) = \frac{(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})}{(4 - 1)(4 - \frac{16}{9})(4 - \frac{9}{4})} = \frac{3}{35}(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})$$

$$f(x) = \sqrt{x} \Rightarrow y_0 = \sqrt{x_0} = 1, y_1 = \sqrt{x_1} = \frac{4}{3}, y_2 = \sqrt{x_2} = \frac{3}{2}, y_3 = \sqrt{x_3} = 2$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)} = -\frac{576}{595}(x - 1)(x - \frac{16}{9})(x - 4)$$

$$L_3(x) = \frac{(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})}{(4 - 1)(4 - \frac{16}{9})(4 - \frac{9}{4})} = \frac{3}{35}(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})$$

$$f(x) = \sqrt{x} \Rightarrow y_0 = \sqrt{x_0} = 1, y_1 = \sqrt{x_1} = \frac{4}{3}, y_2 = \sqrt{x_2} = \frac{3}{2}, y_3 = \sqrt{x_3} = 2$$

$$\text{Therefore, } P(x) = L_0(x) + \frac{4}{3}L_1(x) + \frac{3}{2}L_2(x) + 2L_3(x)$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$L_0(x) = \frac{(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)}{(1 - \frac{16}{9})(1 - \frac{9}{4})(1 - 4)} = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4)$$

$$L_1(x) = \frac{(x - 1)(x - \frac{9}{4})(x - 4)}{(\frac{16}{9} - 1)(\frac{16}{9} - \frac{9}{4})(\frac{16}{9} - 4)} = \frac{729}{595}(x - 1)(x - \frac{9}{4})(x - 4)$$

$$L_2(x) = \frac{(x - 1)(x - \frac{16}{9})(x - 4)}{(\frac{9}{4} - 1)(\frac{9}{4} - \frac{16}{9})(\frac{9}{4} - 4)} = -\frac{576}{595}(x - 1)(x - \frac{16}{9})(x - 4)$$

$$L_3(x) = \frac{(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})}{(4 - 1)(4 - \frac{16}{9})(4 - \frac{9}{4})} = \frac{3}{35}(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})$$

$$\text{Therefore, } P(x) = L_0(x) + \frac{4}{3}L_1(x) + \frac{3}{2}L_2(x) + 2L_3(x)$$

$$P(x) = -\frac{12}{35}(x - \frac{16}{9})(x - \frac{9}{4})(x - 4) + \frac{972}{595}(x - 1)(x - \frac{9}{4})(x - 4) - \frac{864}{595}(x - 1)(x - \frac{16}{9})(x - 4) + \frac{6}{35}(x - 1)(x - \frac{16}{9})(x - \frac{9}{4})$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$\begin{aligned} P(x) &= -\frac{12}{35}\left(x - \frac{16}{9}\right)\left(x - \frac{9}{4}\right)(x - 4) + \frac{972}{595}\left(x - 1\right)\left(x - \frac{9}{4}\right)(x - 4) - \\ &\quad - \frac{864}{595}\left(x - 1\right)\left(x - \frac{16}{9}\right)(x - 4) + \frac{6}{35}\left(x - 1\right)\left(x - \frac{16}{9}\right)\left(x - \frac{9}{4}\right) \\ &= \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85} \end{aligned}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

$$f(x) - P(x) = \frac{f^{(4)}(\zeta)}{4!}(x - x_0)(x - x_1)(x - x_2)(x - x_3)$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

$$f(x) - P(x) = \frac{f^{(4)}(\zeta)}{4!}(x - x_0)(x - x_1)(x - x_2)(x - x_3)$$

$$f(2) - P(2) = \frac{f^{(4)}(\zeta)}{24}(2 - 1)(2 - \frac{16}{9})(2 - \frac{9}{4})(2 - 4) = \frac{f^{(4)}(\zeta)}{216}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

$$f(x) - P(x) = \frac{f^{(4)}(\zeta)}{4!}(x - x_0)(x - x_1)(x - x_2)(x - x_3)$$

$$f(2) - P(2) = \frac{f^{(4)}(\zeta)}{24}(2 - 1)(2 - \frac{16}{9})(2 - \frac{9}{4})(2 - 4) = \frac{f^{(4)}(\zeta)}{216}$$

For some $\zeta \in [1, 4]$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

$$f(2) - P(2) = \frac{f^{(4)}(\zeta)}{24} \left(2 - 1\right) \left(2 - \frac{16}{9}\right) \left(2 - \frac{9}{4}\right) (2 - 4) = \frac{f^{(4)}(\zeta)}{216}$$

For some $\zeta \in [1, 4]$

$$f(x) = \sqrt{x} = x^{\frac{1}{2}} \Rightarrow f'(x) = \frac{1}{2}x^{-\frac{1}{2}}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

$$f(2) - P(2) = \frac{f^{(4)}(\zeta)}{24} \left(2 - 1\right) \left(2 - \frac{16}{9}\right) \left(2 - \frac{9}{4}\right) (2 - 4) = \frac{f^{(4)}(\zeta)}{216}$$

For some $\zeta \in [1, 4]$

$$f(x) = \sqrt{x} = x^{\frac{1}{2}} \Rightarrow f'(x) = \frac{1}{2}x^{-\frac{1}{2}} \Rightarrow f''(x) = -\frac{1}{4}x^{-\frac{3}{2}}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

$$f(2) - P(2) = \frac{f^{(4)}(\zeta)}{24} (2-1)(2-\frac{16}{9})(2-\frac{9}{4})(2-4) = \frac{f^{(4)}(\zeta)}{216}$$

For some $\zeta \in [1, 4]$

$$\begin{aligned} f(x) = \sqrt{x} = x^{\frac{1}{2}} &\Rightarrow f'(x) = \frac{1}{2}x^{-\frac{1}{2}} \Rightarrow f''(x) = -\frac{1}{4}x^{-\frac{3}{2}} \\ &\Rightarrow f^{(3)}(x) = \frac{3}{8}x^{-\frac{5}{2}} \end{aligned}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

$$f(2) - P(2) = \frac{f^{(4)}(\zeta)}{24} (2-1)(2-\frac{16}{9})(2-\frac{9}{4})(2-4) = \frac{f^{(4)}(\zeta)}{216}$$

For some $\zeta \in [1, 4]$

$$\begin{aligned} f(x) = \sqrt{x} = x^{\frac{1}{2}} &\Rightarrow f'(x) = \frac{1}{2}x^{-\frac{1}{2}} \Rightarrow f''(x) = -\frac{1}{4}x^{-\frac{3}{2}} \\ \Rightarrow f^{(3)}(x) &= \frac{3}{8}x^{-\frac{5}{2}} \Rightarrow f^{(4)}(x) = -\frac{15}{16}x^{-\frac{7}{2}} \end{aligned}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

$$f(2) - P(2) = \frac{f^{(4)}(\zeta)}{24}(2-1)(2-\frac{16}{9})(2-\frac{9}{4})(2-4) = \frac{f^{(4)}(\zeta)}{216}$$

For some $\zeta \in [1, 4]$

$$\Rightarrow f^{(4)}(x) = -\frac{15}{16}x^{-\frac{7}{2}} \Rightarrow f(2) - P(2) = \frac{f^{(4)}(\zeta)}{216} = -\frac{5}{1152}\zeta^{-\frac{7}{2}}$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

For some $\zeta \in [1, 4]$

$$\Rightarrow f(2) - P(2) = \frac{f^{(4)}(\zeta)}{216} = -\frac{5}{1152}\zeta^{-\frac{7}{2}}$$

$$-\frac{5}{1152}(1)^{-\frac{7}{2}} < f(2) - P(2) < -\frac{5}{1152}(4)^{-\frac{7}{2}}$$

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

For some $\zeta \in [1, 4]$

$$\Rightarrow f(2) - P(2) = \frac{f^{(4)}(\zeta)}{216} = -\frac{5}{1152}\zeta^{-\frac{7}{2}}$$

$$-\frac{5}{1152}(1)^{-\frac{7}{2}} < f(2) - P(2) < -\frac{5}{1152}(4)^{-\frac{7}{2}}$$

$$-0.004340277778 < f(2) - P(2) < -0.00003390842014$$

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

For some $\zeta \in [1, 4]$

$$\Rightarrow f(2) - P(2) = \frac{f^{(4)}(\zeta)}{216} = -\frac{5}{1152}\zeta^{-\frac{7}{2}}$$

$$-\frac{5}{1152}(1)^{-\frac{7}{2}} < f(2) - P(2) < -\frac{5}{1152}(4)^{-\frac{7}{2}}$$

$$-0.004340277778 < f(2) - P(2) < -0.00003390842014$$

This means $P(2)$ is definitely larger than $f(2) = \sqrt{2}$ and its absolute error is at least 0.00003390842014 and at most 0.004340277778.

Example

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

For some $\zeta \in [1, 4]$

$$-0.004340277778 < f(2) - P(2) < -0.00003390842014$$

This means $P(2)$ is definitely larger than $f(2) = \sqrt{2}$ and its absolute error is at least 0.00003390842014 and at most 0.004340277778.

Notice that $P(2) = \frac{505}{357} \approx 1.414565826$ and

$$f(2) = \sqrt{2} \approx 1.414213562.$$

Example:

- Find the Lagrange interpolating polynomial $P(x)$ for $f(x) = \sqrt{x}$. Let $x_0 = 1$, $x_1 = \frac{16}{9}$, $x_2 = \frac{9}{4}$, and $x_3 = 4$.

$$P(x) = \frac{6}{595}x^3 - \frac{397}{3570}x^2 + \frac{2419}{3570}x + \frac{36}{85}$$

- What are the maximum and minimum absolute errors in using $P(2)$ to estimate $f(2) = \sqrt{2}$?

For some $\zeta \in [1, 4]$

$$-0.004340277778 < f(2) - P(2) < -0.00003390842014$$

This means $P(2)$ is definitely larger than $f(2) = \sqrt{2}$ and its absolute error is at least 0.00003390842014 and at most 0.004340277778.

Notice that $P(2) = \frac{505}{357} \approx 1.414565826$ and

$f(2) = \sqrt{2} \approx 1.414213562$. In particular,

$$f(2) - P(2) \approx -0.000352264.$$